

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

Enquiry into plants and minor works on odoùrs and weather signs

Theophrastus



LELAND · STANFORD JVNIOR · VNIVERSITY


LASSICAL LIBRARY

EDITED BY

. PAGE, LITT.D. W. H. D. ROUSE, LITT.D.

HRASTUS INTO PLANTS

П



THE LOEB CLASSICAL LIBRARY

EDITED BY

E. CAPPS Ph.D., LL.D. T. E. PAGE, LITT.D. W. H. D. ROUSE, LITT.D.

THEOPHRASTUS ENQUIRY INTO PLANTS

 \mathbf{II}

ENQUIRY INTO PLANTS

AND MINOR WORKS ON ODOURS AND WEATHER SIGNS

WITH AN ENGLISH TRANSLATION BY
SIR ARTHUR HORT, BART., M.A.
FORMERLY FELLOW OF TRINITY COLLEGE, CAMBRIDGE

IN TWO VOLUMES



LONDON: WILLIAM HEINEMANN NEW YORK: G. P. PUTNAM'S SONS MOMXVI

234493

Yakali osotakid

BOOK VI

OF UNDER-SHRUBS	
Of the classification of under-shrubs: the wild kinds: the chief distinction that between spinous and spineless	PAGI
Of spineless under-shrubs and their differences	7
Of certain specially important spineless under-shrubs—silphium and magydaris—belonging to ferula-like plants	18
Of spinous under-shrubs and their differences	21
Of cultivated under-shrubs (coronary plants), with which are included those coronary plants which are	35
	ði.
Of the seasons at which coronary plants flower, and of the length of their life	49
BOOK VII	
OF HERBACEOUS PLANTS, OTHER THAN CORONARY PLAN POT-HERBS AND SIMILAR WILD HERBS	its :
Of the times of sowing and of germination of pot-herbs	59
Of the propagation of pot-herbs, and of differences in	
their roots	67
Of the flowers and fruits of pot-herbs	75
Of the various forms of some pot-herbs	81
Of the cultivation of pot-herbs; manure and water	93
Of the pests which infest pot-herbs	95
Of the time for which seed of pot-herbs can be kept	97
Of uncultivated herbs: the wild forms of pot-herbs	99
•	37

Of other uncultivated herbs, which may be classed with pot-herbs	PAGE
Of the differences in stem and leaf found in all herbaceous plants	107
Of other differences seen in herbaceous plants in general,	
as compared with one another and with trees	111
Of the seasons at which herbs grow and flower	115
Of the classes into which herbaceous plants may be divided, as those having a spike and chicory-like	119
plants	-
Of herbs which have fleshy or bulbous roots	125
Of certain properties and habits peculiar to certain herbaceous plants	135
•	
BOOK VIII	
OF HERBACEOUS PLANTS: CEREALS, PULSES, AND 'SUM CROPS'	MER
Of the three classes and the times of sowing and of germination	143
Of differences in the mode of germination and of subsequent development	149
Of differences in development due to soil or climate	155
Of differences between the parts of cereals, pulses, and	
summer crops respectively	159
Of the differences between cereals	165
Of the differences between pulses	173
Of sowing, manuring, and watering	177
Of the degeneration of cereals into darnel	183
Of the peculiar character of chick-pea	183
Of special features of 'summer crops'	185
Of treatment of cereals peculiar to special localities	185
Of cereals which grow a second time from the same	
stock	187
Of the effects of climate, soil, and manuring	189
Of different qualities of seed	191
vi	

Of degeneration of cereals, and of the weeds which infest particular crops
Of the conditions in the seeds of pulses known as 'cookable' and 'uncookable,' and their causes
Of the grains and pulses which most exhaust the soil, or which improve it
Of the diseases of cereals and pulses, and of hurtful winds
Of seeds which keep or do not keep well
Of the age at which seeds should be sown
Of artificial means of preserving seed
Of the effect of heat on seeds
Of certain peculiarities of the seed of lupin and uigilops
BOOK IX
OF THE JUICES OF PLANTS, AND OF THE MEDICINAL PROPERTIES OF HERBS
Of the various kinds of plant-juices and the methods of collecting them
Of resinous trees and the methods of collecting resin and pitch
Of the making of pitch in Macedonia and in Syria
Of frankincense and myrrh: various accounts
Of cinnamon and cassia: various accounts
Of balsam of Mecca
Of other aromatic plants—all oriental, except the iris .
Of the medicinal juices of plants and the collection of
them: general account
Of the cutting of roots for medicinal purposes, and of certain superstitions connected therewith
Of the medicinal uses of divers parts of plants
Of hellebores, the white and the black; their uses and
distribution
Of the various kinds of all-heal
Of the various plants called strykhnos
Of the various kinds of tithymallos
-

	PAGE
Of the two herbs called libanotis	277
Of the two kinds of chamaeleon	277
Of the various plants called 'poppy'	279
Of roots possessing remarkable taste or smell	281
Of the time for which roots can be kept without losing	007
their virtue	287
Of the localities which specially produce medicinal herbs	289
Of the medicinal herbs peculiar to Crete	295
Of wolf's-bane and its habitat, and of meadow-saffron .	299
Of two famous druggists and of the virtues of hemlock.	303
How use diminishes the efficacy of drugs, and how they have not the same effect on all constitutions .	305
Of plants that possess properties affecting lifeless objects	309
Of plants whose properties affect animals other than	
man	309
Of plants possessing properties which affect the mental powers	311
Of plants said to have magical properties	313
A problem as to cause and effect	313
Of certain plants, not yet mentioned, which possess	010
special properties	315
MINOR WORKS	
INTRODUCTION TO THE TREATISES CONCERNING ODOURS AND CONCERNING WEATHER SIGNS	324
CONCERNING ODOURS	
Introductory: Of odours in general and the classifi-	
cation of them	327
Of natural odours; of those of animals and of the effect of odours on animals	329
Of smell and taste	331
Of odours in plants	333
Of artificial odours in general and their manufacture:	
especially of the use of perfumes in wine	333
viii	

	PAGE
Of the oils used as the vehicle of perfumes	341
Of the spices used in making perfumes and their treatment	347
Of the various parts of plants used for perfumes, and of	
the composition of various notable perfumes	351
Of the properties of various spices	355
Of the medicinal properties of certain perfumes	359
Of rules for the mixture of spices, and of the storing of	
various perfumes	36 1
Of the properties of certain perfumes	365
Of other properties and peculiarities of perfumes	373
Of the making of perfume-powders and compound per-	975
fumes	377
Of the characteristic smells of animals, and of certain curious facts as to the smell of animal and vegetable	
products	381
Of odours as compared with other sense-impressions	383
CONCERNING WEATHER SIGNS	
Introductory: general principles	391
The signs of rain	3 97
The signs of wind	407
The signs of fair weather	427
Miscellaneous signs	431
INDEX OF PLANTS	435
KEY TO THE INDEX-	
I.—List of plants mentioned in the Enquiry under botanical names	487
IIList of plants mentioned in the Enquiry unde	
popular names	494

THEOPHRASTUS ENQUIRY INTO PLANTS BOOK VI

VOL, II

ΘΕΟΦΡΑΣΤΟΥ ΠΕΡΙ ΦΥΤΩΝ ΙΣΤΟΡΙΑΣ


 \mathbf{Z}

Ι. Περὶ μὲν οὖν δένδρων καὶ θάμνων εἴρηται πρότερον ἐπόμενον δ' εἰπεῖν περί τε τῶν φρυγανικῶν καὶ ποιωδῶν καὶ εἴ τινες ἐν τούτοις ἔτεραι συμπεριλαμβάνονται φύσεις οἶον ἥ γε σιτηρὰ ποιώδης ἐστί.

Πρώτον δὲ λέγωμεν περὶ τῆς φρυγανικῆς αὕτη γὰρ ἐγγυτέρω τῶν προειρημένων διὰ τὸ ξυλώδης εἶναι. πανταχοῦ μὲν οὖν ἴσως αἰεὶ τὸ ἄγριον τοῦ ἡμέρου πλεῖον, εἰ δὲ μή, περί γε τὴν φρυγανικὴν οὐσίαν ὀλίγον γὰρ τὸ ἤμερον αὐτῆς, ὅπερ σχεδὸν ἐν τοῖς στεφανωτικοῖς ἐστιν, οἶον ῥοδωνία ἰωνία διόσανθος ἀμάρακος ἡμεροκαλλές, ἔτι δὲ ἔρπυλλος σισύμβριον ἐλένιον ἀβρότονον. ἄπαντα γὰρ ταῦτα ξυλώδη καὶ μικρόφυλλα, δι' δ καὶ φρυγανικά. καὶ ἐπὶ τῶν λαχανηρῶν δ'

¹ cf. 6. 6. 1

² This hardly corresponds to the definition of φρύγανα



THEOPHRASTUS ENQUIRY INTO PLANTS

BOOK VI

OF UNDER-SHRUBS.

Of the classification of under-shrubs: the wild kinds: the chief distinction that between spinous and spineless.

I. We have spoken already of trees and shrubs, and next we must speak of under-shrubs and herbaceous plants and of any other natural classes which are included with these; for instance, cereals

come under herbaceous plants.

But first let us tell of under-shrubs, for this class comes near those mentioned above because of its woody character. Now it may be said that with all plants the wild kinds are more abundant than the cultivated, and this is certainly true of the undershrubs. For the cultivated kinds of this class 1 are not numerous, and consist almost entirely of coronary plants, as rose gilliflower carnation sweet marjoram martagon lily, to which may be added tufted thyme bergamot-mint calamint southernwood. For all these are woody and have small leaves; wherefore 2 they are classed as under-shrubs. This class covers

given in 1, 3. 1, nor do some of the plants here mentioned come under the description. St. considers the text defective.

δμοίως, οίον ράφανος πήγανον καὶ ὅσα παραπλήσια τούτοις ἐατίν ὑπὲρ ὧν οὐχ ήττον ἴσως
- ὑρμόττει: κατὰ τὴν οἰκείαν προσηγορίαν εἰπεῖν,
ὅταν περὶ στεφανωμάτων καὶ λαχάνων ποιώμεθα
μνείαν. νῦν δὲ πρῶτον περὶ τῶν ἀγρίων λέγωμεν.
ἔστιν αὐτῶν εἴδη καὶ μέρη πλείω, ἃ δεῖ διαιρεῖν
καὶ τοῖς καθ' ἕκαστον γένος καὶ τοῖς ὅλοις εἴδεσι.

Μεγίστην δ' ἄν τις λάβοι διαφορὰν τῶν ὅλων γενῶν, ὅτι τὰ μὲν ἀνάκανθα τὰ δὲ ἀκανθώδη τυγχάνει. πάλιν δ' ἐν ἑκατέρω τούτων πολλαὶ διαφοραὶ γενῶν καὶ εἰδῶν, ὑπὲρ ὧν καθ' ἑκάτερα

πειρατέον είπεῖν.

Των ἀκανθικών δὴ τὰ μὲν ἀπλως εἰσιν ἄκανθαι, ώσπερ ἀσφάραγος καὶ σκορπίος· οὐ γὰρ ἔχουσι φύλλον οὐδὲν παρὰ τὴν ἄκανθαν. τὰ δὲ φυλλάκανθα, καθάπερ ἄκανος ἤρύγγιον κνῆκος· ταῦτα γὰρ καὶ τὰ τοιαῦτα ἐπὶ τῶν φύλλων ἔχει τὴν ἄκανθαν, δι' ὁ καὶ φυλλάκανθα καλεῖται. τὰ δὲ καὶ παρὰ τὴν ἄκανθαν ἔτερον ἔχει φύλλον, ὥσπερ ἡ ὀνωνὶς καὶ ὁ τρίβολος καὶ ὁ φέως, δν δή τινες καλοῦσι στοιβήν. ὁ δὲ τρίβολος καὶ περικαρπίως δι' ὁ καὶ τοῦτο ἴδιον πρὸς ἄπαντα ὡς εἰπεῖν· ἐπεὶ πτορθάκανθά γε πολλὰ καὶ τῶν δένδρων καὶ τῶν θαμνωδῶν ἐστιν, οἶον ἀχρὰς ῥόα παλίουρος βάτος ῥοδωνία κάππαρις. ἐν μὲν οὖν τοῖς ἀκανθικοῖς ταύτας ἄν τις ὡς εἰπεῖν τύπφ λάβοι τὰς διαφοράς.

¹ ξστιν... εἴδεσι: text needlessly altered by Sch. and W. Sch. himself shews that T. uses εἶδος and γένος almost indiscriminately. Here τῶν ὅλων γενῶν means the same as τοῖς

ENQUIRY INTO PLANTS, VI. 1. 2-3

also pot-herbs, such as cabbage rue and others like them. Of these it is perhaps more appropriate to speak under their proper designation, that is, when we come to make mention of coronary plants and pot-herbs. Now let us first speak of the wild kinds. Of these are several classes and subdivisions, which we must distinguish by the characteristics of each sub-division as well as by those of each class taken as a whole.¹

The most important difference distinguishing class from class which one could find is that between the spineless and the spinous kinds. Again under each of these two heads there are many differences distinguishing kinds and forms, of which we must

endeavour to speak severally.

² Of spinous kinds some just consist of spines, as asparagus and *skorpios*; for these have no leaves except their spines. Then there are the spinous-leaved plants, as thistle eryngo safflower; these and the like have their spines on the leaves, whence their name. Others again have leaves as well as their spines, as rest-harrow caltrop and *pheos*, which some call *stoibe*. Caltrop is also ⁴ spinous-fruited, having spines on the fruit-vessel. Wherefore this peculiarity marks it off from almost all other plants; though many trees and shrubs have spines on the shoots, as wild pear pomegranate Christ's thorn bramble rose caper. Such ⁵ are the general distinctions which may be made among spinous plants.

δλοις είδεσι; and below γενῶν and είδῶν both refer to the smaller divisions called μέρη above. 2 Plin. 21. 91.

⁸ δ φέως δν conj. Sch.; δ φλεώς δ Ald. H.; και δ δή τινες καλ.

στ. P₂. cf. 6. 5. 1 and Index.

⁴ καὶ περικαρπάκανθος conj. Sch.; καὶ ἡ περικαρπία φυλλάκανθον UMVAld. cf. 6. 5. 3. ⁵ οδν add. Sch.

'Εν δε τοις ανακάνθοις ούκ έστιν ούτως διαλαβείν τοίς γένεσιν ή γαρ των φύλλων ανωμαλία μεγέθει καὶ μικρότητι καὶ σχήμασιν ἄπειρος καὶ ἀσαφής ἀλλὰ δεῖ πειρᾶσθαι κατ' ἄλλον τρόπον διαιρείν. πλείω δέ έστι τὰ γένη τὰ τούτων καὶ διαφοράς έγοντα μεγάλας, οίον κίσθος μήλωθρον ερευθεδανών σπειραία κνέωρον δρίγανος θύμβρα σφάκος έλελίσφακος πράσιον κόνυζα μελισσόφυλλον έτερα τοιαθτα πρός τούτοις έτι τὰ ναρθηκώδη καὶ ἐννευρόκαυλα, καθάπερ μάραθον ίππομάραθον ναρθηκία νάρθηξ καὶ τὸ καλούμενον ύπό τινων μυοφόνον καὶ ὅσα ὅμοια τούτοις. άπαντα γάρ ἄν τις καὶ ταῦτα καὶ ὅλως εἴ τι ναρθηκῶδές ἐστι τῆς φρυγανικῆς θείη φύσεως.

ΙΙ. Είδη δὲ καὶ διαφοραὶ καθ' ἔκαστον τῶν είρημένων είσι των μεν φανερώτεραι των δε άφανέστεραι. καὶ γὰρ κίσθου δύο γένη διαιροῦσι, τὸ μὲν ἄρρεν τὸ δὲ θῆλυ, τῷ τὸ μὲν μεῖζον καὶ σκληρότερον καὶ λιπαρώτερον είναι καὶ τὸ ἄνθος έπιπορφυρίζον άμφω δὲ ὅμοια τοῖς ἀγρίοις ρόδοις,

πλην ελάττω και ἄοσμα.

2 Δύο δὲ εἴδη καὶ τοῦ κνεώρου ὁ μὲν γὰρ λευκὸς ό δὲ μέλας. ἔχει δὲ ὁ μὲν λευκὸς τὸ φύλλον

4 θύμβρα σφάκος conj. Sch.; θύμβρας φακός UMVAld.

⁵ cf. 6. 2. 5.

i.e. there is a gradation.

² κίσθος conj. Sch.; κισσός Ald. Η. 3 σπειραία conj. Sch. from Plin. 21. 53; σμηρέα Ald.G.

⁶ ναρθηκώδη = hollow-stemmed, εννευρόκαυλα = plants with a plain unjointed stem, solid with 'immersed' fibres. In the

ENQUIRY INTO PLANTS, VI. I. 4-II. 2

With spineless plants it is not possible to make such 'generic' distinctions; for the variation of the leaves in size and shape is endless, and the differences are not clearly marked 1; but we must try to distinguish on another principle. There are many classes of such plants and they differ widely, as rock-rose 2 bryony madder privet 3 kneoron marjoram savory sphakos 4 (sage) elelisphakos 5 (salvia) horehound konyza balm, and others like these; and in addition to these we have the plants with a ferulalike stem 6 or with a stem composed of fibre, as fennel horse-fennel narthekia (ferula) narthex (ferula) and the plant called by some wolf's bane, s and others like these. All these, as well as any other ferulalike plants, may be placed in the class of undershrubs.

Of spineless under-shrubs and their differences.

II. The various forms and the differences between the above mentioned plants are in some cases more, in some less easy to distinguish. Of rock rose 9 they distinguish two kinds, 'male' and 'female,' in that the one is 10 larger, tougher, more glossy, 11 and has a crimson flower; both however are like the wild rose, 12 save that the flower is smaller and scentless.

There are also two kinds of kneoron, one white, the other black. The white has a leathery oblong

examples given here the two classes are taken together, ναρθηκία being ναρθηκώδης, the others ἐννευρόκαυλα; hence the article is not repeated.

7 cf. 6. 2. 7.

Lit. 'mouse-bane': for other Greek names see Index.
 κίσθου conj. Sch., cf. 6 1.4; κισσοῦ Ald. H.; Plin. 24. 81;

Diosc. 1. 97.

10 είναι conj. W.; έχειν UMVAld. (τό φύλλον έχειν conj. Sch.).

11 i.e. has more glossy leaves.

12 cf. Plin. 21. 55; Theorr. 5. 131. See Index.

δερματώδες πρόμηκες όμοιόσχημον τρόπον τινὰ τῆ ἐλάα, ὁ δὲ μέλας οἰον ἡ μυρίκη σαρκώδες ἐπίγειος δὲ μᾶλλον ὁ λευκός. ἔστι δὲ ὀσμώδης, ὁ δὲ μέλας ἄοσμος. τὴν δὲ ῥίζαν τὴν εἰς βάθος ἄμφω μεγάλην ἔχουσι καὶ τοὺς ἀκρεμόνας πολλοὺς καὶ παχεῖς καὶ ξυλώδεις ἀπ' αὐτῆς τῆς γῆς ἡ μικρὸν ἄνω σχιζομένους, ξυλωδεστάτην δέ. γλίσχρον δὲ σφόδρα, δι' δ καὶ χρῶνται πρὸς τὸ καταδεῖν καὶ περιλαμβάνειν, ὥσπερ τῷ οἴσῳ. βλαστάνει δὲ καὶ ἀνθεῖ μετ' ἰσημερίαν μετοπωρινὴν καὶ ἀνθεῖ πολὺν χρόνον.

Καὶ τῆς ὀριγάνου δὲ ἡ μέλαινα ἄκαρπος ἡ δὲ λευκὴ κάρπιμος. καὶ θύμον τὸ μὲν λευκὸν τὸ δὲ μέλαν εὐανθὲς δὲ σφόδρα: περὶ τροπὰς γὰρ ἀνθεῖ θερινάς. ἀφ' οὖ καὶ ἡ μέλιττα λαμβάνει τὸ μέλι, καὶ τούτω φασὶν οἱ μελιττουργοὶ δῆλον εἶναι πότερον εὐμελιτοῦσι ἡ οὖ· καλῶς γὰρ ἀπανθήσαντος εὐμελιτεῖν· βλάπτει δὲ καὶ ἀπόλλυσι

την άνθησιν έαν ύδωρ έπιγένηται.

Σπέρμα δὲ κάρπίμον ἡ μὲν θύμβρα καὶ ἔτι μᾶλλον ἡ ὀρίγανος ἔχει φανερόν, τοῦ θύμου δ' οὐκ ἔστι λαβεῖν, ἀλλ' ἐν τῷ ἄνθει πως ἀναμέμικται· σπείρουσι γὰρ τοῦτο καὶ ἀναβλαστάνει. 4 ζητοῦσι δὲ καὶ λαμβάνουσιν οἱ ἐξάγειν 'Αθήνησι βουλόμενοι τὸ γένος. ἴδιον δὲ ἔχει καὶ πρὸς ταῦτα καὶ σχεδὸν πρὸς τὰ ἄλλα τὸ κατὰ τοὺς τόπους· οὐ γάρ φασι δύνασθαι φύεσθαι καὶ

¹ cf. 1. 10. 4.

3 περιλαμβάνει conj. W. from G; περιλαμβάνειν Ald.

⁴ Plin. 21. 55.

² Apparently an afterthought, suggested by the mention of the woodiness of the branches.

ENQUIRY INTO PLANTS, VI. 11. 2-4

leaf, somewhat like that of the olive; the leaf of the black is like that of the tamarisk 1 and fleshy; the white grows more on the ground and is scented, while the black is scentless. In both the root, which runs deep, is large (and the branches which divide at the ground level are numerous thick and woody), and the root is also very woody. 2 It is also very tough, wherefore it is used for binding and to put 3 round things, like the withy. It grows and flowers after the autumnal equinox, and remains in flower a long time.

⁴Of marjoram the black form is barren, the white bears fruit. ^b There is a black and a white thyme, and it flowers very freely: it is in bloom about the summer solstice. It is from this flower that the bee gets the honey, and by it ⁶ beekeepers say that it is made known whether they have a good yield of honey or not; for, if the thyme flowers abundantly, ⁷ they have a good yield, but the bloom is injured or

even destroyed if it is rained upon.

Savory, and still more marjoram, has a conspicuous fruitful seed, but in thyme it is not easy to find, being somehow mixed up with the flower; for men sow the flower and plants come up from it. 8 This plant is sought and obtained by those in Athens who wish to export such herbs. But it has a peculiarity as compared both with similar plants and with most others, namely the kind of region which it affects 9; they say that it can not be grown or become

⁵ Plin. 21. 56 and 154.

⁶ τούτφ conj. Sch.; τοῦτο Ald.

⁷ καλώς mBod.; άλλως UMVAld. H.

⁸ Plin. 21. 57.

⁹ το κατά τοὺς τόπους conj. W.; και κατά τοὺς τ. Ald.; και κατά τόπους P.

λαμβάνειν ὅπου μὴ ἀναπνοὴ διῖκνεῖται ἡ ἀπὸ τῆς θαλάττης δι' ὁ οὐδ' ἐν ᾿Αρκαδίᾳ γίνεται · θύμβρα δὲ καὶ ὀρίγανος καὶ τὰ τοιαῦτα πολλὰ καὶ πολλαχοῦ. παραπλήσιον οὖν τὸ συμβαῖνον τοῦτο καὶ ἐπὶ τῆς ἐλάας · οὐδὲ γὰρ οὐδ' ἐκείνη δοκεῖ τριακοσίων σταδίων ἀπὸ θαλάττης ἐπάνω φύεσθαι.

Σφάκος δὲ καὶ ἐλελίσφακος διαφέρουσιν ώσὰν τὸ μὲν ἥμερον τὸ δὲ ἄγριον λειότερον γὰρ τὸ φύλλον τοῦ σφάκου καὶ ἔλαττον καὶ αὐχμηρότε-

ρον, τὸ δὲ τοῦ ἐλελισφάκου τραχύτερον.

Δύο δὲ γένη καὶ τοῦ πρασίου το μὲν γὰρ ἔχει ποῶδες το φύλλον καὶ μᾶλλον ἐπικεχαραγμένου, ἔτι δὲ τὰς ἐντομὰς ἐνδήλους σφόδρα καὶ βαθείας, ῷ καὶ οἱ φαρμακοπῶλαι χρῶνται πρὸς ἔνια· τὸ δὲ ἔτερον στρογγυλότερον καὶ αὐχμῶδες σφόδρα, καθάπερ τοῦ σφάκου, καὶ τὰς ἐντομὰς ἀμαυρο-

τέρας έχου καὶ ἐπικεχαραγμένου ήττου.

Κονύζης δε το μεν άρρεν το δε θήλυ. διαφορας δε έχει καθάπερ τα άλλα τα ούτω διαιρούμενα το μεν γαρ θήλυ λεπτοφυλλότερον και
ξυνεστηκός μαλλον και το δλον έλαττον, το
δε άρρεν μειζόν τε και παχυκαυλότερον και
πολυκλωνότερον και το φύλλον μειζον και
λιπαρώτερον έχον, έτι δε το άνθος λαμπρότερον. καρποφόρα δε άμφω το δε όλον όψιβλαστει και όψιανθει περι 'Αρκτούρον και μετ'

² σφάκοs conj. Sch.: σφάκελος UMVP₂Ald.; Plin. 22, 146

¹ λαμβάνειν P₂Ald.: lit. 'take hold,' cf. 6. 2. 6; βλαστάνειν conj. W.

ENQUIRY INTO PLANTS, VI. 11. 4-6

established where a breeze from the sea does not reach. This is why it does not grow in Arcadia, while savory marjoram and such plants are common in many parts. (A similar peculiarity is found in the olive; for it appears that it likewise will not grow more than three hundred furlongs from the sea.)

The difference between sphakos² (sage) and elelisphakos (salvia) is like that between cultivated and wild; for the leaf of sphakos³ is smoother smaller and less succulent,⁴ while that of elelisphakos is

rougher.5

There are also two kinds of horehound: one has a narrow leaf with a more jagged edge, and the notches are very conspicuous and deep, and this is the plant used by druggists for certain purposes; the other has a rounder 6 leaf, which, like that of *sphakos*, is not at all succulent; the notches are less conspicuous and

the edge less jagged.

Of konyza, there is a 'male' and a 'female' kind, the differences between them being such as are usual between forms so distinguished; the 'female' has slenderer leaves, is more compact, and a smaller plant; the 'male' is larger, has thicker stalks, is more branched, has larger glossier leaves, and moreover the flower is more conspicuous. Both bear fruit; the plant as a whole is late in growing and in blooming; it blooms about the rising of

4 W. omits ήττον before αὐχμηρότερον.
5 τραχύτερον conj. Scal. from G; βραχύτερον Ald. H.

6 στρογγυλότερον: cf. 1. 10. 4 n.

³ σφάκου conj. Sch.; σφακέλου UMVP2Ald.

⁷ See Index. Plin. 26. 58. cf. Nic. Ther. 875; Diosc. 3. 121.

'Αρκτοῦρον λαμβάνει. βαρεῖα δὲ ἡ ὀσμὴ τοῦ ἄρρενος, ἡ δὲ τῆς θηλείας δριμυτέρα, δι' ὃ καὶ πρὸς τὰ θηρία χρησίμη.

Ταθτα μεν οθν και τὰ τοιαθτα ὅσπερ διαφέροντα. πάλιν δε ἄλλα μονοειδη τυγχάνοντα και των πρότερον είρημένων και ετερα παρὰ ταθτα·

πλείω γάρ ἐστι.

Τὸ δὲ ναρθηκῶδες, καὶ γὰρ καὶ τοῦτο τῶν φρυγανικών, πολλάς περιείληφεν ίδέας έν οίς πρώτον ύπερ του κοινού πασι λεκτέον, ύπερ νάρθηκός τε καὶ ναρθηκίας, εἴτε τὸ αὐτὸ γένος ἐστὶν ἀμφοῖν διαφέρον δὲ κατὰ μέγεθος, εἴτε καὶ ετερον ωσπερ τινές φασιν. ή δ' οὖν φανερὰ φύσις ἀμφοῖν ὁμοία πλην κατὰ τὸ μέγεθος ὁ μὲν γὰρ νάρθηξ γίνεται μέγας σφόδρα ή δὲ ναρθηκία μικρά. μονόκαυλα δ' ἄμφω καὶ γονατώδη, ἀφ' ὧν τά τε φύλλα βλαστάνει καὶ καυλοί 8 τινες μικροί. βλαστάνει δὲ παραλλάξ τὰ φύλλα. λέγω δὲ παραλλάξ ὅτι οὐκ ἐκ τοῦ αὐτοῦ μέρους τῶν γονάτων ἀλλ' ἐναλλάξι περιειληφότα δὲ τὸν καυλὸν ἐπὶ πολύ, καθάπερ τὰ τοῦ καλάμου, πλὴν άποκεκλιμένα ταθτα μάλλον διὰ τὴν μαλακότητα καὶ τὸ μέγεθος μέγα γὰρ τὸ Φύλλον καὶ μαλακόν καὶ πολυσχιδές, ὅστε εἶναι σχεδόν τριχῶδες· ἔχει δὲ μέγιστα τὰ κάτω πρὸς τὴν γῆν καὶ ἀεὶ κατὰ λόγον. ἄνθος δὲ μηλινοειδὲς άμαυρόν, καρπὸν δὲ παρόμοιον τῷ ἀνήθω πλην

¹ λαμβάνει Ald.; ἀδρύνει conj. W. But cf. the somewhat similar use 6. 2. 4.

² contra bestiarum morsus Plin. l.c.

³ Plin. 13, 132 and 133.

⁴ The form of expression in the repeated ὑπὲρ seems loose,

ENQUIRY INTO PLANTS, VI. 11. 6-8

Arcturus and is full grown 1 after his setting. The smell of the 'male' plant is strong, but that of the 'female' more pungent; wherefore both of them are of use against wild beasts.²

These plants then and others like them have, as it were, different forms. Again there are some which have but one form both among those already mentioned and others as well; for there are numerous

plants of this class.

3 The class of ferula-like plants (for this too belongs to the under-shrubs) comprises many kinds: here we must first speak of the characteristic which is common to all, including ferula itself 4 (narthex) and narthekia, whether they both belong to the same kind and differ only in size, or whether, as some sav. they are distinct. The obvious character of both is alike, except as to size; for narthex grows very tall, while narthekia is a small plant. Each of them has a single stalk, which is jointed; from this spring the leaves and some small stalks; the leaves come alternately-by which I mean that they do not spring from the same part of the joint, but in alternating rows. For a considerable distance they embrace the stalk, like the leaves of the reed, but they turn back from it more owing to their softness and their size; for the leaf is large soft and much divided, so that it is almost hair-like; the largest leaves are the lowest ones next the ground, and so on in proportion. The flower is quince-yellow 5 and inconspicuous, the fruit 6 like dill, but larger.7

and above $\ell\nu$ of s is hardly satisfactory. Sch. suspects corruption.

5 μηλινοειδès: cf. 7. 3. 1.

7 μείζω conj. Sch.; μείζον Ald.

⁶ άμαυρόν, καρπόν δέ conj. Sch.; άμαυρόκαρπον Ald.

μείζω. ἐξ ἄκρου δὲ σχίζεται καὶ ἔχει τινὰς οὐ μεγάλους καυλούς ἐνταῦθα δὲ τό τε ἄνθος καὶ ὁ καρπός. ἔχει δὲ καὶ ἄνθος καὶ καρπὸν καὶ ἐν τοῦς παρακαυλίζουσι δι' ὅλου, καθάπερ τὸ ἄνηθον. ἐπετειόκαυλον δέ, καὶ ἡ βλάστησις τοῦ ἡρος πρῶτον μὲν τῶν φύλλων ἔπειτα τοῦ καυλοῦ, καθάπερ τῶν ἄλλων. ῥίζαν δὲ ἔχει βαθεῖαν, ἔστι δὲ μονόρριζον. ὁ μὲν οὖν νάρθηξ τοιοῦτος.

Τῶν δ' ἄλλων τὰ μὲν ὁμοιότερα τούτω τὸν καυλὸν ἔχει <κοῖλον>, καθάπερ ὁ μανδραγόρας καὶ τὸ κώνειον καὶ ὁ ἐλλέβορος καὶ ὁ ἀνθέρικος· τὰ δ' οἶον ἐννευρόκαυλα τυγχάνει, καθάπερ μάραθον μυοφόνον τὰ ὅμοια τούτοις. ἴδιος δὲ ὁ καρπὸς τοῦ μανδραγόρου τῷ μέλας τε καὶ ῥαγώδης καὶ οἰνώδης εἶναι τῷ χυμῷ.

III. Μέγισται δὲ καὶ ἰδιώταται φύσεις ἥ τε τοῦ σιλφίου καὶ ἡ τοῦ παπύρου ἐν Αἰγύπτω ναρθηκώδη γὰρ καὶ ταῦτά ἐστιν ὑπὲρ ὧν τοῦ μὲν παπύρου πρότερον εἴπομεν ἐν τοῦς ἐνύδροις,

ύπερ θατέρου δε νῦν λεκτέον.

Τὸ δὲ σίλφιον ἔχει ρίζαν μὲν πολλὴν καὶ παχεῖαν, τὸν δὲ καυλὸν ἡλίκον νάρθηξ, σχεδὸν δὲ καὶ τῷ πάχει παραπλήσιον, τὸ δὲ φύλλον, δ καλοῦσι μάσπετον, ὅμοιον τῷ σελίνῳ· σπέρμα

² See Index: the stalk is specially in question here.

⁴ Plin. 25. 147-150 describes mandragoras, but his description is not taken from T. cf. Diosc. 4. 75, where three kinds

¹ κοιλον add. W.

³ cf. 6. 1. 4 n. ἐκνευρόκαυλα P₂Ald., cf. ἔκλευκος, 'whitish'; ἐννευρόκαυλα conj. Sch. as in 6. 1. 4; but οἶον indicates the coinage of a fresh term. κώνειον seems to be placed in the wrong list.

ENQUIRY INTO PLANTS, VI. 11. 8-111. 1

plant divides at the top and has some small branches, on which grow the flower and the fruit. It also bears flowers and fruit on the side-stalks all the way up, like dill. The stalk only lasts a year, and the growth takes place in spring, the leaves growing first and then the stem, as with other plants. It roots deep and has but a single root. Such is the ferula.

Of the others some to a certain extent resemble ferula, that is, in having a hollow stem 1; for instance deadly nightshade hemlock hellebore asphodel 2: while some have a stem more or less, as it were, consisting of fibre, 3 as fennel aconite and others like these. The fruit of deadly nightshade 4 is peculiar in being black and like a grape and like wine in taste.

Of certain specially important spineless under-shrubs—silphium and magydaris—belonging to ferula-like plants.

III. Most important and peculiar in their characters are the silphium and papyrus of Egypt. These too come under the class of ferula-like plants; of these we have spoken ⁵ of the papyrus already under the head of plants living in water; of the other we have now to speak.

⁶The silphium has a great deal of thick root; its stalk is like ferula in size, and is nearly as thick; the leaf, which they call maspeton, is like celery: it has a broad fruit, which is leaf-like,

of $\mu a \nu \delta \rho a \gamma \delta \rho a s$ are described: there being only two known species of mandragora, the third may be atropa Belladonna; and to this plant may also refer an interpolated sentence in Diosc. 4. 73 ($\delta \nu \delta o s \ldots \sigma \tau a \phi \nu \lambda \delta \nu$).

⁶ 4. 8. 3 and 4. Papyrus is loosely classed with ferula-like plants, as it has not a hollow stem.

⁶ Plin, 19, 42-45.

Digitized by Google

δ' έχει πλατύ, οίον φυλλώδες, τὸ λεγόμενον φύλλον. ἐπετειόκαυλον δ' ἐστίν, ὥσπερ ὁ νάρθηξ. αμα μεν οθν τω ήρι το μάσπετον τοθτο άφίησιν, δ καθαίρει τὰ πρόβατα καὶ παχύνει σφόδρα καὶ τὰ κρέα θαυμαστὰ ποιεῖ τῆ ἡδονῆ· μετὰ δὲ ταῦτα καυλόν, <ον> ἐσθίεσθαι πάντα τρόπον ἐφθὸν οπτόν, καθαίρειν δε και τουτόν φασι τα σώματα 2 τετταράκοντα ήμέραις. όπον δε διττον έχει, τον μεν ἀπὸ τοῦ καυλοῦ τὸν δε ἀπὸ τῆς ρίζης, δι δ καλοῦσι τὸν μὲν καυλίαν τὸν δὲ ῥιζίαν. ἡ δὲ ρίζα του Φλοιον έγει μέλανα, καὶ τοῦτον περιαιροῦσιν. ἔστι δὲ ώσπερ μέταλλα τῶν ριζοτομιῶν αὐτοῖς, ἐξ ὧν ὁπόσον ἂν δοκῆ συμφέρειν ταμιευόμενοι πρός τὰς τομὰς καὶ τὸ προϋπάργον τέμνουσιν οὐκ ἔξεστι γὰρ οὔτε παρατέμνειν ούτε πλείον των τεταγμένων και γαρ διαφθείρεται καὶ σήπεται τὸ ἀργὸν ἐὰν χρονίζη. κατεργάζονται δε άγοντες είς τον Πειραιά τόνδε τον τρόπου δταν βάλωσι είς άγγεία καὶ ἄλευρα μίξωσι, σείουσι χρόνον συχνόν, δθεν καὶ τὸ γρώμα λαμβάνει και έργασθέν ἄσηπτον ήδη διατὰ μὲν οὖν κατὰ τὴν ἐργασίαν καὶ τομὴν μένει. ούτως έγει.

Τόπου δὲ πολὺν ἐπέχει τῆς Λιβύης πλείω γάρ φασιν ἢ τετρακισχίλια στάδια πλείστα δὲ γίνεσθαι περὶ τὴν σύρτιν ἀπὸ τῶν Εὐεσπερίδων. ἴδιον δὲ τὸ φεύγειν τὴν ἐργαζομένην καὶ ἀεὶ συνεργαζομένης καὶ συνημερουμένης ἐξαναχωρεῖν,

4 cf. 9. 1. 7; Diosc. 3. 80.

¹ οΐον φυλλ. τό λεγ. conj. W.; φυλλ. οΐον τό λεγ. Ald. H.

I have added δν.
 μέταλλα U; μέτρα Ald.; ? ἐν μετάλλοις μέτρα.

ENQUIRY INTO PLANTS, VI. III. 1-3

as it were, and is called the phyllon. The stalk lasts only a year, like that of ferula. spring it sends up this maspeton, which purges sheep and greatly fattens them, and makes their flesh wonderfully delicious; after that it sends up a stalk. which 2 is eaten, it is said, in all ways, boiled and roast, and this too, they say, purges the body in forty days. It has two kinds of juice, one from the stalk and one from the root; wherefore the one is called 'stalk-juice,' the other 'root-juice.' root has a black bark, which is stripped off. have regulations, like those in use in mines.3 for cutting the root, in accordance with which they fix carefully the proper amount to be cut, having regard to previous cuttings and the supply of the plant. For it is not allowed to cut it wrong nor to cut more than the appointed amount; for, if the juice is kept and not used, it goes bad and decays. When they are conveying it to Peiraeus, they deal with it thus 4:having put it in vessels and mixed meal with it, they shake it for a considerable time, and from this process it gets its colour, and this treatment 5 makes it thenceforward keep without decaying. are the facts in regard to the cutting and treatment.

The plant is found over a wide tract of Libya, for a distance,⁶ they say, of more than four thousand furlongs, but it is most abundant⁷ near the Syrtis, starting from the Euesperides islands. It is a peculiarity of it that it avoids cultivated ground, and, as the land is brought under cultivation and tamed,

17

δ ἐργασθὲν: ἐξοργασθὲν conj. Salm.; from Plin. l.c., argumentum erat maturitatis color siccitasque sudore finito.

⁶ cf. Strabo 2. 5. 20; 17. 3. 20: Scyl. Periplus, Libya.
⁷ πλείστα conj. W.; πλείονα U; τὰ πλείονα MAld.; γίνεσθαι conj. W.; γενέσθαι Ald.

ώς οὐ δεομένου δῆλον ὅτι θεραπείας ἀλλ' ὅντος ἀγρίου. φασὶ δ' οἱ Κυρηναῖοι φανῆναι τὸ σίλφιον ἔτεσι πρότερον ἡ αὐτοὶ τὴν πόλιν ὤκησαν ἐπτά· οἰκοῦσι δὲ μάλιστα περὶ τριακόσια εἰς

Σιμωνίδην ἄρχοντα 'Αθήνησιν.

Οἱ μὲν οὖν οὕτω λέγουσιν. οἱ δὲ τοῦ σιλφίου τὴν ῥίζαν φασὶ γίνεσθαι πηχυαίαν ἢ μικρῷ μείζω. ταύτην δὲ ἔχειν ἐπὶ τοῦ μέσου κεφαλήν, δ καὶ μετεωρότατόν ἐστι καὶ σχεδὸν ὑπὲρ γῆς, καλεῖσθαι δὲ γάλα ἐξ ῆς δὴ φύεσθαι μετὰ ταῦτα καὶ τὸν καυλόν, ἐκ δὲ τούτου μαγύδαριν τὸ καὶ καλούμενον φύλλον τοῦτο δ' εἶναι σπέρμα καὶ ὅταν νότος λαμπρὸς πνεύση μετὰ Κύνα διαρρίπτεσθαι, ἐξ οὖ φύεσθαι τὸ σίλφιον. τῷ αὐτῷ δὲ ἔτει τήν τε ῥίζαν γίνεσθαι καὶ τὸν καυλόν οὐθὲν δὲ τοῦτο ἴδιον, καὶ γὰρ ἐπ' ἄλλων, εἰ μὴ τοῦτο λέγουσιν ὅτι εὐθὺς φύεται μετὰ τὴν διάρριψιν.

Καὶ τοῦτο ἴδιον καὶ διάφορον τοῖς πρότερον, ὅτι φασὶ δεῖν ὀρύττειν ἐπέτειον· ἐὰν δὲ ἐαθῆ, φέρειν μὲν τὸ σπέρμα καὶ τὸν καυλόν, χείρω δὲ γίνεσθαι καὶ ταῦτα καὶ τὴν ρίζαν, ὀρυττομένας δὲ βελτίους γίνεσθαι διὰ τὸ μεταβάλλεσθαι τὴν γῆν. ἐναντίον δὲ τοῦτο τῷ φεύγειν τὴν ἐργάσιμον. ἐσθίεσθαι καὶ τὰς ρίζας προσφάτους κατατεμνομένας εἰς ὄξος. τὸ δὲ φύλλον τῆ χροιῷ χρυσοειδὲς

¹ cf. Hdt. 4. 158. ² B.C. 310.

7 τοῦτο conj. Salm.; τούτον UMVAld.; τοῦτον P2.

³ ταύτην δὲ... τὸ σίλφιον: text as restored conjecturally by W., chiefly by alteration in the order of the words in Ald.
⁴ καλεῖσθαι δὲ γάλα after φύεσθαι in Ald.

⁶ In 6. 3. 7 this name is applied to a distinct plant. μαγ. τὸ καὶ conj. Salm.; μαγ. καὶ τὸ P₂Ald. H. ⁶ cf. 6. 3. 2.

ENQUIRY INTO PLANTS, VI. 111. 3-5

it retires, plainly shewing that it needs no tendance but is a wild thing. The people of Cyrene say that the silphium appeared ¹ seven years before they founded their city; now they had lived there for about three hundred years before the archonship at Athens of Simonides.²

Such is their account. Others however say that the root of the silphium grows to the length of a cubit or a little longer, and in the middle of this is a head,³ which is the highest part and almost comes above ground, and is called the 'milk',⁴ from this then presently grows the stalk, and from that the magydaris,⁵ which is also called the phyllon ⁶; but it ⁷ is really the seed, and, when a strong south wind blows after the setting of the dog-star, it is scattered ⁸ abroad and the silphium grows from it. The root and the stalk grow in the same year; nor is this a singular feature—unless they mean that it grows immediately after the dispersal ⁹ of the seed—since the same thing occurs with other ¹⁰ plants also.

There is this singular statement, which is inconsistent with what was said above, that, it is said, it is necessary to dig the ground every year, and that, if it be left alone, it bears 11 the seed and the stalk, but these are inferior and so is the root; on the other hand, that with digging they are improved because the soil is changed. (This is inconsistent with the statement that silphium avoids cultivated land.) They add that the roots are cut up into vinegar and eaten fresh, and that the leaf is of a golden

c 2

 $^{^{8}}$ διαρρίπτεσθαι conj. Sch.; διαρρίπτεται Ald.; διαρριπτείται U; διαρρίπτηται M.

ο διάρριψιν conj. Sch.; δίριψιν UM; έκριψιν Ald.

 ¹⁰ ἐπ' ἄλλων conj. W.; τῶν ἄλλων Ald. H.
 ¹¹ μἐν conj. Sch.; πᾶν Ald.

- 8 εἶναι. ἐναντίον δὲ καὶ τὸ μὴ καθαίρεσθαι τὰ πρόβατα τὸ φύλλον ἐσθίοντα· φασὶ γὰρ καὶ τοῦ ἤρος καὶ τοῦ χειμῶνος εἰς ὅρος ἀφιέναι, νέμεσθαι δὲ τοῦτό τε καὶ ἔτερον ὅμοιον ἀβροτόνῳ· θερμαντικὰ δ' ἄμφω δοκεῖ εἶναι καὶ κάθαρσιν μὲν οὐ ποιεῖν, ἀναξηραίνειν δὲ καὶ συμπέττειν· ἐὰν δέ τι νοσοῦν ἡ κακῶς ἔχον εἰσέλθη πρόβατον, ὑγιάζεσθαι ταχέως ἡ ἀποθνήσκειν, ὡς δ' ἐπὶ τὸ πολὺ σώζεσθαι μᾶλλον. ταῦτα μὲν ὁποτέρως ἔχει σκεπτέον.
 - 'Η δὲ καλουμένη μαγύδαρις ἔτερόν ἐστι τοῦ σιλφίου μανότερόν τε καὶ ἦττον δριμὰ καὶ τὸν ὀπὸν οἰκ ἔχει διάδηλος δέ ἐστι καὶ τἢ ὄψει τοῖς ἐμπείροις. γίνεται δὲ περὶ Συρίαν καὶ οἰκ ἐν Κυρήνη· φασὶ δὲ καὶ ἐν τῷ Παρνασίῳ ὅρει πολλήν· ἔνιοι δὲ σίλφιον τοῦτο καλοῦσιν. εἰ δὲ φεύγει τὴν ἐργάσιμον ὥσπερ τὸ σίλφιον σκεπτέον, ὡσαύτως δὲ καὶ εἴ τι ὅμοιον ἢ παραπλήσιον ἔχει φύλλου τε πέρι καὶ καυλοῦ, καὶ εἰ ὅλως ἀφίησί τι δάκρυον. τὴν μὲν οῦν ναρθηκώδη [καὶ ὅλως τὴν ἀκανθώδη] φύσιν ἐν τοῖς τοιούτοις θεωρητέον.

IV. Τῆς δ' ἀκανθικῆς, ἐπόμενον γὰρ τοῦτο εἰπεῖν, ἐπειδὴ διήρηται τὸ μὲν ἀκανθῶδες ὅλως τὸ δὲ φυλλάκανθον, ὑπὲρ ἐκατέρου χώρις λεκτέον,

¹ cf. Arr. Anab. 3. 28. 6 and 7.

² Artemisia camphorata: Index App. (24). ⁸ Plin. 19. 46; Diosc. 3. 94; Hesych. s.v.; Photius, Gloss. s.v.; cf. 6. 3. 4 n.

ENQUIRY INTO PLANTS, VI. III. 5-IV. 1

colour. We have also the inconsistent statement that sheep are not purged by eating the leaves; for they say that in spring and in winter they are driven into the hill-country, where 1 they feed on this and on another plant 2 which is like southernwood; both these plants appear to be heating and not to cause purging, but, on the contrary, to have a drying effect and promote digestion. It is also said that, if a sheep which is sick or in bad condition comes to that district, it is quickly cured or else dies, but usually it recovers. Which of these accounts is true is matter for enquiry.

The plant called magydaris is distinct from silphium, being of later growth and less pungent, and it does not produce the characteristic juice; experts can also easily distinguish it by its appearance. It grows in Syria and not in Cyrene, and they say that it is also abundant on Mount Parnassus, and some call it silphium. Whether however, like silphium, it avoids cultivated ground is matter for enquiry, as also whether it has any resemblance or likeness in leaf and stalk, and, in general, whether it produces a juice. In these examples we may consider the class of ferula-like plants [and, in general, that of spinous plants.⁴]

Of spinous under-shrubs and their differences.

IV. Taking next the class of spinous plants (for we must next speak of them), we have already distinguished 5 those which are altogether spinous and those which have spinous leaves, and now we must

 $^{^4}$ καl... ἀκανθώδη. These words occur only in U: they cannot belong here. Note that τ δ μὲν ἀκανθώδες δλως occurs just below. 5 6. 1. 3.

καὶ τρίτου δὴ περὶ τοῦ καὶ παρὰ τὴν ἄκανθαν έχοντος φύλλον, ώσπερ ο τε φέως και ο τρίβολος. καὶ ἡ κάππαρις ἴδιον ἔχει τὸ μὴ μόνον τὴν ἐκ τῶν καυλῶν ἄκανθαν ἔχειν ἀλλὰ καὶ τὸ φύλλον έπακανθίζον. των δε διηρημένων είδων πλείστον μέν έστι τὸ φυλλάκανθον, ἐλάχιστον δὲ ώς εἰπεῖν τὸ ἀκανθῶδες ὅλως. βραχὺ γάρ τι πάμπαν ἐστίν, ωσπερ ελέχθη, καὶ σχεδον οὐ ράδιον λαβεῖν παρά

τε τὸν ἀσφάραγον καὶ τὸν σκορπίον:

2 'Αμφότερα δὲ ταῦτα ἀνθεῖ μετὰ ἰσημερίαν φθινοπωρινήν. ὁ μὲν σκορπίος ἐν τῷ σαρκώδει τῷ ἐποιδοῦντι τῷ ὑπὸ τὸ ἄκρον τῆς ἀκάνθης έχων τὸ ἄνθος έξ ἀρχῆς μὲν λευκὸν ὕστερον δ' έπιπορφυρίζου. ὁ δὲ ἀσφάραγος ἐκφύων παρὰ τὰς ἀκάνθας κορυνῶδες μικρόν, ἐκ τούτου δέ ἐστι τὸ ἄνθος μικρόν. ὁ δὲ σκορπίος μονόρριζον καὶ βαθύρριζον, ὁ δὲ ἀσφάραγος βαθύρριζόν τε εὖ μάλα καὶ πολύρριζον πυκναῖς ταῖς ρίζαις, ώστε τὸ ἄνω συνεχὲς είναι αὐτῶν, ἀφ' οὐ καὶ αί βλαστήσεις αὐτῶν τῶν καυλῶν ἀναβλαστάνει δὲ ὁ καυλὸς ἐκ τῆς ἀσφαραγιᾶς τοῦ ἦρος καὶ έδωδιμός έστιν είθ' ούτως αποτραχύνεται καὶ έξακανθούται προϊούσης της ώρας ή δε άνθησις ούκ έκ τούτου μόνον άλλα και έκ των πρότερον. οὐ γὰρ ἐπετειόκαυλόν ἐστι. τὰ μὲν οὖν ὅλως άκανθώδη τοιαύτην τινά έχει φύσιν.

Των δε φυλλακάνθων το πλείστον γένος ώς

¹ φέως conj. St.; φλεώς Ald. cf. 6. 1. 3. ² cf. 6. 1. 3.

³ Plin. 21. 91; 22. 39.

ENQUIRY INTO PLANTS, VI. IV. 1-3

speak of each of these classes separately, and also, in the third place, of those which have leaves as well as their spines, such as pheos 1 and caltrop. Moreover caper has the peculiarity of possessing not only spines on its stems but also a spinous leaf. Of the classes thus distinguished that with spinous leaves is the largest, while that which is altogether spinous is about the smallest. It is indeed, as was said, a very small class, and it would not be easy to find examples of such plants besides asparagus and

skorpios.2

³ Both of these flower after the autumnal equinox. Skorpios produces its flower in the fleshy swelling 4 below the top 5 of the spinous twig; at first it is white, but afterwards it becomes purplish. Asparagus produces alongside of the spines a small knob, and from this grows the flower, which is of small size. Skorpios has a single root which runs deep; asparagus roots very deep and its roots are numerous and matted, the upper part of them being in one piece,6 and from this the actual shoots spring. The stalk comes up from the plant in spring and is edible; afterwards, as the season advances, it acquires its rough and spinous character 7; the bloom appears not only on this stalk, but on those of previous years, for the Such is the character of stalk is not annual. plants which are altogether spinous.

⁸ Of those which have spinous leaves the largest class, one may say, consists of those plants which

⁴ ἐποιδοῦντι conj. Scal.; ἐπωδοῦντι U; ἐσποδοῦντι MAld.
5 τὸ ἄκρον conj. Scal.; τὸ ἄκρατον UMAld.; τῆς ἀκάνθης om. Scal.

⁶ i.e. tuberous. cf. Col. 11. 3. 43; Pall. 3. 24. 8; 4. 9. 11.
⁷ ἐξακανθοῦται conj. Link. ap. Sch.; ἐξανθοῦται UM; ἐξανθεῖται Ald.

8 Plin. 21. 94,

άπλως είπειν άκανωδες τυγχάνει λέγω δὲ τὸ ακανώδες, ὅτι τὸ κύημα καὶ ἐν ῷ τὸ ἄνθος ἡ καὶ ὁ καρπός ἄκανος ἡ ἀκανῶδες πάντων ἐστί. διαφοραν δε έχει εν εαυτώ και μεγέθει και σχήματι καὶ γρώματι καὶ πλήθει καὶ ολιγότητι τῶν ἄκανθων καὶ των άλλων. ἔξω γὰρ ολίγων πάνυ, καθάπερ τοῦ στρουθίου τε καὶ τοῦ σόγκου καὶ εἴ τινων έτέρων, τὰ λοιπὰ πάντα ὡς εἰπεῖν τοιαύτην έχει την φύσιν έπει και ο σόγκος την γε φύσιν άκανθώδη έχει, τὸ δὲ σπερματικὸν οὐχ ὅμοιον. άλλὰ τά γε τοιαῦτα πάντα οἶον ἄκορνα λευκάκανθα γάλκειος κνήκος πολυάκανθος άτρακτυλίς ονόπυξος ίξίνη χαμαιλέων πλην ούτος ού φυλλάκανθος, σκόλυμος δέ, δς καὶ λειμωνία, φυλλ-4 άκανθος καὶ τάλλα, πλείω γάρ ἐστι. διαφέρουσι δ' άλλήλων πρὸς τοῖς εἰρημένοις τῷ τὰ μὲν πολύκαυλα είναι καὶ ἀποφύσεις έχειν, ώσπερ ὁ ἄκανος, τὰ δὲ μονόκαυλα καὶ μὴ ἔχειν, ὥσπερ ὁ κνῆκος, ένια δ' ἄνωθεν έχειν έξ ἄκρου, καθάπερ το ρύτρος. καὶ τὰ μὲν εὐθὺς τοῖς πρώτοις ὑετοῖς βλαστάνειν τὰ δ' ὕστερον, ἔνια δὲ καὶ τοῦ θέρους, ὥσπερ καὶ ἡ τετράλιξ ὑπό τινων καλουμένη καὶ ἡ ἰξίνη· καὶ έπὶ τῶν ἀνθῶν ὁμοίως οψιανθής γὰρ ὁ σκόλυμος καὶ ἐπὶ πολὺν χρόνον.

² ἀκανῶδες conj. Sch.; ἀκανθῶδες Ald.H.; acanaceum G.

* άκανος ἡ ἀκανῶδες Ald.; ἄκανθος ἡ ἀκανθῶδες mBas. v. supra.

⁴ σόγκος conj. Sch.; κνῆκος Ald. The correction seems necessary in view of 6. 4. 8.

⁵ ἄκορνα conj. Sch.; ἄκαρνα Ald. cf. Plin. l.c.

¹ ἀκανῶδες mBas.; ἀκανθῶδες Ald. cf. 1. 13. 3, where ἀκανῶδες is restored by W.'s certain conj.

⁶ bs και λειμωνία I conj.; η και λειμωνία conj. Scal. from

ENQUIRY INTO PLANTS, VI. IV. 3-4

are thistle-like.1 by which 2 I mean that the swollen part, that part which contains the flower, or, it may be, the fruit, is in all cases a thistle-head,3 or has that appearance. However there are differences in the 'head' itself, in size shape colour number of spines and in other respects. For, apart from quite a few plants, such as soap-wort sow-thistle and possibly some others, nearly all the rest have this character (even sow-thistle 4 has a spinous character, but its seed-process is different). The list includes all the following: akorna 5 milk-thistle khalkeios safflower polyakanthos distaff-thistle onopyxos ixine chamaeleon (the last-named, however, has not spinous leaves, though golden thistle, which is also called 'meadow-thistle,' 6 has 7), and so on, for there are many more. These differ from one another not only in the aforesaid ways, but in that some of them have many stalks and side-growths, like the pinethistle, while some have a single stalk and no sidegrowths, like the safflower, and some again have out-growths above from the top of the plant, like the globe-thistle.8 Again some grow directly the first rains come, others at a later time, some again in summer, as the plant which some call yellow starthistle, and ixine. So too 10 the flowering-time differs: golden thistle blooms late 11 and is in bloom for a long time.

Plin. 22. 86; ήλυλειμωνία UM,; ήλυλειμωνία MaAld. καί λειμωνία conj. W. But λειμωνία is not mentioned again in the following description, which is against its being a distinct plant from σκόλυμος.

⁷ φυλλάκανθος I conj.; φυλλάκανθα MSS.
 ⁸ ρύτρος: rhutrum G; but Plin. l.c. has eryngen.

10 καὶ ἐπὶ conj. Sch.; καὶ ἡ ἐπὶ Ald. H. 11 δψιανθής conj. Bod. from Plin. l.c., floret sero et diu; εὐανθης Ald.

25

Διαφοραί δὲ τῶν μὲν ἀκάνων οὐκ εἰσί, τῆς κυήκου δ' εἰσίν· ή μεν γαρ άγρία ή δ' ήμερος. της δ' άγρίας δύο είδη, τὸ μὲν προσεμφερές σφόδρα τῷ ἡμέρω πλὴν εὐθυκαυλότερον, δι' δ καὶ πηνίοις ένιαι των άργαίων έγρωντο γυναικών, καρπον δε έχει μέλανα και μέγαν και πικρόν. δ' έτέρα δασεία καὶ τοὺς καυλοὺς ἔχει σογκώδεις. ώστε τρόπον τινὰ ἐπιγειόκαυλος γίνεται διὰ γὰρ μαλακότητα των καυλών κατακλίνεται πρός τάς άρούρας· καρπου δ' έχει μικρου πώγωνος· σπερ-ματώδεις πασαι, πλην μειζόσι και πυκνοτέροις αί ἄγριαι. ἴδιον δὲ ἔχει πρὸς τὰ ἄλλα ἄγρια· τὰ μεν γάρ σκληρότερα και άκανθωδέστερα των ήμέρων, αΰτη δὲ μαλακωτέρα καὶ λειοτέρα.

'Η δ' ἄκορνα προσεμφερής ώς άπλως είπειν κατά την πρόσοψιν τη κνήκω τη ήμέρω, χρώμα δ' ἐπίξανθον ἔχει καὶ χυλον λιπαρόν. ἀτρακτυλὶς δέ τις καλεῖται καὶ λευκοτέρα τούτων ἴδιον δὲ ἔχει τὸ περὶ τὸ φύλλον, ὅτι ἀφαιρούμενον καὶ τη σαρκί προσφερόμενον αίματώδη ποιεί τον χυλόν, δι' δ καὶ φόνον ἔνιοι καλοῦσι τὴν ἄκανθαν ταύτην έχει δε καὶ τὴν όσμὴν δεινὴν καὶ φονώδη. όψε δε και τελειοί τον καρπον προς το μετό-

 2 πηνίοις conj. R. Const.; σπληνείοις U; σπληνίοις MAld; colu G and Plin. l.c.; cf. Diose. 3. 107.

⁴ καυλών conj. Scal. from Plin. l.c.; φύλλων Ald.

¹ ἀκάνων Ald.; ἀκαρνῶν mBas.; ἀκάνου or ἀκόρνης conj. Sch., the plural being awkward.

³ σογκώδεις: Plin. l.c. seems to have read δγκώδεις (torosiore caule).

⁵ μικρον conj. Spr. from Plin. l.c. (minutum semen); πικρον

⁶ πώγωνος· σπερματώδεις Ald.: so U, but πωγωνος, and M, but 26

ENQUIRY INTO PLANTS, VI. IV. 5-6

Pine-thistle 1 has but one kind, but there are different kinds of safflower, the wild and the culti-Again of the wild kind there are two forms, one very like the cultivated except that the stalk is straighter: wherefore in ancient times women sometimes used it to make distaffs.2 It has a fruit which is black large and bitter. The other is leafy, and its stalks are like those of the sow-thistle,3 so that to some extent it comes to have a prostrate stem: for on account of the softness of the stalks 4 it bends down towards the ground; and it has a small 5 fruit, which is bearded. All the forms produce abundant seed,6 but it is larger 7 and more crowded in the wild forms. This kind has also a peculiarity as compared with other wild plants; these are usually coarser and more spinous than the cultivated forms, but in this plant the wild form is softer and smoother.

The akorna resembles in a general way in appearance the cultivated safflower, but has a yellowish colour and a sticky juice. There is also a plant called distaff-thistle, which is whiter than these. A peculiarity of the leaf of this is that, if it is stripped off and applied to the flesh, the contact makes the juice blood-coloured, wherefore some call this kind of spinous plant 'blood-wort'; also it has an abominable smell, like that of blood; it matures its fruit late,

πωγωνδs: G. has fructum amarum (see last note) frequentem barbaeque modo hirsutum gignit: sunt ambo seminosa. W. conj. πωγωνοσπέρματα δ' είσὶ πᾶσαι, which is not convincing. I have retained the corrupt text and translate in the light of G.

τ μείζοσι: sc. σπέρμασι, but σπερματώδειs cannot be right.
 * χυλον add. St.; om. Ald.; succo pingui G; pinguiore succo Plin. l.c.
 * cf. 9. 1, 1. Plin. 21. 95.

πωρου. τὸ δ' ὅλον ὡς ἀπλῶς εἰπεῖν ἄπασα ἡ ἀκανικὴ φύσις ὀψίκαρπος. ἄπαντα δὲ ταῦτα φύεται καὶ ἀπὸ τοῦ σπέρματος καὶ ἀπὸ τῆς ρίζης, ὥστε βραχύν τινα γίνεσθαι τὸν ἀνὰ μέσον χρόνον τῆς ἐκφύσεως τε καὶ τῆς τοῦ σπέρματος τελειώσεως.

Τοῦ σκολύμου δὲ οὐχ ὅτι τοῦτο μόνον ἴδιον, ὅτι τὴν ρίζαν ἐδώδιμον ἔχει καὶ ἑφθὴν καὶ ὡμήν, ἀλλὰ καὶ ὅτι τότε ἀρίστην ὅταν ἀνθῆ καὶ ὅτι σκληρυνομένη ἀφίησιν ὀπόν. ἴδιον δὲ καὶ τὸ τῆς

άνθήσεως έπεὶ περὶ τροπάς.

Σαρκώδης δὲ καὶ ἐδώδιμος ἡ τοῦ σόγκου· ἡ δὲ κύησις οὐκ ἀκανώδης ἀλλὰ προμήκης αὐτοῦ· καὶ τοῦτ' ἴδιον μόνον ἔχει τῶν φυλλακάνθων ἀντεστραμμένως ἡ ὁ χαμαιλέων· ὁ μὲν γὰρ ἀφυλλάκανθος ὢν ἀκανίζει. γηράσκον δὲ τὸ ἄνθος ἐκπαπποῦται, καθάπερ τὸ τῆς ἀπάπης καὶ τὸ τῆς μυρίκης καὶ ὅσα παραπλήσια τούτοις. παρακολουθεῖ δὲ μέχρι τοῦ θέρους τὸ μὲν κυοῦν τὸ δὲ ἀνθοῦν τὸ δὲ σπέρμα τίκτον, μικρὰν ἰκμάδα καὶ κέντρον ἔχον· ξηραινόμενον δὲ τὸ φύλλον διαχείται καὶ οὐκέτι κεντεῖ.

'Η ἰξίνη δὲ φύεται μὲν οὐ πολλαχοῦ, ῥιζόφυλλον δέ ἐστιν. ἀπὸ δὲ τῆς ῥίζης μέσης ὁ σπερματικὸς ἄκανος ἐπιπέφυκεν, ὧσπερ μῆλον εὖ

² cf. Hes. Op. 582.

3 σόγκου conj. C. Hoffmann; ύγκου Ald.

¹ ἀκανική conj. Bod., cf. 6. 4. 4 nn.; ἀκανθική Ald.

⁴ κύησις: i.e. flower-head. cf. κύημα 6. 4. 3; Plin. 21. 94.

⁵ ἀκανώδης conj. Scal.; ἀκανθώδης Ald. cf. 6. 4. 3 nn.

⁶ cf. 6. 4. 3. T.'s information seems to be incorrect, as 28

ENQUIRY INTO PLANTS, VI. iv. 6-9

towards autumn. Indeed, generally speaking, all plants like the thistle-tribe 1 are late fruiting. All these plants grow both from seed and from the root, so that there is but a short period between the beginning of growth and the maturing of the seed.

Golden thistle has not only this peculiarity, that it has a root which is edible, whether boiled or raw, but the root is best when the plant is in flower, and, as it becomes hard, it produces a juice. The flowering

time 2 is also peculiar, about the solstice.

The root of the sow-thistle³ is also fleshy and edible; but the swollen part is elongated and not thistle-like ; and, alone of the spinous-leaved plants, it has this peculiarity, in which it is the reverse of the chamaeleon, (for that plant, though it has not spinous leaves, has a thistle-like flower-head). The flower of the sow-thistle, as it ages, turns into down, as do that of the dandelion the tamarisk and other plants like these. In its growth there is a succession up to the summer, part forming flowers, part flowering, and part producing seed to; this the leaf, as it dries, becomes flaccid and no longer pricks.

Ixine does not grow in many places, and it has leaves on the root. From the middle of the root grows the seed-bearing thistle-head, which is like

both of the plants which he calls $\chi a \mu a i \lambda \ell \omega \nu$ (see Index) have spinous leaves.

⁷ ἀπάπης conj. Sch., cf. 7. 8. 3; πάπνης U; δαπάνης P; δάφνης Ald.

8 μυρίκης conj. Sch.; μυρίνης M; μυρρίνης Ald.

9 cf. Plin. l.c.

10 σπέρμα τίκτον Ι conj.; σπέρματος μέν Ald.H.; σπερμοτόκουν conj. Sch.

11 Text perhaps defective.

μάλα ἐπικεκρυμμένον ὑπὸ τῶν φύλλων οὖτος δὲ ἐπὶ τοῦ ἄκρου φέρει τὸ δάκρυον εὔστομον, καὶ τοῦτό ἐστιν ἡ ἀκανθικὴ μαστίχη. τα μέν οὖν καὶ τὰ τοιαῦτα πανταγοῦ σγεδόν έστιν.

10 'Η δὲ κάκτος καλουμένη περὶ Σικελίαν μόνον, έν τῆ Ἑλλάδι δὲ οὐκ ἔστιν. ἴδιον δὲ παρὰ τἄλλα τὸ φυτόν· ἀφίησι γὰρ εὐθὺς ἀπὸ τῆς ρίζης καυλούς ἐπιγείους, τὸ δὲ φύλλον ἔχει πλατύ καὶ ἀκανθώδες καλοῦσι δὲ τούς καυλούς τούτους κάκτους εδώδιμοι δέ είσι περιλεπόμενοι μικρον επίπικροι, και θησαυρίζουσιν αὐτούς εν

11 "Ετερον δε καυλον ορθον αφίησιν, ον καλουσι πτέρνικα γίνεται δὲ καὶ οὖτος ἐδώδιμος πλην άθησαύριστος. τὸ δὲ περικάρπιον, ἐν ῷ τὸ σπέρμα, την μεν μορφην ακανώδες, αφαιρεθέντων δὲ τῶν παππωδῶν σπερμάτων ἐδώδιμον καὶ τοῦτο καὶ ἐμφερὲς τῷ τοῦ φοίνικος ἐγ-κεφάλῳ· καλοῦσι δὲ αὐτὸ σκαλίαν. τὰ μὲν οὖν φυλλάκανθα σκεπτέον ἐν τοιαύταις διαφοραίς.

V. Τὰ δὲ καὶ παρὰ τὴν ἄκανθαν ἔγοντα φύλλον, οίον τὰ τοιαῦτα φέως ὄνωνις παντάδουσα τρίβολος ίππόφεως μυάκανθος τε σφόδρα καὶ τὸ φύλλον έχει σαρκώδες πολυ-

⁴ Plin. 21. 97; Athen. 2. 83.

6 ἀκανῶδες conj. Sch.: ἀκανθῶδες Ald.

¹ ύπὸ conj. Sch; ἐπὶ Ald.; Plin. l.c. malum contectum sua ² cf. 9. 1. 3. ⁸ cf. Plin. 12. 72.

⁵ πλατὸ add. Scal. from Athen. l.c., cf. Plin. l.c.; om. Ald. H. The 'stems' are the petioles of the leaves.

ENQUIRY INTO PLANTS, VI. IV. Q-V. 1

an apple and well hidden by 1 the leaves; this on its head produces its gum,2 which is pleasant to the taste, and this is the 'thorn-mastich.' 3 plants and others like them are found almost

everywhere.

4 But the plant called kaktos (cardoon) grows only in Sicily, and not in Hellas. It is a plant quite different from any other; for it sends up straight from the root stems which creep on the ground, and its leaf is broad 5 and spinous: these stems are called kaktoi: they are edible, if peeled, and are slightly

bitter, and men preserve them in brine.

There is another kind which sends up an erect stem, called the pternix. This too is edible, but cannot be preserved. The fruit-vessel, which contains the seed, is in shape like a thistle-head 6: and when the downy seeds are taken off, this too is edible and resembles the 'brain' 7 of the palm; and it is called Such are the different characteristics in skalias.8 the light of which we may observe the spinousleaved plants.

V. Examples of plants which have leaves as well as spines are pheos 9 rest-harrow star-thistle caltrop 'horse-pheos' 10 (spurge) butcher's broom 11 . . . , 12 and it has a fleshy leaf: it is much divided and has

⁷ i.e. 'cabbage,' cf. 2. 6. 2.

11 Diosc. 2. 125; Plin. 19. 151.

⁸ ascaliam Plin. l.c.; ἀσκάληρον Athen. l.c. Modern Greek σκάληρα. English 'bottom.' See Index κάκτος (2).

⁹ φέως conj. St.; φλέως Ald. cf. 6. 1. 3.

¹⁰ iπποφέως conj. Salm., cf. 6. 5. 2; iππόφυον Ald. cf. Plin. 21. 91.

¹² Text defective: the end of one sentence is missing and the beginning of the next, containing the name of a plant. G attaches the following description to pews. The plants presently described do not correspond to this list.

σχιδές δέ καὶ πολύρριζον, οὐ μὴν κατά βάθους γε τὰς ρίζας ἔχον. βλαστάνει δὲ ἄμα Πλειάδι καὶ τοῖς πρώτοις ἀρότοις καὶ ἀφίησι τότε τὸ φύλλον· οὐ γάρ ἐστιν ἐπέτειον ἀλλὰ χρονιώ-

τερον.

Τὸ δὲ τῆς καππάριος ἴδιον, ὥσπερ ἐλέχθη, παρὰ ταῦτα καὶ γὰρ τὸ φύλλον ἐπακανθίζον ἔχει καὶ τὸν καυλόν, οὐχ ὅσπερ ὁ φέως καὶ ίππόφεως ἀνάκανθα τοις φύλλοις μονόρριζον δὲ καὶ ἐπίγειον καὶ χαμαίκαυλον βλαστάνει δὲ καὶ ἀνθεῖ τοῦ θέρους καὶ διαμένει τὸ φύλλον χλωρὸν ἄχρι Πλειάδος. χαίρει δὲ ὑφάμμοις καὶ λεπτογείοις χωρίοις λέγεται δὲ ὡς ἐν τοῖς ἐργασίμοις οὐ θέλει φύεσθαι, καὶ ταῦτα περὶ τὰ άστη και έν εύγείοις τόποις φυομένη και ούχ ωσπερ σίλφιον εν δρεινοίς τούτο μεν <ούν> ού πάντως άληθές.

Ο δὲ τρίβολος ἴδιον ἔχει, διότι περικαρπι-άκανθός ἐστι. δύο δ' αὐτοῦ γένη τὸ μὲν γὰρ έχει φύλλον έρεβινθώδες, έτερος δε φυλλάκανθος. επίγειοι δε άμφω και πολλαχη σχιζόμενοι όψιβλαστής δὲ μᾶλλον ὁ φυλλάκανθος καὶ φύεται περὶ τὰς αὐλάς. τὸ δὲ σπέρμα τοῦ μὲν πρωΐου σησαμῶδες, τοῦ δὲ ὀψίου στρογγύλον ἐπίμελαν έν λοβφ. καὶ τὰ μὲν οὖν παρὰ τὰ φύλλα καὶ ἄκανθαν ἔχοντα σχεδὸν ἐν τούτοις.

Ή δ' ὄνωνίς ἐστι πτορθάκανθον ἐπέτειον δὲ τὸ φύλλον ἔχει πηγανώδες παραπεφυκὸς παρ'

⁵ Plin. 21, 91.

ἀρότοις conj. Bod.; ἀρότροις Ald. cf. 8. 1. 2.
 τότε conj. St.; τοῦτο Ald. s. f. Pall. 10. 13. 2.

⁴ δ φέως conj. St.; δφεως Ald. Bas. Cam. H.; δ φλεως mBas.

ENQUIRY INTO PLANTS, VI. v. 1-3

many roots, but is not deep-rooting. It grows at the rising of the Pleiad, the first seed-time, and then 2 puts forth its leaf; for it is not annual, but lives

longer than one year.

³Caper, as was said, is quite distinct from these; it has a spinous leaf and a spinous stem, whereas pheos and 'horse-pheos' have no spines on their leaves 5; it has a single 6 root, is low-growing.7 and has a creeping stem; it grows and flowers in summer, and the leaf remains green till the rising of the Pleiad. It rejoices in sandy light soils, and it is said that it is unwilling to grow on cultivated land, and that though it grows near towns and in good soil, and not, like silphium, in mountain country. This account however 8 is not altogether accurate.

⁹ A peculiarity of caltrop is that it is spinousfruited. 10 There are two kinds; one has a leaf like that of chick-pea, the other has spinous leaves. Both are low-growing and much divided, but the spinous-leaved form grows later and is found near enclosures. The seed of the early kind is like that of sesame, that of the late kind is round and blackish and enclosed in a pod. These may serve as examples of plants which have spines as well as leaves.11

12 Rest-harrow has spines on the shoots; the leaf, which is annual,13 is like that of rue, and grows right along the stem, so that the general appearance is

7 cf. 7. 8. 1.

8 obv add. W. (in comm.) from G. 10 cf. 6. 1. 3. ⁹ Plin. 21. 98.

13 I have altered the punctuation; πτορθάκανθον, ἐπέτειον δέτὸ φ. κτλ. W. after UMP.

33

VOL. II.



D

⁶ Diosc. 2. 173 gives a different account.

¹¹ τὰ μέν οδυ παρά τὰ φύλλα conj. Sch. (οδυ add. W.); τὰ μέν ¹² Plin. 21. 98. οδν ωσπερ ανάφυλλα Ald.H.

όλον τὸν καυλόν, ὥστε καθάπερ στεφάνου τὴν όλην είναι μορφήν, διαλαμβανομένων έπαλλήλων κολοβοανθής δέ καὶ έλλοβόκαρπος άδια-4 Φράκτως φύεται δ' έν τη γλίσχρα καλ γανώδει καὶ μάλιστα ἐν τῆ σπορίμω καὶ γεωργουμένη, δι' δ καὶ πολέμιον τοῖς γεωργοῖς καὶ ἔστι δυσώλεθρος όταν γὰρ λάβη χώρας βάθος, ώθεῖται κάτω εὐθὺς καὶ καθ' ἕκαστον ἔτος ἀποφύσεις άφεμένη είς τὰ πλάγια πάλιν είς τὸ έτερον ώθειται κάτω σπαστέα μέν οὖν ὅλη τοῦτο δὲ Βραγείσης γίνεται της γης καὶ ἀπόλλυται ράον ἐὰν δὲ καὶ μικρὸν ἀπολειφθή, ἀπὸ τούτου πάλιν βλαστάνει ἄργεται δὲ τῆς βλαστήσεως θέρους τελειουται δὲ μετοπώρου. τὰ μὲν οὖν άγρια τῶν Φρυγανικῶν ἐκ τούτων θεωρείσθω.

VI. Τὰ δὲ ημερα βραχεῖάν τινα ἔχει θεωρίαν,

άπερ εν τοίς στεφανωματικοίς έστι.

Τὰ δὲ καθ' ὅλου πειρατέον περὶ στεφανωμάτων εἰπεῖν, ὅπως ἄπαν περιληφθἢ τὸ γένος. ἡ γὰρ στεφανωματικὴ φύσις ἰδίαν τινὰ ἔχει τάξιν, ἐπιμιγνυμένη δὲ τὰ μὲν τοῖς φρυγανικοῖς τὰ δὲ τοῖς ποιώδεσι· δι' δ κἀκεῖνα συμπεριληπτέον ἐπιμιμνησκομένους ὡς ἀν ἢ ὁ καιρός, ἀρξαμένους 2 πρῶτον ἀπὸ τῶν φρυγανικῶν. διχῆ δὲ ἡ τούτων

3 κολοβοανθής; cf. 8. 3. 3.

¹ Evidently some conventional way of making a wreath.

² διαλαμβανομένων ἐπαλλήλων conj. W.; διαλαμβανομένην ὑπ' ἀλλήλων Ald. cf. Plat. Prot. 346 E, where the verb means 'to punctuate.'

ENQUIRY INTO PLANTS, VI. v. 3-vi. 2

that of a garland, the leaves being set at intervals alternately along it 2; the flower is irregular, 3 and the fruit contained in a pod,4 which is not divided into compartments.⁵ It grows in sticky rich soil and especially in sown and cultivated land; wherefore it is an enemy to husbandmen, and it is hard, to kill; for, when it gets hold of a piece of ground, it immediately pushes its roots down deep,6 and every year it sends up new growths at the sides and the next year 7 it roots these again. Wherefore it has to be dragged up entire 8; this is done when the ground has been moistened, and then it is easier to destroy. But, if but a small piece is left, it shoots again from this. It begins to grow in summer and completes its growth in autumn. Let these examples serve for a survey of the wild forms of under-shrubs.

Of cultivated under-shrubs (coronary plants), with which are included those coronary plants which are herbaceous.

VI. The cultivated kinds need but a brief survey; these 9 come under the class of coronary plants.

Of coronary plants we must endeavour to give a general account, so that the whole class may be included. This group has a somewhat peculiar position, since it overlaps partly the under-shrubs, partly the herbaceous plants; wherefore the latter must also be included and we must mention them as occasion serves, taking first the under-shrubs.

⁹ ἄπερ conj. Sch.; εἴπερ UMAld.G.

⁴ ελλοβόκαρπος conj. Sch.; ελλοβοάνθης Ald.

 ⁶ ὧθείται κάτω conj. Sch.; ὧθεί τὰ κάτω Ald.
 ⁷ εἰς τὸ ἔτερον, sc. ἔτος; τῷ ἐτέρφ conj. Sch.

⁸ σπαστέα μεν οδυ δλη conj. W.; σταθείσα μεν ούτως δλη Ald.

διαίρεσις ή κατά τὴν χρείαν. τῶν μὲν γὰρ τὸ άνθος μόνον χρήσιμον καὶ τούτων τὸ μὲν εὔοσμον. ώσπερ ζον, τὸ δ' ἄνοσμον, ώσπερ διόσανθος Φλόξ. των δέ καὶ οι κλώνες καὶ τὰ φύλλα καὶ όλως ή πασα φύσις εὔοσμος, οἶον έρπύλλου έλενίου σισυμβρίου των άλλων. άμφω δε φρυγανικά. κακείνων των ανθικών πολλών ή φύσις φρυγανώδης, ή μὲν ἐπέτειος οὖσα μόνον, ή δὲ πολυγρονιωτέρα, πλην ιωνίας της μελαίνης αυτη γαρ ακλων όλως αλλά προσριζόφυλλος καὶ αείφυλλος, ώς δέ τινές φασι καὶ δυναμένη δι' όλου φέρειν τὸ ἄνθος, ἐὰν τρόπω τινὶ θεραπεύηται. τοῦτο μὲν ἴδιον ἂν ἔχοι.

Των δὲ ἄλλων μαλλον δὲ των πάντων αί μὲν όλαι μορφαί σχεδον πάσι φανεραί εί δέ τινας άλλας ιδιότητας έχουσι, ταύτας λεκτέον, οίον εί τὰ μὲν ἀπλα δοκεί τοις εἴδεσιν εἶναι τὰ δὲ ἔγειν

διαφοράς.

Απλα μεν οὖν τὰ ξυλώδη, καθάπερ έρπυλλος σισύμβριον έλένιον πλην εί τὰ μὲν ἄγρια τὰ δὲ ημερα καὶ <τὰ μὲν> εὔοσμα τὰ δὲ ἀοσμότερά ἐστι· τούτων δὲ καὶ αἱ θεραπεῖαι καὶ αἱ χῶραι διάφοροι καὶ οἱ ἀέρες. ἔνια δὲ καὶ τῶν ἀνθῶν, οἶον τὸ μέλαν ἴον· οὐ γὰρ ἔχειν δοκεῖ τοῦτο διαφοράν

Plin. 21, 59.

4 olov el conj. W.; oti Ald. 5 odv conj. W.; olov Ald.

² So Plin. l.c.; but Nic. ap. Athen. 15. 31 calls this flower fragrant. ⁸ πολλών conj. W.; πολλά UMAld.

ENQUIRY INTO PLANTS, VI. vi. 2-3

¹These may be divided into two groups according to their uses. Of some only the flower is serviceable: and of these some are sweet-scented, as gilliflower, some scentless, as carnation 2 and wallflower. Of others again the branches leaves and in fact the whole growth are sweet-scented, as with tufted thyme calamint bergamot-mint and the rest. Both groups however belong to the under-shrubs. And of the first-mentioned, those valued for their flowers, the growth is in most 3 cases that of undershrubs, (in some annual merely, in others of longer duration) except in the violet; for this is altogether without branches, its leaves grow close to the root, and it is always in leaf; while, as some say, it is able to bear flowers continuously, if it is tended in a certain way. This may be considered a peculiar characteristic of this plant.

Of the others, or rather of all the group, the general appearance is in each case plain to all; any peculiarities that they may exhibit we must mention, for instance, if 4 some appear to have but a single form, while others have various forms.

Thus 5 those of woody character, as tufted thyme bergamot-mint calamint, have but one form, unless one counts wild and cultivated, scented and scentless plants, as belonging to distinct forms; and again there are with these plants differences of culture of position and of climate. Some also 6 of the group valued for their flowers 7 have each but one form, for instance, the black ion (violet); for this does not appear to have different forms

⁶ ξνια δὲ καὶ conj. W.; ξνιοι δὲ UMAld.

⁷ ἀνθῶν in the sense of ἀνθικῶν § 2, which perhaps should be read here.

ῶσπερ τὸ λευκόν ἐμφανης γὰρ ἡ τούτων χροιὰ διαλλάττουσα, καὶ ἔτι δὴ μᾶλλον ἡ τῶν κρίνων, εἴπερ δή, καθάπερ φασίν, ἔνια καὶ πορφυρὰ ἐστι.
Τῶν δὲ ρόδων πολλαὶ διαφοραὶ πλήθει τε φύλλων καὶ ὀλιγότητι καὶ τραχύτητι καὶ λειότητι καὶ εὐχροίᾳ καὶ εὐοσμίᾳ. τὰ μὲν γὰρ πλεῖστα πεντάφυλλα, τὰ δὲ δωδεκάφυλλα καὶ εἰκοσίφυλλα, τὰ δ' ἔτι πολλῷ πλεῖον ὑπεραίροντα τούτων ἔνια γὰρ εἶναί φασιν ἃ καὶ καλοῦσιν ἐκατοντάφυλλα πλεῖστα δὲ τὰ τοιαῦτά ἐστι περὶ Φιλίππους οὐτοι γὰρ λαμβάνοντες ἐκ τοῦ Παγγαίου φυτεύουσιν ἐκεῖ γὰρ γίνεται πολλά σμικρὰ δὲ σφόδρα τὰ ἐντὸς φύλλα ἡ γὰρ ἔκφυσις αὐτῶν οὕτως ὥστε εἶναι τὰ μὲν ἐκτὸς τὰ δ' ἐντός οὐκ εὔοσμα δὲ οὐδὲ μεγάλα τοῖς μεγέθεσιν. ἐν δὲ τοῖς μεγά-

όλον, ώσπερ ελέχθη, καὶ ἡ εὐχροια καὶ ἡ εὐοσμία παρὰ τοὺς τόπους ἐστίν· ἐπεὶ καὶ τὰ ἐν γἢ τἢ αὐτἢ γινόμενα ποιεῖ τινα παραλλαγὴν εὐοσμίας καὶ ἀοσμίας. εὐοσμότατα δὲ τὰ ἐν Κυρήνῃ, δι δ καὶ τὸ μύρον ἤδιστον. ἀπλῶς δὲ καὶ τῶν ἴων καὶ τῶν ἄλλων ἀνθῶν ἄκρατοι μάλιστα ἐκεῦθι

5 λοις εὐώδη μᾶλλον ὧν τραχὺ τὸ κάτω. τὸ δὲ

¹ cf. 6. 8. 1 n.; Diosc. 3. 102.

² Plin. 21. 14-21; Athen. 15. 29.

³ i.e. of the bark. cf. Plin. 21. 17, scabritia corticis.

⁴ Sc. in 'double' roses.

⁵ i.e. the hip; called $\ddot{\nu}\mu\phi\alpha\lambda$ os Arist. Probl. 12. 8, where the same statement is made; called $\mu\hat{\eta}\lambda$ oν below, § 6.

ENQUIRY INTO PLANTS, VI. vi. 3-5

like the white ion (gilliflower) in which the colour evidently varies; as does still more that of the lilies, if it be true, as some say, that there is a crimson kind.¹

² Among roses there are many differences, in the number of petals, in roughness,3 in beauty of colour, and in sweetness of scent. Most have five petals, but some have twelve or twenty, and some a great many more than these; for there are some, they say, which are even called 'hundredpetalled.' Most of such roses grow near Philippi; for the people of that place get them on Mount Pangaeus, where they are abundant, and plant them. However the inner petals 4 are very small, (the way in which they are produced being such that some are outside, some inside). Some kinds are not fragrant nor of large size. Among those which have large flowers those in which the part 5 below the flower is rough are the more fragrant. In general, as has been said, good colour and scent depend upon locality; for even bushes which are growing in the same 6 soil shew some variation in the presence or absence of a sweet scent. Sweetestscented of all are the roses of Cyrene, wherefore the perfume made from these is the sweetest. (Indeed it may be said generally that the scents 7 of the gilliflowers 8 also and of the other flowers of that place are the purest, and especially the

⁸ ? violets and gilliflowers : see Index.

⁶ τῆ αὐτῆ conj. Sch.; τοιαύτη U; τοιαῦτα Μ.

⁷ ἄκρατοι μάλιστα ἐκείθι αἱ ὁσμαί conj. Sch. after Saracenus on Diosc. 1. 25; Athen. l.c. (ἄκρατοι μάλιστα καὶ θείαι αἱ ὀσμαί); ἄκρατοι μάλιστα δὶ ἐκείνου αἱ ὀσμαί Ald.; ἐκεῖ αἱ ὀσμαί (rest uncertain) U. cf. C.P. 6. 18. 3.

αί όσμαί, διαφερόντως δὲ τοῦ κρόκου πλεῖστον γὰρ οὖτος δοκεῖ παραλλάττειν. φύεται μὲν οὖν ἡ ροδωνία καὶ ἐκ τοῦ σπέρματος ἔχει δὲ ὑπὸ τὸ ἄνθος ἐν τῷ μήλῳ κνηκῶδες ἡ ἀκανῶδες, ἔχον δέ τινα χνοῦν ὥστε ἐγγὺς εἶναι τῶν παππωδῶν σπερμάτων οὐ μὴν ἀλλὰ διὰ τὸ βραδέως παραγίνεσθαι κατακόπτοντες, ὡς ἐλέχθη, τὸν καυλὸν φυτεύουσιν. ἐπικαιομένη δὲ καὶ ἐπιτεμνομένη βέλτιον φέρει τὸ ἄνθος ἐωμένη γὰρ ἐξαύξεται καὶ ἀπολοχμοῦται. δεῖ δὲ καὶ μεταφυτεύειν πολλάκις καὶ γὰρ οὕτω φασὶ κάλλιον γίνεσθαι τὸ ῥόδον. αἱ δ' ἄγριαι τραχύτεραι καὶ ταῖς ράβδοις καὶ τοῖς φύλλοις, ἔτι δὲ ἄνθος ἀχρού στερον ἔχουσι καὶ ἔλαττον.

Τὸ δὲ ἴον τὸ μέλαν τοῦ λευκοῦ διαφέρει κατά τε ἄλλα καὶ κατ' αὐτὴν τὴν ἰωνίαν, ὅτι πλατύφυλλός τε καὶ ἐγγειόφυλλος καὶ σαρκόφυλλός

έστι, πολλην έχουσα ρίζαν.

Τὰ δὲ κρίνα τῆ μὲν χροιᾶ τὴν εἰρημένην ἔχει διαφοράν. μονόκαυλα δέ ἐστιν ὡς ἐπὶ πᾶν, δικαυλεῖ δὲ σπανίως· τάχα δὲ τοῦτο χώρας καὶ ἀέρος διαφορᾶς. καθ' ἔκαστον δὲ καυλὸν ότὲ μὲν ἐν κρίνον ότὲ δὲ πλείω γίνεται· βλαστάνει γὰρ τὸ ἄκρον· σπανιώτερα δὲ ταῦτα· ρίζαν δὲ ἔχει πολλὴν σαρκώδη καὶ στρογγύλην· ὁ δὲ καρπὸς

² ἀκανῶδες conj. Sch. from G, acanaceum; ἀνθῶδες UMAld.

3 παππωδών conj. Sch.; πρώτων Ald.

4 Plin. 21, 27.

¹ διαφερόντως δὲ τοῦ κρόκου conj. Saracenus from Athen. l.c.; διαφερόντως δὴ τοῦ χρόνου Ald. cf Callim. Hymn to Apollo 83, whence it appears that an autumnal crocus (crocus sativus) is meant. See below § 10.

ENQUIRY INTO PLANTS, VI. vi. 5-8

scent of the saffron-crocus, a plant which seems to vary in this respect more than any other). Roses can be grown from seed, which is to be found below the flower in the 'apple,' and is like that of safflower or pine-thistle, but it has a sort of fluff, so that it is not unlike the seeds which have a pappus. As however the plant comes slowly from seed, they make cuttings of the stem, as has been said, and plant them. If the bush is burnt or cut over, it bears better flowers; for, if left to itself, it grows luxuriantly and makes too much wood. Also it has to be often transplanted; for then, they say, the roses are improved. The wild kinds are rougher both in stem and in leaf, and have also smaller flowers of a duller colour.

⁴ The black ion (violet) differs from the white ion (gilliflower) not only in other respects but in the plant itself, in that in the former the leaves are broad, lie close to the ground, and are fleshy, and there is much root.

⁵ Krina (lilies) shew the variation in colour which has been already mentioned.⁶ The plant has in general a single stem, but occasionally divides into two, which may be due to differences⁷ in position and climate. On each stem grows sometimes one flower, but sometimes more; (for it is the top of the stem which produces the flower⁸) but this sort is less common. There is an ample root, which is fleshy and round. If the fruit is taken off, it

⁷ διαφορᾶς U; διαφορᾶ W. after Sch.

⁵ Plin. 21. 25. The account of herbaceous coronary plants seems to begin here. cf. 6. 6. 10. ⁶ 6. 6. 3.

⁸ βλαστάνει. But this word in T. has usually a more general sense. ? 'for in that case the top of the stem branches' (lit. 'makes fresh growth').

άφαιρούμενος έκβλαστάνει καὶ άποδίδωσι τὸ κρίνον πλην έλαττον. ποιεί δέ τινα καλ δακρυώδη συοροήν, ην και φυτεύουσιν, ώσπερ είπομεν.

Ο δε νάρκισσος ή το λείριον, οι μεν γαρ τοῦτο οί δ' ἐκεῖνο καλοῦσι, τὸ μὲν ἐπὶ τῆ γῆ Φύλλον άσφοδελώδες έχει, πλατύτερον δὲ πολύ, καθάπερ ή κρινωνία, τὸν δὲ καυλὸν ἄφυλλον μὲν ποώδη δε και εξ άκρου το άνθος, και εν υμένι τινί καθάπερ εν άγγείω <καρπον> μέγαν εθ μάλα καλ μέλανα τη χροιά σχήματι δὲ προμήκη. οὖτος δ' έκπίπτων ποιεί βλάστησιν αὐτόματον οὐ μὴν άλλα και συλλέγοντες πηγνύουσι και την ρίζαν φυτεύουσιν. έχει ρίζαν σαρκώδη στρογγύλην μεγάλην. όψιον δὲ σφόδρα μετὰ γὰρ Αρκτοῦρον ή άνθησις καὶ περὶ ἰσημερίαν.

'Ο δὲ κρόκος ποώδης μὲν τῆ φύσει, καθάπερ καὶ 10 ταῦτα, πλην φύλλω στενώ, σχεδον γάρ ώσπερ τριχόφυλλόν έστιν όψιανθές δέ σφόδρα καί οψιβλαστές ή πρωϊανθές, οποτέρως τις λαμβάνοι την ώραν <μετά> Πλειάδα γὰρ ἀνθεῖ καὶ ὀλίγας ήμέρας εὐθὺς δ' ἄμα τῷ φύλλω καὶ τὸ ἄνθος ώθει δοκεί δὲ καὶ πρότερον ρίζα δὲ πολλή καὶ σαρκώδης, καὶ τὸ ὅλον εὔζωον φιλεῖ δὲ καὶ πατεισθαι και γίνεται καλλίων κατατοιβομένης

¹ cf. 2. 2. 1 n., 9. 14; C.P. 1. 4. 4-6. Plin. 21. 26 describes a method of artificially producing crimson lilies from the bulbils of a white lily. cf. Geop. 11. 20.

² cf. 6. 8. 1 and 3. See Index.

³ cf. 7. 13. 1.

⁴ ποώδη: cf. 4. 10. 3.

ENQUIRY INTO PLANTS, VI. vi. 8-10

germinates and produces a fresh plant, but of smaller size; the plant also produces a sort of tear-like exudation, which men also plant, as we have said.1

The narcissus 2 or leirion (for some call it by the one name, some by the other) has its ground-leaves like those of the asphodel,3 but much broader, like those of the krinon (lily); its stem is leafless and grass-green 4 and bears the flower at the top; the fruit 5 is in a kind of membrane-like vessel, and is very large, black in colour, and oblong in shape. This as it falls germinates of its own accord; however men collect and set 6 the seed, and also plant the root, which is fleshy round and large. The plant blooms very late,7 after the setting of Arcturus about the equinox.

8 The saffron-crocus is herbaceous in character, like the above-mentioned plants,9 but has a narrow leaf; indeed the leaves are, as it were, hair-like; it blooms very late, and grows either late or early, according as one looks at the season 10; for it blooms after 11 the rising of the Pleiad and only for a few days. It pushes up the flower at once with the leaf, or even seems to do so earlier. The root 12 is large and fleshy, and the whole plant vigorous; it loves even to be trodden on and grows fairer when the root is crushed into the ground by the

⁵ καρπὸν omitted in MSS.; add. Dalec. from Diosc. 4. 158.

⁶ πηγνύουσι: cf. 7. 4. 3 n. ⁷ cf. C.P. 1. 10. 5; Plin. l.c. (a much confused passage).

⁸ Plin. 21. 31-34.

⁹ Sc. κρίνον and νάρκισσος; cf. 6. 6. 8 n.

¹⁰ i.e. whether at the end of one season or the beginning of the next. cf. C.P. 1. 10. 5. λαμβάνοι U; λαμβάνει Ald.

11 μετὰ add. W.

12 cf. 7. 9. 4.

πάτφ τῆς ρίζης· δι' δ καὶ παρὰ τὰς όδοὺς καὶ ἐν τοῖς κροτητοῖς κάλλιστος. ἡ δὲ φυτεία ἀπὸ

ρίζης.

1 Ταῦτα μὲν οὖν οὕτω γεννᾶται. τὰ δ΄ ἄλλα ἄνθη τὰ προειρημένα πάντα σπείρεται, οἶον ἰωνία διόσανθος ἴφυον φλὸξ ἡμεροκαλλές· καὶ γὰρ αὐτὰ καὶ αἱ ῥίζαι ξυλώδεις· σπείρεται δὲ καὶ ἡ οἰνάνθη· καὶ γὰρ καὶ τοῦτο ἀνθῶδες. τὰ μὲν οῦν ἀνθικὰ σχεδὸν ἐν τούτοις καὶ τοῖς ὁμοίοις ληπτέον.

VII. Τὰ δὲ ἔτερα πάντα μὲν ἀνθεῖ καὶ σπερμοφορεῖ, δοκεῖ δὲ οὐ πάντα διὰ τὸ μὴ φανερὸν εἶναί τινων τὸν καρπόν ἐπεὶ καὶ τὸ ἄνθος ἐνίων ἀμαυρόν ἀλλ' ὅτι βραδέως καὶ χαλεπωτέρως παραγίνεται, τῆ φυτεία χρῶνται μᾶλλον, εὤσπερ ἐλέχθη καὶ κατ' ἀρχάς. καίτοι διατείνονταί τινες ὡς οὐκ ἐχόντων καρπόν οἵ τε πεπειρᾶσθαι φάσκοντες καὶ τούτων εἰσίν, αὐτοὶ γὰρ ξηρᾶναι πολλάκις καὶ ἀποτρίψαι καὶ σπεῖραι, καὶ οὐδεπώποτε βλαστεῖν οὔτε ἔρπυλλον οὔτε ἐλένιον οὔτε σισύμβριον οὔτε μίνθαν πεπειρᾶσθαι γὰρ καὶ ταύτης. ἀλλ' ὅμως ἐκεῖνο ἀληθέστερον, ἥ τε τῶν ἀγρίων φύσις ἐπιμαρτυρεῖ· καὶ γὰρ ἔρπυλλός ἐστιν ἄγριος, ὃν κομίζοντες ἐκ τῶν ὀρῶν φυτεύουσι καὶ ἐν Σικυῶνι καὶ ᾿Αθήνησιν ἐκ τοῦ Ὑμηττοῦ· παρ' ἄλλοις δὲ ὅλως ὅρη πλήρη καὶ λόφοι, καθάπερ ἐν τῆ Θράκη· καὶ σισύμβριον

¹ πάτφ conj. Turneb. and others; κάτω Ald,

² κροτητοις: Plin. l.c. iuxta semitas ac fontes. Did he read κρουνοις?

ἀνθικὰ conj. Scal.; ἀκανθικὰ Ald. cf. 6. 6. 2.
 ἀλλ' ὅτι conj. W. from G; ἄλλα δὲ UMPAld.

ENQUIRY INTO PLANTS, VI. vi. 10-VII. 2

foot¹: wherefore it is fairest along the roads and in well-worn places.² It is propagated from the root.

These are the ways then in which the above plants are grown. All the above-mentioned flowers are grown from seed, as gilliflower carnation spike-lavender wall-flower martagon-lily; these plants themselves, as well as their roots, are woody. Drop-wort is also grown from seed; for that too is a plant grown for its flower. These and other plants like them may serve as examples of plants

grown for their flowers.8

VII. All the others flower and bear seed, though they do not all appear to do so, since in some cases the fruit is not obvious. Indeed in some the flower too is inconspicuous, but, because 4 these grow slowly and with some difficulty, men propagate them rather by off-shoots, as was said at the beginning. However some contend that they have no fruit: and there are men who have actually tried with the following plants 5; they have, they say, themselves often dried and rubbed out and sown the apparent fruit of thyme calamint bergamot-mint and green mint (for even that they have tried) and there was no germination from such sowing. However, the account given above is the truer, and the character of the wild forms testifies to this; for there is also a wild thyme (Attic thyme 6), which they bring from the mountains and plant at Sicyon, or from Hymettus and plant at Athens; and in other districts the mountains and hills 7 are quite covered with it, for instance in Thrace. There is also a

⁵ οῖ τε... εἰσίν transposed by Sch.; in MSS. after ἀληθέστερον.

⁶ Plin. 19. 172; Athen. 15. 28. λόφοι conj. W.; τόποι Ald.

δὲ καὶ τάλλα δριμυτέραν ἔχοντα τὴν ὀσμήν. έρπυλλος δ' ενίστε καὶ παντελώς θυμώδης α

δήλον ὅτι ταύτην τὴν γένεσιν λαμβάνει.

Αβρότονον δὲ μᾶλλον ἀπὸ σπέρματος βλαστάνει ή ἀπὸ ρίζης καὶ παρασπάδος γαλεπώς δὲ καὶ ἀπὸ σπέρματος προμοσχευόμενον <δέ> ἐν οστράκοις, ώσπερ οι 'Αδώνιδος κήποι, του θέρους. δύσριγον γάρ σφόδρα καὶ ὅλως ἐπίκηρον καὶ ὅποι ό ήλιος σφόδρα λάμπει εμβιῶσαν δὲ καὶ αὐξηθεν μέγα καὶ ἰσχυρὸν καὶ δενδρῶδες ὥσπερ τὸ πήγανον, πλην ξυλωδέστερον πολύ τοῦτο καὶ ξηρότερον

καὶ αὐχμωδέστερον.

Ο δε αμαρακος αμφοτέρως φύεται, καὶ απὸ παρασπάδος καὶ ἀπὸ σπέρματος πολύσπερμον δέ, καὶ τὸ σπέρμα εὔοσμον ὀσμῆ μαλακωτέρα· δύναται δὲ καὶ μεταφυτεύεσθαι. πολύσπερμον δὲ καὶ τὸ ἀβρότονον καὶ οὐκ ἄοσμον, τοῦτο δὲ ῥίζας μεν έχει όρθας καὶ κατά βάθους. ἔστι γὰρ ὥσπερ μονόρριζον τῆ παχεία τὰς δ' ἄλλας <ἀφίησιν> ἀπ' αὐτῆς ὁ δ' ἀμάρακος καὶ ὁ ἔρπυλλος καὶ τὸ σισύμβριον καὶ τὸ έλένιον ἐπιπολαίους καὶ πολυσχιδείς και ταρρώδεις ξυλώδεις δὲ πᾶσαι, πολύ δὲ μᾶλλον ἡ τοῦ ἀβροτόνου καὶ διὰ τὸ μέγεθος καὶ τη ξηρότητι.

1 evlote conj. W.; evlois Ald.

³ i.e. from seed. ταύτην conj. W.; πάντα UMAld.; ? τάντα

4 Plin. 21. 34: cf. C.P. 1. 4. 2. ἀβρότονον . . . θέρους, text nearly as given by Ald. and by UM (?)—supported by Plin.

² Plin. 21. 57. Description of various forms of έρπυλλος has perhaps dropped out after this word: cf. § 5, καθάπερ

ENQUIRY INTO PLANTS, VI. vII. 2-4

wild bergamot-mint, and wild forms of the other plants mentioned, having a more pungent smell. Thyme is sometimes ¹ quite like cultivated thyme.² Now it is plain that these wild forms possess this

means of reproducing themselves.3

Southernwood actually grows more readily from seed than from a root 4 or a piece torn off (though it grows even from seed with difficulty); however it can be propagated by layering in pots in summertime, like the 'gardens of Adonis'; it is indeed very sensitive 6 to cold and generally delicate even where the sun shines brightly; but, when it is established and has grown, it becomes tall and strong and tree-like, like rue, except that the latter is much more woody drier and less succulent.

⁷ Sweet marjoram grows in either way, from pieces torn off or from seed; it produces a quantity of seed, which is fragrant with a delicate scent; it can also be transplanted.⁸ Southernwood also produces much seed, which has some scent. This plant has straight roots which run deep; it has, as it were, its single stout root, from which the others spring; ⁹ while sweet marjoram thyme bergamot-mint and calamint have surface ¹⁰ roots which are much divided and matted; in all these plants the roots are woody, but especially in southernwood, because of its size and because it is so dry.

l.c. so far as that passage is intelligible—but δè before èν οστράκοις add. W.; after èν οστρ. supply βλαστάνει.

⁵ cf. Plat. Phaedo 276 B and Thompson's n. Sir W. Thiselton-Dyer in Companion to Greek Studies, § 99, p. 65.

⁶ cf C.P. 4. 3. 2. ⁷ Plin. 21. 61.

8 μεταφυτεύεσθαι conj. Sch. from G; μεταφύεσθαι Ald.

° αφίησιν add. W.

10 ἐπιπολαίους conj. Scal.; ἐπὶ πολλούς MAld. cf. C.P. 2. 16. 5.

Τοῦ δὲ ἐρπύλλου ἴδιος ἡ αὕξησις ἡ τῶν βλαστῶν δύναται γὰρ ἐφ' ὁσονοῦν προϊέναι κατὰ μῆκος χάρακα λαβὼν ἡ πρὸς αἰμασιὰν φυτευθεὶς ἡ κάτω καθιέμενος εὐαυξέστατος δὲ εἰς φρέαρ. εἴδη δὲ τοῦ μὲν ἡμέρου λαβεῖν οὐκ ἔστι, καθάπερ ἐλέχθη. τοῦ δὲ ἀγρίου φασὶν εἶναι. τοῦ γὰρ ἐν τοῖς ὄρεσιν τὸν μὲν θυμβρώδη τινὰ καὶ δριμύτατον τὸν δ' εὔοσμον εἶναι καὶ μαλακώτερον.

"Ωρα δὲ τῆς φυτείας πλείστων μετόπωρον, ἐν ῷ σπεύδουσιν ὡς πρῶτα φυτεύειν οὐ μὴν ἀλλ' ἔνια καὶ τοῦ ἦρος φυτεύουσιν. ἄπαντα φιλόσκια καὶ φίλυδρα καὶ φιλόκοπρα μάλιστα αὐχμὸν δὲ δέχεται καὶ ὅλως ὀλιγοϋδρότατος ὁ ἔρπυλλος. κόπρφ δὲ χαίρει, μάλιστα δὲ καὶ τῆ τῶν λοφούρων φασὶ δὲ καὶ μεταφυτεύειν δεῖν πολλάκις καλλίω γάρ. τὸ δὲ σισύμβριον, ὥσπερ ἐλέχθη, καὶ ἐξίσταται μὴ μεταφυτευόμενον.

VIII. Τῶν δ ἀνθῶν τὸ μὲν πρῶτον ἐκφαίνεται τὸ λευκόῖον, ὅπου μὲν ὁ ἀὴρ μαλακώτερος εὐθὺς τοῦ χειμῶνος, ὅπου δὲ σκληρότερος ὕστερον, ἐνιαχοῦ τοῦ ἦρος. ἄμα δὲ τῷ ἔφ ἡ μικρόν τι ὕστερον καὶ τὸ φλόγινον καλούμενον τὸ ἄγριον·

⁸ Plin. 21. 61.

 $^{^{1}}$ cf. Plin, 20. 245 and 246 (not from T.); C.P. 2. 18. 2; Diosc. 3. 38; Index ξρπυλλος.

 $^{^2}$ cf. Plin. 19. 172, which refers however to σισύμβριον; Nic. ap. Athen. 15. 31.

ENQUIRY INTO PLANTS, VI. vii. 5-viii. 1

¹The growth of the shoots of thyme is peculiar. If it has a stake, or is planted against a wall, it can send them out to any length; so also if it is let grow downwards; indeed it is most vigorous when grown into a pit.² It is not possible to distinguish different forms of the cultivated kind, as has been said, but they say that of the wild kind (Attic thyme) there is more than one form; for that of the kind which grows on the mountains one form is like savory and very pungent, while the other is fragrant and more delicate.

⁸ The season for planting most of these is autumn, and then men hasten to plant them as early as possible; however some are planted also in spring. All of them love shade, water, and especially dung; however thyme is patient of drought and, in general, needs moisture less than the others. These plants especially delight in the dung of beasts of burden; and it is said that they should often be transplanted, for that it improves them, while bergamot-mint, as has been said, actually degenerates ⁵ if it is not transplanted.

Of the seasons at which coronary plants flower, and of the length of their life.

VIII. ⁶ Of the flowers the ⁷ first to appear is the gilliflower; where the air is mild, it appears as soon as winter comes, but, where it is more severe, later, sometimes in spring. Along with the gilliflower, or a little later, appears the flower called the wild wall-

⁷ τὸ conj. Scal.; τοῦ Ald.

49

E

φιλόσκια conj. Scal. from G; φιλοίκια UMAld. cf. Plin. l.c.
 ἐξίσταται conj. Scal. from G, degenerat; ἐξήτασται MAld.

⁶ Plin. 21. 64-66; Athen. 15. 26 and 27. ἀνθῶν: ? in the sense of ἀνθικῶν, as in 6. 6. 3.

ταθτα γὰρ ὧν οἱ στεφανήπλοκοι χρῶνται πολὺ έκτρέχει τῶν ἄλλων. μετὰ δὲ ταῦτα ὁ νάρκισσος και τὸ λείριον, <και τῶν ἀγρίων ἀνεμώνης γένος τὸ καλούμενον ὅρειον,> καὶ τὸ τοῦ Βολβοῦ κώδυον ἐμπλέκουσι γὰρ ἔνιοι καὶ τοῦτο είς τους στεφάνους. ἐπὶ δὲ τούτοις ἡ οἰνάνθη καὶ τὸ μέλαν ἴον καὶ τῶν ἀγρίων ὅ τε ἐλειόγρυσος καὶ τῆς ἀνεμώνης ἡ λειμωνία καλουμένη και τὸ ξίφιον και υάκινθος και σχεδον 2 όσοις άλλοις χρώνται των όρείων. τὸ δὲ ῥόδον ύστερεί τούτων καὶ τελευταίον μέν φαίνεται, πρώτον δ' ἀπολείπει των ἐαρινών ολιγογρονία γάρ ή ἄνθησις. ὀλιγοχρόνια δὲ καὶ τῶν ἀγοίων τὰ λοιπὰ πλην της ὑακίνθου και της ἀγρίας καὶ τῆς σπαρτῆς αὕτη δὲ διαμένει καὶ τὸ λευκὸν ἴον καὶ ἔτι πλείω τὸ φλόγινον· τὸ δὲ δὴ μέλαν ἴον, ώσπερ είρηται, δι' ενιαυτοῦ θεραπείας τυγχάνον. ώσαύτως δὲ καὶ ἡ οἰνάνθη, καὶ γὰρ τοῦτο ἀνθικὸν μὲν ποῶδες δὲ τὴν φύσιν, ἐάν τις ἀποκνίζη καὶ ἀφαιρῆ τὸ ἄνθος καὶ μὴ ἐᾳ σπερματοῦσθαι καὶ ἔτι τόπον εὔειλον ἔχη· τὸ δὲ ἄνθος βοτρυῶδες καὶ λευκὸν καθάπερ τῶν ἀγρίων . . . ταῦτα μὲν οὖν ὥσπερ ἐαρινὰ φαίνεται.

Τὰ δὲ θερινὰ μᾶλλον η τε λυχνὶς καὶ τὸ διόσανθος καὶ τὸ κρίνον καὶ τὸ ἴφυον καὶ ὁ

² καl τῶν . . . ὄρειον ins Sch. from Athen. l.c. with

alteration of opelwe to applied. cf. Plin. l.c.

⁴ cf. 9. 19. 3. ⁵ See Index.

¹ Evidently both distinct from the νάρκισσος ή λείριον of 6. 6. 9; 6. 8 3. See Index.

³ i.e. the flower of muscari, mentioned in this way because elsewhere (e.g. 7. 12. 1) the edible root is in question, which was properly called $\beta o \lambda \beta \delta s$.

ENQUIRY INTO PLANTS, VI. vIII. 1-3

flower. These, of all the flowers that the garlandmakers use, far outrun the others. After these come pheasant's eye1 and polyanthus1 narcissus (and, among wild plants, the kind of anemone which is called the 'mountain anemone')2 and the 'head'3 of pursetassels: for this too some interweave in their garlands. After these come dropwort violet, and of wild plants, gold-flower,4 the meadow kind of anemone corn-flag hyakinthos (squill), and pretty well all the mountain flowers that are used. The rose comes last of these, and is the first of the spring flowers to come to an end, as it is the first to appear, for its time of blooming is short. So too is that of the rest of the wild plants mentioned, except hyakinthos,5 the wild kind (squill), and also the cultivated (larkspur); this lasts on, and so does the gilliflower, and for a still longer time the wallflower, while the violet, as has been said, blooms throughout the year, if it receives tendance. So too dropwort 7 (for that too is one of the plants valued for their flowers, though it is herbaceous 8 in character) if one pinches off and removes the flower instead of letting it go to seed, and if, further,9 it has a sunny position. The flower is clustering and white, like that of the wild 10 These then are, we may say, the plants of spring.

¹¹ The following belong rather to summer: rose-campion carnation *krinon* ¹² (lily) spike-layender and

⁶ 6. 6. 2; cf. C.P. 1. 13. 12. ⁷ cf. 6. 6. 11.

ποωδες: sense not obvious; εὐωδες conj. Dalec. cf. C.P.
 13. 12.

 ⁹
 ^ℓτι conj. W.; δτι UMAld.

¹⁰ Ut labruscae G, perhaps a guess : see οἰνάνθη in Index.

¹¹ Plin. 21, 67 and 68.

¹² κρίνον Sch. from Athen. l.c.; so also Plin, l.c.; κήρινθον Ald.

άμάρακος ὁ Φρύγιος. ἔτι δὲ ὁ πόθος καλούμενος. ούτος δ' έστι διττός, ό μεν έχων το άνθος δμοιον τη ὑακίνθω, ὁ δὲ ἔτερος ἄχρους λευκός, ώ χρώνται περί τους τάφους και χρονιώτερος ούτος. ανθεί δὲ καὶ ή Ιρις τοῦ θέρους καὶ τὸ στρούθιον καλούμενον τῆ μὲν ὄψει καλὸν τὸ ἄνθος ἄοσμον δέ. μετοπώρου δὲ τὸ λείριον τὸ ἔτερον καὶ ὁ κρόκος, ο τε όρεινος ἄσσμος και ο ημερος εὐθὺς γαρ ανθούσι τοίς πρώτοις ύδασι. χρώνται δε καί τῶν ἀγρίων τῷ τῆς ὀξυακάνθου καρπῷ καὶ τῷ άνθει τῷ τῆς μίλακος.

Καὶ ταῖς μὲν ὥραις οὕτως ἐκάστων ἡ γένεσις. ώς δὲ ἀπλῶς εἰπεῖν οὐδεὶς διαλείπεται χρόνος οὐδ' ἔστιν ἀνανθής, ἀλλὰ καὶ ὁ χειμών ἔχει καίπερ άγονος δοκών είναι διὰ τὴν κατάψυξιν τών μετοπωρινών μεταλαμβανόντων, έαν δε δη καί μαλακὸς ή, πολλώ μᾶλλον. άπλως γὰρ πάντ' ή τὰ πολλά καὶ ἐπεκτείνεται τῆς οἰκείας ώρας, καὶ έὰν ὁ τόπος εὔειλος ἢ μᾶλλον δι' δ καὶ συνέγεια χρόνοι μέν οθν οθτοι καὶ ώραι κατά γίνεται. τας γενέσεις.

Βίος δὲ ἰωνίας μὲν τῆς λευκῆς ἔτη μάλιστα τρία γηράσκουσα δὲ ἐλαττοῦται καὶ ἴα λευκότερα φέρει. ροδωνίας δὲ πέντε τὰ πρὸς τὴν άκμὴν μὴ ἐπικαομένης· χείρω δὲ καὶ ταύτης τὰ ῥόδα γηρασκούσης. πρὸς εὐοσμίαν δὲ καὶ ῥόδων καὶ ἴων καὶ τῶν ἄλλων ἀνθῶν μέγιστον ὁ τόπος

¹ cf. C.P. 1. 4. 1.

3 λευκός. έκλευκος, 'whitish,' Athen. l.c.

² cf. the Eng. plant-name 'love-in-absence'; see πόθος in Index.

⁴ Evidently the νάρκισσος ή λείριον of 6. 6. 9; cf. 6. 8. 1 n.

ENQUIRY INTO PLANTS, VI. vIII. 3-5

the Phrygian sweet marjoram 1; also the plant called 'regret,' 2 of which there are two kinds, one with a flower like that of larkspur, the other not coloured but white, 3 which is used at funerals; and this one lasts longer. The iris also blooms in summer, and the plant called soap-wort, which has a beautiful flower but is scentless. In autumn bloom the other kind of narcissus, 4 the crocus, both the scentless mountain form and the cultivated one (saffroncrocus); for these bloom directly the first rains come. The fruit 5 of the cotoneaster and the flower of the smilax, both of them wild plants, are also used in garlands.

Such are the seasons at which each appears; and, to speak generally, there is no interval of time nor flowerless period, but even winter produces flowers, for all that it seems to be unproductive by reason of the cold, since the autumn flowers continue into winter, and to a much greater extent if the season be mild. For all things, one may say, or at least most of them, extend beyond their proper season, and all the more if the place be sunny; so that there is a continuous succession. These then are the periods and seasons at which the various flowers are produced.

⁷The life of the gilliflower is at most three years; as it ages it degenerates and produces paler flowers.⁸ A rose-bush lives five years, after which its prime ⁹ is past, unless it is pruned by burning ¹⁰; with this plant too the flowers become inferior as it ages. Position and a suitable climate contribute most to

καρπφ: Plin. l,c. apparently read ἄνθει.

⁶ πάντ' η conj. St.; πάντη Ald. H. ⁷ Plin. 21. 69.

^{8 7}a conj. St.; del Ald.

⁹ ἀκμὴν conj. Scal.; ἀκτὴν Ald. ¹⁰ cf. 6. 6. 6.

συμβάλλεται καὶ ὁ ἀὴρ πρὸς ἔκαστον οἰκεῖος. έν Αἰγύπτω γὰρ τὰ μὲν ἄλλα πάντ' ἄοσμα καὶ άνθη καὶ ἀρώματα, αἱ δὲ μυρρίναι θαυμασταὶ τῆ εὐοσμία. προτερεῖν δέ φασι τῶν ἐνταῦθα καὶ ρόδα καὶ ἴα καὶ τὰ ἄλλα ἄνθη καὶ διμήνω, καὶ διαμένειν πλείω των παρ' ήμιν ή οὐκ ελάττω χρόνον ταθτα.

Δοκεί δὲ πολύ πρὸς εὐοσμίαν διαφέρειν, ὥσπερ έλέχθη, καὶ ὁ ἐνιαυτὸς τοῖος ἡ τοῖος γενόμενος, ού μόνον έπομβρίαις καὶ αὐχμοῖς ἀλλὰ καὶ τῶ κατά καιρόν γίνεσθαι καὶ ύδατα καὶ πνεύματα καὶ άπλῶς τὰς τοῦ ἀέρος μεταβολάς. τὰ δὲ ἐν τοῖς ὄρεσιν ώς άπλως εἰπεῖν καὶ ρόδα καὶ ἴα καὶ τὰ άλλα <καλώς μὲν ἀνθεῖν> τη δὲ ὀσμή πολλὰ χείρω γίνεσθαι. καὶ περὶ μὲν τῶν στεφανωματικών καὶ άπλώς των φρυγανικών σχεδον έν τούτοις και τοις όμοίοις έστιν ή ίστορία.

² Plin. l.c.; cf. C.P. 6. 18. 3.

4 Plin. 15, 37,

^{1 ?&#}x27;violets and gilliflowers; so also below.

³ άνθη conj. St. from G; άνανθη Ald. cf. C.P. 6. 19. 4.

ENQUIRY INTO PLANTS, VI. vIII. 5-6

the fragrance of roses gilliflowers¹ and other flowers. Thus in Egypt,² while all other flowers³ and sweet herbs are scentless, the myrtles⁴ are marvellously fragrant. In that country it is said that the roses gilliflowers and other flowers are as much as two months ahead of those in our country, and also that they⁵ last a longer, or at least not a shorter, time than those of our country.

And, as has been said, the particular season according to its character, makes a great difference to the fragrance, not only by reason of rains and droughts, but also according as rain, wind, and in general, the changes of climate occur or do not occur at the fitting moment. Also it appears that in general roses gilliflowers and the rest bloom well on the mountains, but many of them have there an inferior scent.⁶ Concerning coronary plants and under-shrubs in general these examples and others like them suffice for our enquiry.

⁵ ταῦτα conj. W.; τούτου Ald.

 $^{^6}$ ἄνθη τῆ ὀσμῆ πολλῷ Ald.; ἄνθη τῆ δὲ ὀσμῆ πολλὰ UM, whence Sch. and W. conj. that some such words as καλῶς μέν have dropped out and ἀνθεῖν has been altered to ἄνθη. cf. C.P. 6. 20. 1.

BOOK VII

Н

Ι. 'Επόμενον δὲ τοῖς εἰρημένοις περὶ τῶν ποιωδῶν εἰπεῖν· τοῦτο γάρ ἐστι λοιπὸν τῶν ἐξ ἀρχῆς διαιρεθέντων γενῶν, ἐν ῷ συμπεριλαμβάνονταί πως τὸ λαχανηρὸν καὶ τὸ σιτῶδες. καὶ πρῶτον περὶ τοῦ λαχανώδους λεκτέον ἀρξαμένους ἀπὸ τῶν ἡμέρων, ἐπεὶ γνώριμα μᾶλλον τυγχάνει τῶν

ἀγρίων.

Είσὶ δὴ τρεῖς ἄροτοι πάντων τῶν κηπευομένων, ἐν οἶς ἔκαστα σπείρουσι διαιροῦντες ταῖς ὥραις. εἶς μὲν οὖν ὁ χειμερινός, ἄλλος δὲ ὁ θερινός, τρίτος δὲ ὁ μεταξὺ τούτων μεθ' ἡλίου τροπὰς χειμερινάς. καλοῦσι δ' οὕτως οὐ πρὸς τὴν σπορὰν βλέποντες ἀλλὰ πρὸς τὴν γένεσιν καὶ τὴν χρείαν ἐκάστου· ἐπεὶ ἥ γε σπορὰ σχεδὸν ἐν τοῖς ἐναντίοις γίνεται. τοῦ χειμερινοῦ μὲν γὰρ ἀρχὴ μετὰ τροπὰς θερινὰς τοῦ Μεταγειτνιῶνος μηνός, ἐν ῷ σπείρουσι ῥάφανον ῥαφανίδα γογγυλίδα καὶ τὰ καλούμενα ἐπίσπορα· ταῦτα δ' ἐστὶ τεύτλιον θριδακίνη εὕζωμον λάπαθον νᾶπυ κορίαννον ἄνηθον κάρδαμον· καλοῦσι δὲ

¹ cf. C, P, 3, 20, 7 and 8.

BOOK VII

OF HERBACEOUS PLANTS, OTHER THAN CORONARY PLANTS:
POT-HERBS AND SIMILAR WILD HERBS.

Of the times of sowing and of germination of pot-herbs.

I. Next we have to tell of herbaceous plants: for this class remains of those which we distinguished at the outset, and it includes to some extent the classes of pot-herbs and of cereals. And first we must speak of the class of pot-herbs, beginning with the cultivated kinds, since it happens that these are better known than the wild kinds.

¹ There are three seed-times for all things grown in gardens, at which men sow the various herbs, distinguishing by the season. One is the 'winter' seed-time, another the 'summer,' and the third is that which falls between these, coming after the winter solstice. These terms however are given in regard not to the sowing, but to the growth and use of each kind; for the actual sowing takes place, one might almost say, at the opposite seasons. Thus, the 'winter' period begins after the summer ² solstice in the month Metageitnion,³ in which they sow cabbage radish turnip, and what are called 'secondary crops,' that is to say, beet lettuce rocket monk's rhubarb mustard coriander dill cress; and

* July. & before M. om. Sch.

² θ eρινàs conj. Scal.; χ ειμερινàs U(?)MP₂Ald.G (ed. Bas. and Par. but not ed. Tarv.).

καὶ πρώτον τοῦτον τών ἀρότων. τοῦ δὲ δευτέρου πάλιν μεθ' ἡλίου τροπὰς τοῦ Γαμηλιώνος μηνός, ἐν ῷ σπείρουσι καὶ πηγνύουσι πράσον σέλινον γήθυον ἀδράφαξυν. τοῦ τρίτου δέ, δν καλοῦσι θερινόν, τοῦ Μουνυχιώνος· ἐν τούτφ δὲ σπείρεται σίκυος κολοκύντη βλίτον ἄκιμον ἀνδράχνη θύμβρον. ποιοῦνται δὲ πλείους ἀρότους τῶν ὁμοίων καθ' ἐκάστην ὥραν, οἶον ἡαφανίδος ἀκίμου τῶν ἄλλων. πᾶσι δὲ σπείρεται τοῖς

άρότοις τὰ ἐπίσπορα.

Διαφύεται δ' οὐκ ἐν ἴσοις πάντα χρόνοις, ἀλλὰ τὰ μὲν θᾶττον τὰ δὲ βραδύτερον όσα δυσφυή. τάχιστα μεν ουν ὤκιμον καὶ βλίτον καὶ εύζωμον καὶ τῶν χειμερινῶν ραφανίς τριταῖα γὰρ ώς είπειν. θριδακίναι δέ τεταρταίαι ή πεμπταΐαι. σίκυος δὲ καὶ κολοκύντη περὶ τὰς πέντε ή έξ, οἱ δέ φασιν ἐπτά· πρότερον δὲ καὶ θᾶττον ο σίκυος. ἀνδράχνη δ' ἐν πλείοσι τούτων. θον δὲ τεταρταῖον. κάρδαμον δὲ καὶ νᾶπυ πεμπταία. τεύτλιον δε θέρους μεν έκταιον χειμώνος δὲ δεκαταίον. ἀδράφαξυς δὲ ὀγδοαία. ῥάφανος δὲ δεκαταία. πράσον δὲ καὶ γήθυον οὐκ ἐν ἴσοις, ἀλλὰ τὸ μὲν ἐννεακαιδεκαταῖον ἐνιαχοῦ δὲ εἰκοσταῖον, γήθυον δὲ δεκαταῖον ἡ δωδεκαταίον. κορίαννον δε δυσφυές οὐδε γάρ εθέλει βλαστάνειν τὸ νέον ἐὰν μὴ βρεχθῆ. θύμβρα δὲ καὶ ὀρίγανος ἐν πλείοσιν ἡ τριάκοντα. δυσφυέστατον δὲ πάντων τὸ σέλινον τεσσαρακοσταίον γάρ φασιν οί τὰ συντομώτερα λέγοντες,

60

January.
 April.
 Plin. 19. 117.
 τῶν χειμερινῶν: cf. 7. 1. 1.

ENQUIRY INTO PLANTS, VII. 1. 2-3

this is also called the 'first' period of cultivation. The second period begins after the winter solstice in the month Gamelion, in which they scatter or plant the seed of leeks celery long onion orach. The third period, which is called the 'summer' period, begins in the month Munychion : in this are sown cucumber gourd blite basil purslane savory. Moreover they make several sowings of the same herb at each season, as of radish basil and the others. And at all the periods are sown the

'secondary crops.'

8 Not all herbs germinate within the same time, but some are quicker, others slower, namely those which germinate with difficulty. The speediest are basil blite rocket, and of those sown for winter 4 use, radish; for these germinate in about three days. Lettuce takes four or five, cucumber and gourd about five or six, or, as some say, seven; however, cucumber is earlier and quicker than the others. Purslane takes a longer time, dill four days, cress and mustard five. Beet in summer takes six days, in winter ten, orach takes eight, and turnip ten. Leek 5 and long onion do not take the same time, but the former nineteen to twenty days, the latter ten to twelve. Coriander germinates with difficulty; indeed fresh seed will not come up at all unless it is moistened.⁶ Savory ⁷ and marjoram take more than thirty days; but celery germinates with the greatest difficulty of all: for those who make the time comparatively short say forty days, and others fifty, and

⁵ πράσον conj. Bod.; πράσιον P₂Ald. H.

⁷ cf. C.P. 4. 3. 1; Plin. 19. 7.

 $^{^{6}}$ βρεχθή conj. Bod. cf. C.P. 4. 3. 1; ἐλιχθή Ald.; ἐλιχθή P_2 Bas.; so also G.

οί δὲ πεντηκοσταίον, καὶ τοῦτο κατὰ πάντας τούς ἀρότους: ἐπισπείρουσι γάρ τινες ἐπὶ πᾶσιν.

"Ολως δὲ ὅσα κατὰ πλείους ώρας σπείρεται, ταῦτ' οὐδὲν θᾶττον τέλεια γίνεται τοῦ θέρους. καὶ θαυμαστὸν εἰ καὶ μηθὲν ἡ ώρα συμβάλλεται καὶ ὁ ἀὴρ πρὸς τὸ θᾶττον, ἐὰν δὲ μοχθηρὰ καὶ ψυγρά καὶ τῶ ἀέρι περισκεπης βραδύτερον ἐπεὶ καὶ γειμώνων ή εὐδιῶν ἐπιγινομένων τοῖς ἀρότοις ότὲ μὲν βραδύτερον ότὲ δὲ θᾶττον ή βλάστησις. διαφέρει δὲ ταῦτα κατὰ τοὺς ἀρότους ἐκάστων: πρωϊαίτατον γάρ έν τοῖς εὐείλοις καὶ εὐκρᾶσιν.

'Ως γὰρ ἀπλῶς εἰπεῖν ἐν πλείοσι δεῖ τὰς αἰτίας ύπολαβείν των τοιούτων, έν τε τοίς σπέρμασιν αὐτοῖς καὶ ἐν τῆ χώρα καὶ τῷ ἀέρι καὶ ταῖς ὥραις αίς εκαστα σπείρουσι και χειμώνων και εὐδιῶν. άλλὰ τοῦτο μὲν σκεπτέον, ἐφ' ὧν τε παραλλάττουσιν οί χρόνοι καὶ ἐφ' ὧν οὖ καὶ γὰρ τὴν ραφανίδα φασί τινες τριταίαν καὶ θέρους καὶ χειμώνος, τὸ δὲ τεύτλιον, ὥσπερ εἴρηται, παραλλάττει κατὰ τὰς ὥρας. χρόνοι δ' οὖν οὖτοι τῆς βλαστήσεώς είσι καὶ λέγονται καθ' έκαστον.

Διαφέρει δὲ πρὸς τὸ θᾶττον καὶ βραδύτερον καὶ ή τῶν σπερμάτων παλαιότης. τὰ μὲν γὰρ άπὸ νέων παραγίνεται θᾶττον, οἶον πράσον γήθυον

Δρας Vo. H.; χώρας UM; so also G.
 τέλεια conj. W. (comm.); γε πολλά MSS.; τὰ πολλά Vo. Sch. W. (text); γίνεται conj. Sch. from G; γίνεσθαι Ald. 3 και τη άέρι . . . Βραδύτερον: grammar doubtful and text perhaps defective: so given in UM; και δ άηρ περισκεπής 62

ENQUIRY INTO PLANTS, VII. 1. 3-6

that too, at whichever period it is sown, for some sow it as a 'secondary crop' at all the periods.

Generally speaking, those herbs which are sown at more than one season 1 do not mature 2 faster in the summer. Howbeit it is strange if the season and the state of the atmosphere do not contribute at all to quicker growth, and if, when there is an unfavourable cold season and the atmosphere is cloudy. these conditions do not tend to make growth slower, s seeing that, when stormy or fair weather follows the sowing, germination is slower or quicker accordingly. And there is another thing which makes a difference as to the raising of the various herbs; germination begins earlier in sunny places which have an even temperature.

As a matter of fact, to speak roundly, the causes of such differences must be found in several different circumstances, in the seeds themselves, in the ground. in the state of the atmosphere, and in the season at which each is sown, according as it is stormy or fair. However it is a point for consideration with which herbs the time of sowing makes a difference and with which it makes none; thus it is said that radish germinates on the third day whether it be sown in summer or in winter, while beet, as has been said, behaves differently according to the season. Anyway such are and are said to be the seasons of germination in each case.

Another thing which makes a difference as to the rapidity with which the seeds germinate is their age; for some herbs come up quicker from fresh seed, as

Digitized by GOOGLE

πρός το βραδύτερον conj. Sch. (with μοχθ. κ. ψυχρά supply ή ἄρα ή).
4 Plin. 19. 118. δè conj. Scal.; γàρ Ald.H.

σίκυος κολοκύντη ένιοι δὲ καὶ προβρέχουσι τὸν σίκυον πρὸς τὸ θᾶττον ἡ ἐν γάλακτι ἡ ἐν ὕδατι. τὰ δ΄ ἀπὸ παλαιῶν, οἶον σέλινον τεύτλιον κάρ-δαμον θύμβρα κορίαννον ὀρίγανον εἴπερ μὴ <φυτεύεται> αὐτὰ ἀπὸ τοῦ νέου, καθάπερ εἴπομεν. ἴδιον δέ φασιν ἐπὶ τοῦ τευτλίου συμβαίνειν οὐ γὰρ διαφύεσθαι πᾶν εὐθὺς ἀλλ' ὕστερον πολλῷ, τὸ δὲ καὶ τῷ ἐχομένῳ ἔτει καὶ τῷ τρίτῳ, δι' δ καὶ

έκ πολλοῦ σπέρματος ολίγον βλαστάνειν.

Έκαστον δὲ τῶν σπερμάτων, ἐὰν ἀδρυνθέντα ἀποπέση, διαμένει πρὸς τὴν ὥραν τὴν ἑαυτοῦ καὶ οὐ πρότερον ἐκβλαστάνει· καὶ κατὰ λόγον ἐστί· καὶ γὰρ ἐπὶ τῶν ἀγρίων ὁρῶμεν συμβαῖνον, ἐὰν μὴ φθαρἢ. αἱ δὲ τελειώσεις τῶν καρπῶν ἀπάντων γίνονται τοῦ θέρους, πρότερον δὲ καὶ θᾶττον ὡς ἀπλῶς εἰπεῖν τῶν πρότερον σπαρέντων. διαφέρει δὲ καὶ ἡ ὥρα· τὰ γὰρ ἐν ταῖς θερμημερίαις σπαρέντα θᾶττον ἐκκαυλεῖ καὶ ἐκσπερματοῦται, καθάπερ ῥαφανὶς γογγυλίς. ἔνια δὲ οὐκ ἐνιαύσια φέρει τὸν καρπὸν ἀλλὰ δίενα, καθάπερ σέλινον πράσον γήθυον, ἃ καὶ διαμένει χρόνον πλείονα καὶ οὐκ ἔστιν ἐπέτεια· τὰ γὰρ πολλὰ τούτων ἄμα τῆ τελειώσει τῶν σπερμάτων αὐαίνεται.

Πάντα δὲ ὡς εἰπεῖν ὅσα ἐκκαυλεῖ καὶ τελειοῖ τὸν καρπὸν ἀποτελειοῦται κατὰ τὸ σχῆμα τοῦ παραβλαστήσεις ἐκ τῶν καυλῶν ἔχειν ἀκρεμονικάς, πλὴν ὅσα μονόκαυλα, καθάπερ πράσον

καὶ γήθυον καὶ κρόμυον καὶ σκόροδον.

Φίλυδρα δὲ καὶ φιλόκοπρα πάντα, μᾶλλον δὲ

² Sc. soaking.

¹ φυτεύεται αὐτὰ conj. W.; οὐ τὸ UMAld.

ENQUIRY INTO PLANTS, VII. 1. 6-8

leek long onion cucumber gourd; (some even soak the seed of cucumber first in milk or water, to make it germinate quicker). Some come up quicker from old seed, as celery beet cress savory coriander marjoram (unless indeed they are raised ¹ from fresh seed in the manner ² which we have mentioned). There is, they say, a singular feature about beet ³; the seed does not all germinate at once, but some of it not for some time, some even in the next or in the third year; wherefore it is said that little comes up from much seed.

Any of the seeds, if they are ripe when they fall, last till their own proper season and do not sprout till then. And in this they are consistent; for we note that the same thing happens with the seed of wild plants, unless it is destroyed. However all mature their fruits in the summer, though sooner and quicker, generally speaking, when they are sown earlier. The season also 4 makes a difference; things sown in the hot season push up their shoots and go to seed sooner, as radish and turnip. Some however bear their fruit not in the same year but in the next, as celery 5 leek long onion, which plants also last a longer time, and are not annual; for most herbs wither with the ripening of their seed.

Generally speaking, all those that push up shoots and mature their fruit reach their perfection of form in having side-shoots branching from the main stem—except those which have but a single stem, as leek long onion onion garlic.

All these herbs are lovers of water and of dung,

65

³ cf. C.P. 4. 3. 2; Plin. l.c.

δè conj. W.; γὰρ Ald.H.

⁵ Plin. *l.c.*

τὰ ἀσθενέστερα καὶ πλείονος ἐπιμελείας δεόμενα. τὰ δὲ καὶ τροφής.

ΙΙ. Φύεται δὲ πάντα ἀπὸ τοῦ σπέρματος, ἔνια δὲ καὶ ἀπὸ παρασπάδος καὶ κλωνὸς καὶ ῥίζης. άπὸ μὲν παρασπάδος ἡ ῥάφανος δεῖ γάρ τι καὶ οιζώδες προσλαβείν. ἀπὸ δὲ τῶν βλαστῶν πήγανον δρίγανος ὤκιμον ἀποφυτεύουσι γὰρ καὶ τοῦτο ὅταν σπιθαμιαῖον ἡ μεῖζον γένηται τεμόντες είς τὸ ημισυ. ἀπὸ ρίζης δὲ σκόροδον καὶ κρόμυον καὶ βολβὸς καὶ ἄρον καὶ άπλῶς τὰ τοιαῦτα τῶν κεφαλορρίζων. φύεται δὲ καὶ εἴ τινων αἱ ρίζαι διαμένουσιν έπλ πλείονα χρόνον έπετειοκαύλων όντων. ότι δε άπο σπέρματος πάντα βλαστάνει φανερόν καὶ γὰρ τὸ πήγανον, ὅπερ οὔ φασί τινες, άλλα Βραδέως, δι' δ καὶ ἀποφυτεύουσιν.

"Όσα δὲ ἀπὸ ρίζης Φύεται, τούτων ή μὲν ρίζα γρόνιος αὐτὰ δὲ ἐπετειόκαυλα, δι' δ καὶ παραβλαστάνουσιν αι ρίζαι των τοιούτων και γίνονται πλείους οὐ μόνον ἐν τοῖς ἡμέροις καὶ κηπευομένοις άλλα και έν τοις αγρίοις, ώσπερ είπομεν, οίον βολβοίς γηθύοις σκίλλαις καὶ τοίς άλλοις. παραβλαστάνει δ' ένια καὶ τῶν μὴ κεφαλορρίζων γρονιωτέρων δέ, οίον σέλινον καὶ τεύτλιον ἀφιᾶσι γαρ ρίζας αφ' ων φύονται φύλλα και καυλοί.

βλαστών corresponds to κλωνδς above.

Plin. 19. 121.
 ² cf. C.P. 1. 4. 2.
 δεῖ γάρ τι UP₂; ἀεὶ γάρ τι Ald. H.G; Sch. suggests δὲ for γàρ, missing the sense.

ENQUIRY INTO PLANTS, VII. 1. 8-11. 2

and especially the weaker ones, which require more attention or in some cases more feeding,

Of the propagation of pot herbs, and of differences in their roots.

II. ¹All these herbs are propagated from seed, and some also by a piece torn off, a shoot, or a piece of root. Cabbage is propagated by a piece torn off.2 since it is essential 3 in this case to take a piece which has root attached to it. From cuttings 4 are grown rue marjoram basil; for slips of this too men plant when it has grown to the height of a span or more, cutting off half the plant. By root are planted garlic onion purse-tassels cuckoo-pint and in general such bulbous plants. Such propagation is also possible in cases where the roots persist for more than a year, though the shoots last but for a year. And it is plain that all these herbs can be grown from seed; for even rue can (which some deny), though the process is slow, and so cuttings are also taken.

Of those which are propagated by a piece of root the root is long-lived, though the plant itself may be annual; wherefore the roots of such plants make offsets and so increase; and this is true not only of plants cultivated in the garden, but also of wild plants, as we have said, for instance of purse-tassels long onion 7 squill and so forth. Some plants even which are not bulbous 8 but longer-lived make offsets, as celery and beet; for these send out roots from which grow leaves and stems. Long onion and

⁵ cf. C.P. 1. 4. 3. 6 i.e. offsets.

γηθύοις om. some editors, as not being wild.
 i.e. and so annual.

παραβλαστάνει δὲ καὶ γήθυον καὶ πράσον καὶ παραφύει κάτωθεν οίον βολβώδη τινά κεφαλήν, έξ ής ή βλάστησις γίνεται τῶν φύλλων, αὐανθέντος δε τοῦ καυλοῦ καὶ τοῦ σπέρματος ἀφαιρεθέντος άλλα δια το μη χρησίμας είναι τας τούτων κεφαλάς οὐ συλλέγουσιν εἰς ξηρασίαν, δι' 3 δ καὶ οὐ φυτεύουσι. τάχα δὲ ταῦτα καὶ ὁμογενῆ καὶ σύνεγγύς πως τη τοῦ κρομύου φύσει, δι' δ καὶ οὐ θαυμαστόν. ἀλλ' ὁμοίως [καὶ] ἐπὶ πάντων καὶ ἡμέρων καὶ ἀγρίων, ὅσα χρονιώτερα μέν έστιν έπετειόκαυλα δέ, τούτων και αι ρίζαι έπιβλαστάνουσιν, ώσπερ καὶ ἐπὶ τῶν Φρυγανικῶν καὶ τῶν θαμνωδῶν ἀλλ' ἐπὶ τῶν κρομύων καὶ σκορόδων καὶ βολβῶν καὶ ὥσπερ ἀριθμός τις γίνεται τούτων. ή δη γένεσις, ώσπερ είρηται, τριχῶς ἐστιν, ἀπὸ σπέρματος μὲν πάντων, ἀπὸ δὲ καύλοῦ καὶ ρίζης τῶν εἰρημένων.

Τῶν δὲ καυλῶν κολουσθέντων πάντα μὲν ὡς εἰπεῖν βλαστάνει πλὴν τῶν ἀποκαύλων, ἐμφανέστατα δ' ὥσπερ καὶ εἰς χρείαν ὤκιμον θρίδαξ ράφανος. καὶ τῆς μὲν θρίδακος ἡδίους φασὶ τοὺς παλιμβλαστεῖς εἶναι καυλούς τὸν γὰρ πρῶτον ὀπώδη καὶ πικρὸν εἶναι ὡς ἄπεπτον οἱ δὲ τὸ ἐναντίον ὀπωδεστέρους τούτους ἀλλ' ἔως ἃν ὧσιν ἀπαλοὶ φαίνεσθαι γλυκυτέρους. ἀλλ' ἐπὶ τῆς

¹ πράσυν conj. St.; πράσιον Ald. H.

² διὰ τὸ μη conj. W.; μη διὰ τὸ UM(?) Ald.

³ i.e. offset bulbs.

⁴ W. omits μέν (Ald. UM(?)) after συλλέγουσι.

i.e. the plant is increased by seed only and not by offsets.
 of. 7. 4. 10; Plin. 19. 103.

⁶ δμοίως conj. Sch.; δμως PAld. H. (UM?).

⁷ επετειόκαυλα conj. Sch.; επιγειότερα PAld.H.

ENQUIRY INTO PLANTS, VII. 11. 2-4

leek 1 also make offsets, sending out a 'head' below, like the bulb of purse-tassels, from which the leaves spring; but this only takes place when the stem has withered and the seed has been removed. But. as 2 the 'heads' 3 of such plants are not useful, they do not collect them 4 for storing dry; wherefore also they do not plant these.5 It may be that somehow these are akin and closely allied to onion, wherefore what has been said is not surprising. However in all those plants, both wild and cultivated alike,6 which have an annual stem,7 but yet live longer than a year, there is an outgrowth of the roots, just as there is in under-shrubs and shrubby plants: while in onions garlic and purse-tassels even a number,8 as it were, of such roots is formed. In fact, they are reproduced in three ways, as has been said; from seed in all cases and from the stem 10 and root in those specified.

¹¹ Almost all shoot again if the stem is broken (except those which are stemless), but most obviously basil lettuce cabbage, which are, as it were, broken for a practical reason. Indeed they say that the stems of lettuce which thus grow again are sweeter, ¹² for that the original stem has a taste like fig-juice and is bitter, as being not properly ripened. Some however say that the later stems have the taste of fig-juice more than the original one, but that, so long as they are tender, they appear sweeter. Be that as

⁹ δη conj. Sch.; δè Ald.

69

δριθμός is clearly corrupt, and has displaced an unusual word for which ὅσπερ apologises.

¹⁰ καυλοῦ is here that part of the plant which is above ground.

11 Plin. 19. 122.

¹² ήδίους Vo.mBas.H., so too G, Plin. l.c., Athen. 2. 69; iblous UAld. of. C.P. 2. 15. 6.

ραφάνου τοῦτο όμολογούμενον, ώς εἰ πάλιν βλαστήσειεν ἡδίων ἀφαιρεθέντων γε τῶν φύλλων πρὸ τοῦ διακαυλίσαι.

Διαμένουσι δὲ αἱ ῥίζαι πλειόνων, ἀλλ' αἱ μὲν βλαστάνουσι πάλιν αἱ δὲ οὔ. ῥαφανὶς γοῦν καὶ γογγυλὶς διαμένουσι γῆς ἐπιβληθείσης ἄχρι θέρους καὶ αὔξησιν λαμβάνουσιν, ὅπερ ποιοῦσί τινες ἐξεπίτηδες τῶν κηπουρῶν· οὐ βλαστάνουσι δὲ οὐδ' ἀμιᾶσι φύλλον οὐδ' εἴ τις ἀφέλοι τὴν ἐπισεσαγμένην γῆν. ἰδεῖν δὲ τοῦτο καὶ ἐπὶ τῶν ἄλλων ἐστί. τὰ δὲ πλεῖστα τῶν λαχάνων μονόρριζα τῆ παχεία κατὰ βάθους ῥίζη· καὶ γὰρ ὅσα παραφύει τὰς ἰσοπαχεῖς ταύτας, ὥσπερ σέλινον καὶ τεύτλιον, ἀπὸ τῆς μέσης πως ἡ παράφυσίς ἐστι καὶ οὐκ εὐθὺς ἀπὸ τῆς ἀρχῆς ἡ σχίσις· ἐκ δὲ ταύτης τῆς μιᾶς ἀπήρτηνται αἰ ἀποφυάδες αἱ μικραὶ καὶ τῆς ῥαφανίδος καὶ τῆς γογγυλίδος. καὶ αὖται μὲν δὴ πᾶσι φανεραὶ διὰ τὴν χρείαν.

Ή δὲ τοῦ τευτλίου μία μὲν μακρὰ καὶ παχεῖα καὶ ὀρθή, καθάπερ ἡ τῶν ῥαφανίδων, ἀποφύσεις δὲ ἔχει παχείας ὁτὲ μὲν δύο ὁτὲ δὲ καὶ τρεῖς ὁτὲ δὲ καὶ μίαν, τὰς δὲ μικρὰς ἐκ τούτων. σαρκώδης δὲ ἡ ῥίζα καὶ τῆ γεύσει γλυκεῖα καὶ ἡδεῖα, δι ὁ καὶ ἀμὴν ἐσθίουσί τινες ὁ δὲ φλοιὸς οὐ παχὺς οὐδὲ ἀφαιρετός, ὥσπερ ὁ τῶν ῥαφανίδων, ἀλλὰ μᾶλλον οἰος ὁ τῶν ἱπποσελίνων. ὡσαύτως δὲ καὶ ἡ τῆς ἀδραφάξυος μία μὲν εἰς βάθος ἐκ

ταύτης δὲ ἄλλαι.

1 βλαστήσειεν conj. Sch.; βλαστήσει Ald.

 ² οὐδ' εἴ τις Ald. H.; εἰ μή τις conj. Scal. supported by G.
 ³ ἐκ... μικραὶ conj. W.; εἰς δὲ ταύτην τὴν μίαν ἡ ἀπ' αὐτῆς

ENQUIRY INTO PLANTS, VII. II. 4-6

it may, it is admitted that in the case of cabbage the stem is sweeter if it should have grown 1 again after being broken, provided that the leaves are stripped

off before the plant runs to stalk.

In most cases the roots persist, but they do not in all cases produce fresh growth. Thus radish and turnip persist till summer, if earth is thrown on them, and they increase in size; and some gardeners do this deliberately; but they do not make fresh growth nor send out leaves, even if one 2 removes the earth heaped over them. And this may also be observed in other plants. However, most pot-herbs have the single stout root which runs deep; for even in those which produce these side-roots of equal stoutness, as celery and beet, the side-growth comes, as it were, from the middle root and it is not separate to start with; but to this single root are attached the small out-growths,8 both in radish and in turnip. These instances are familiar to all because of the use 4 which is made of these plants.

The beet has a single long stout straight root like that of the radish, and has stout out-growths, sometimes two, sometimes three, sometimes only one, and the small ones are attached to these. The root is fleshy and sweet and pleasant to the taste, wherefore some even eat it raw. The 'bark' is not thick and cannot be detached, like that of the radish, but rather resembles that of alexanders. In like manner the root of orach is single and runs deep, and other

roots are attached to it.

τε καl τῆς ἀποφυάδος καl μικρὰ Ald. H.; so also M, omitting τε. W.'s restoration of a very corrupt text is at least consistent with what follows in $\S 6$.

4 i.e. for food.

Μονορριζότατον δὲ τούτων πάντων τὸ λάπαθον οὐ γὰρ ἔχει παχείας ἀποφύσεις ἀλλὰ τὰς λεπτάς βαθυρριζότατον δὲ πάντων, ἔχει γὰρ μείζω τριῶν ἡμιποδίων τὸ δ' ἄγριον βραχυτέραν, πολύκαυλον δὲ καὶ πολύκλαδον καὶ ἡ ὅλη μορφὴ τελειωθεῖσα παραπλησία τῆ τοῦ τευτλίου πολυχρονιώτερον δὲ καὶ τοῦ ἀγρίου καὶ ὅλως δὲ πάντων τῶν λαχάνων ὡς εἰπεῖν διαμένει γὰρ ὁποσονοῦν χρόνον ὡς φασιν. ἔχει δὲ σαρκώδη τὴν ρίζαν καὶ ἔνικμον, δι' δ καὶ ἐξαιρεθεῖσα ζῆ πολὺν χρόνον.

Τὸ δ΄ ὥκιμον μίαν μὲν τὴν παχεῖαν τὴν κατὰ βάθους τὰς δ' ἄλλας τὰς ἐκ πλαγίου λεπτὰς

έπιεικώς εὐμήκεις.

"Ενια δ' οὐκ ἔχει τὴν μίαν τὴν ὀρθήν, οἶον τὸ βλίτον, ἀλλ' εὐθὺ πολλὰς έξ ἄκρου καὶ εὐπαχεῖς

καὶ μακροτέρας τῆς ἀδραφάξυος.

Τῶν δὲ ριζῶν ξυλωδέσταται πασῶν αἱ τοῦ ἀκίμου, καθάπερ καὶ ὁ καυλός. ἡ γὰρ τοῦ βλίτου καὶ τῆς ἀδραφάξυος καὶ τῶν τοιούτων ἦττον ξυλώδης. εἰσὶ γὰρ ὡς ἀπλῶς εἰπεῖν πασῶν αἱ μὲν σαρκώδεις αἱ δὲ ξυλώδεις. <σαρκώδεις>, οἶον ἡ τοῦ τευτλίου καὶ τοῦ σελίνου καὶ ἱπποσελίνου καὶ λαπάθου καὶ ἡαφανίδος καὶ γογγυλίδος καὶ πάντων μάλιστα τῶν κεφαλοβαρῶν οὐδὲ γὰρ ἀναξηραινόμεναι σκληρύνονται τελείως. ξυλώδεις δέ,

3 cf. 7. 6. 1; C.P. 3. 1. 4. See Index.

 ² τὰs Ald., cf. τὰs δὲ μικρὰs § 6; τινας conj. W. cf. Plin. 19.
 98 (who mistranslates).

ENQUIRY INTO PLANTS, VII. 11. 7-8

Monk's rhubarb 1 however has a single root in a truer sense than any of the others, for it has no stout out-growths of root, but only the 2 slender ones; its root also runs deeper than that of the others, being more than a foot and a half long. The wild sort 3 however has a shorter root, and has several stems and branches, and its shape, as a whole, when fully grown resembles that of beet. Cultivated monk's rhubarb moreover is longer lived than the wild form, 4 and, in general, we may say, than any other pot-herb, for, they say, it may live any time. It has a fleshy root, 5 full of moisture, wherefore, if pulled up, it will live some time.

Basil has the single stout root, the one which runs deep, and the others at the sides are slender and

fairly long.

Some herbs, as blite, have not the single straight root, but a number of roots which start directly from the top and are of a good stoutness⁶ and longer than

those of orach.

The roots of basil are woodier than those of any of the other herbs, as also is its stem; for those of blite orach and the like are less woody. In general we may say that the roots of any of these herbs are either woody or fleshy. Examples of fleshy roots are beet celery alexanders monk's rhubarb radish turnip, and especially all heavy-headed hinds, for the roots of these do not wither up altogether even when they are dried. Examples of those with woody roots

δίζαν conj. Sch.; σάρκα Ald.

⁷ πασῶν conj. W.; παρ' ὧν UMP; also Ald. H., omitting ai.

8 σαρκώδειs add. Scal. from G.

9 i.e. bulbous; cf. 1. 6. 8.

⁶ Plin. l.c. seems to have read a different word from εὐπαχεῖs, or to have misunderstood it.

ώσπερ αί τοῦ ἀκίμου καὶ βλίτου καὶ ἀδραφάξυος καὶ εὐζώμου καὶ ἀνήθου [καὶ λαπάθου] καὶ κοριάννου καὶ ἀπλῶς τῶν νευροκαύλων· ἔχει γὰρ δὴ καὶ τὸ ἄνηθον καὶ τὸ κορίαννον ὄντα μονόρριζα ξυλώδη τε τὴν ῥίζαν καὶ οὐ μακρὰν οὐδὲ τὰς λεπτὰς ἀποφυάδας ἔχουσαν πολλάς· πολύκαυλα δὲ ἄμφω καὶ πολύοζα, δι' δ καὶ οὐ κατὰ λόγον οὐδενὶ τούτων τὸ ἄνω πρὸς τὸ κάτω.

Βραχύρριζα δὲ ταῦτά ἐστιν, οἶον θρῖδαξ ἀνδράχνη, τῆ ὀρθῆ καὶ ταῖς εἰς τὰ πλάγια. ἡ δὲ
θρῖδαξ, ὥσπερ οὐκ ἔχει τὰς τοιαύτας ἀποφύσεις
ἀλλὰ μόνον τὰς λεπτάς, καὶ μάλιστα δὴ μονόρριζον ὡς εἰπεῖν. ἀπλῶς δὴ πάντα τὰ θερινὰ
βραχύρριζα· καὶ γὰρ ὁ σίκυος καὶ ἡ κολοκύντη
καὶ ἡ σικύα καὶ διὰ τὴν ὥραν καὶ ἴσως ἔτι μᾶλλον
διὰ τὴν φύσιν, ἤπερ συνηκολούθηκε τῆ ὥρα. ἡ δὲ
μεταφυτευομένη θρῖδαξ βραχυτέραν ἔχει τὴν
ρίζαν τῆς σπαρείσης· παραβλαστάνει γὰρ ἐκ τῶν
πλαγίων μᾶλλον· βραχυτέραν δὲ καὶ ἡ ἀγρία τῆς
ἡμέρου, καὶ ἐκ τῶν ἄνωθεν πολυκαυλοτερα.

ΙΙΙ. 'Ανθεί δὲ τῶν μὲν ἄλλων ἔκαστον ἀθρόον, τὸ δὲ ἄκιμον κατὰ μέρος, τὰ κάτω πρῶτον εἶτ' ὅταν ταῦτα ἀπανθήση τὰ ἄνω, δι' ὁ καὶ πολυ-

² ἀποφυάδας conj. Scal.; ἀποφυλλάδας Ald.

ταῦτα conj. Sch.; τὰ τοιαῦτα UM; τοιαῦτα Ald.
 Athen. 2. 79. Sch suggests that the name of a plant has dropped out after ωσπερ: ? ἡ ἀνδράχνη.

¹ After ἀνήθου Ald. H. have καὶ λαπάθου: bracketed by W. after Sch.

ENQUIRY INTO PLANTS, VII. II. 8-III. I

are basil blite orach rocket dill 1 coriander, and in general, those with fibrous stems; for in dill and coriander, which have a single root, the root is woody and not long, and the slender side-roots from it are not numerous; but both plants have several stems and branches; wherefore in neither of these plants does the part above ground correspond to the part which is below.

The following 3 have short roots: lettuce and purslane, in which both the straight main root and the side ones are short. 4 Lettuce may be said to have no such side-roots, but only the slender ones, and may be called in the strictest sense a plant of a single root. In general all summer herbs have short roots: we may include cucumber gourd and bottlegourd, both because of the season to which they belong and perhaps still more because of their character, which corresponds to the season. ever the transplanted lettuce has a shorter root than one that is raised from seed, since it is more apt to send out side-growths; also the wild kind has a shorter 5 root than the cultivated, and the part above ground has more stems.6

Of the flowers and fruits of pot-herbs.

III. 7All, except one, of these herbs produce all their bloom at once, but basil has a succession of flowers. the lower part of the plant flowering first, and then, when that bloom is over, the upper part. Wherefore its season of bloom is a long one, like that of the

⁵ βραχυτέραν conj. Sch.; βραχυτέρα Ald.

 δ ἄνωθεν πολυκαυλοτέρα conj Sch. from G; ἄνω· τὰ δὲ πολυκ.
 Id. of. Diosc. 2. 136,
 Plin. 19. 100. Ald. cf. Diosc. 2. 136,

χρόνιον ἐν τῷ ἀνθεῖν, καθάπερ κύαμος καὶ τῆς πόας τὸ ἡλιοτρόπιον καλούμενον καὶ ἄλλα δὲ τῶν ἀγρίων. ἀνθεῖ δὲ καὶ ὁ σίκυος πολὺν χρόνον καὶ γὰρ ἐπιβλαστάνειν τούτῷ γε συμβαίνει. τὰ δὲ ἄνθη τῶν μὲν ἔκλευκα τῶν δὲ μηλινοειδῆ τῶν δὲ μικρὸν ἐπιπορφυρίζοντα, εὕχρουν δ' οὐθέν.

Τὰ δὲ σπέρματα διαφέρει καὶ τοῖς σχήμασι τὰ μὲν γὰρ πλεῖστα στρογγύλα τὰ δὲ προμήκη τὰ δ᾽ αὖ πλατέα καὶ ψυλλώδη, καθάπερ τὰ τῆς ἀδραφάξυος ὅμοιον γὰρ τῷ τοῦ σιλφίου τὰ δὲ στενὰ καὶ γραμμώδη, καθάπερ τοῦ κυμίνου. καὶ τοῖς χρώμασιν ὁμοίως, τὰ μὲν μέλανα τὰ δὲ ξυλώδη τὰ δὲ λευκότερα. πάντα δὴ ἐλλοβοσπέρματα ἡ παπποσπέρματα ἡ ἀμφλοιοσπέρματα ἡ παπποσπέρματα ἡ άμφλοιοσπέρματα, καὶ γογγυλὶς ἐλλοβοσπέρματα, κορίαννον δὲ καὶ μάραθον καὶ ἄνηθον καὶ κύμινον γυμνοσπέρματα, βλίτον δὲ καὶ τεύτλιον καὶ ἀδράφαξυς καὶ ἄκιμον ἐμφλοιοσπέρματα, θριδακίνη δὲ παπποσπέρματον.

Πάντα δὲ πολύκαρπα καὶ πολυβλαστῆ, πολυκαρπότατον δὲ τὸ κύμινον. ἔδιον δὲ καὶ δ λέγουσι κατὰ τούτου· φασὶ γὰρ δεῖν καταρᾶσθαί τε καὶ βλασφημεῖν σπείροντας, εἰ μέλλει καλὸν ἔσεσθαι

καὶ πολύ.

Δυσξήραντα δὲ πάντα μὲν ὡς εἰπεῖν πλὴν τοῦ κυμίνου, οὐχ ὡς ὁ σῖτος οὖτος γὰρ κὰν ἄπαξ

3 μηλινοειδέs: cf. 6. 2. 8.

¹ For the collective sense of πόα (= τὰ ποώδη) cf 1. 3. 1.
² πολὺν χρόνον conj. W., which at least gives the required sense; καλούμενος Ald.

^{? &#}x27;orange.' 5 Plin. 19. 119.

ENQUIRY INTO PLANTS, VII. III. 1-3

bean, and among herbaceous plants 1 that of the plant called *heliotropion*, and also other wild plants. Cucumber also has a long period 2 of bloom, for this plant has a second growth. The flowers are in some cases whitish, in others quince-yellow, 3 in others somewhat reddish 4; but the flower is never of a

bright colour.

⁵The seeds too differ in shape; most are round, but some are oblong; some again are broad and leaf-like, as those of orach, for the seed of this is like that of silphium; others again are narrow or marked in lines, as those of cummin. They also vary in colour, some being black, some the colour of wood, some paler. The seeds of all are either in pods or naked, or have an integument or have a pappus. Radish mustard and turnip have their seeds in pods; coriander fennel dill and cummin have naked seeds; those of blite beet orach and basil are enclosed in an integument; those of lettuce have a pappus on them.

All have numerous fruits and numerous shoots, but cummin has the most ⁸ fruits of all. ⁹ And there is another peculiarity told of this plant: they say that one must curse and abuse it, while sowing, if

the crop is to be fair and abundant.

Nearly all of these, except cummin, are hard to dry for keeping,—unlike corn 10; for this, when once

⁷? 'brown' cf. 7 9. 3. • cf. 8. 3. 5; C. P. 4. 15. 2.

⁹ cf 9. 8. 8; Plin. l.c. applies this to ωκιμον, Pall. 4. 9. 5

10 σίτος· οδτος γάρ I conj.; σίτος γάρ UMH.; P omits γάρ; σίτος δε W. after Sch.; nec modo frumenti consistunt, quod G.

^{*} γραμμώδη: cf 4. 12. 2.; canaliculata Plin. l.c.

άδρυνθή ταχὺ ξηραίνεται καὶ ἀποπίπτει· δυσξηραντότερα δὲ τὰ ἐμφλοιοσπέρματα καὶ τούτων 4 <μάλιστα τὸ ὤκιμον. ἄπαντα δὲ ξηρανθέντα πολυκαρπότερα γίνεται, δι' δ> καὶ προαφαιροῦντες αὐτὰ ξηραίνουσιν. ἄπαντα δὲ πολύχοα καὶ πολυσπέρματα, πολυκαρπότατον δὲ τὸ

ὥκιμον.

Έστι δὲ τὰ μὲν ἀκρόκαρπα, καθάπερ ὤκιμον πράσον κρόμυον· τὰ δὲ πλαγιόκαρπα μᾶλλον, οἰον ῥαφανὶς γογγυλὶς καὶ τὰ τοιαῦτα· τὰ δ᾽ ἀμφοτέρως, οἰον βλίτον ἀδράφαξυς· ἀμφότερα γὰρ ταῦτα καὶ ἐκ τοῦ πλαγίου, καὶ τό γε βλίτον εὐθὺς παρ᾽ ἔκαστον ὄζον προσκαθήμενον ἔχει τὸ σπέρμα βοτρυῶδες. τὰ δ᾽ ἐκ παλαιοτέρων σπερμάτων θᾶττον ἐκκαυλεῖ, τάχιστα δὲ τὰ ἐκ τῶν ἀκμαζόντων· ἔστι γάρ τις ἀκμὴ καὶ τούτων. ἀνὰ λόγον δὲ καὶ τὸ κάλλος ἀκολουθεῖ τῶν . . . ἐὰν τὰ ἄλλα τὴν αὐτὴν ἔχωσι θεραπείαν.

Δοκεῖ δὲ καὶ εἰς το αὐτο ἀθρόα θεμένων καλλίω γίνεσθαι καὶ βλαστάνειν οὕτω γὰρ το τοῦ πράσου καὶ τὸ τοῦ σελίνου τιθέασιν ἀποδήσαντες

είς δθόνιον καὶ γίνονται μεγάλα.

Συμβάλλεται δέ τι καὶ ὁ τόπος πρὸς αὔξησιν· κελεύουσι γοῦν, ὅταν τις μεταφυτεύη τὰ σέλινα, πάτταλον κατακρούειν ἡλίκον ἃν βούληται ποιεῖν τὸ σέλινον· τιθέναι δὲ καὶ ἐν ὀθονίφ πάτταλον κατακρούσαντα καὶ πλήσαντα κόπρου καὶ γῆς.

² τό γε βλίτυν conj. W.; τό γε πλείστον U; τό τε πλείστοι Ald.H.

¹ μάλιστα . . . δι' δ missing in UMAld. Bas.; text as restored by Sch. from Cam., G and Plin. l.c.
² τό γε βλίτον conj. W.; τό γε πλεῖστον U; τό τε πλεῖστον

³ ἐκκαυλεῖ: cf. 7. 1. 7; 7. 4. 3, and esp. C.P. 4. 3. 5.

After ἀκολουθεῖ τῶν follows a lacuna ôf one and a half lines
 78

ENQUIRY INTO PLANTS, VII. 111, 3-5

it is ripened, quickly dries and is shed, and the herbs whose seed have an integument are harder still to dry. especially basil. All however, when dried, produce more fruit: wherefore 1 it is the custom to gather the seed early and dry it. All of them are prolific and produce many seeds, but basil produces most of all.

Examples of those which produce their fruit at the top of the stem are basil leek onion: of those which produce it rather at the sides, radish turnip and the like; of those which produce it in both ways, blite and orach; both of these produce it at the side as well as at the top; in fact blite 2 has its seed in clusters, closely attached to each branch. Some push up their shoots 3 fairly soon from old seed, but seed from plants in their prime is the most rapid; for these plants too have a time when they are at their best. The beauty of the plant also corresponds 4 in proportion, provided that equal care in 5 other respects is shewn in cultivation.

⁶ It likewise appears that, if a quantity of seed is sown in the same place, the resulting crop comes up and germinates better; thus they tie up seed of leek and celery in a piece of cloth before sowing, and then there is a large 8 crop.

The position also contributes to growth; at least, when celery is transplanted, they suggest that one should hammer 9 in a peg of whatever size one wishes to make the celery; and also that one should sow the seed in a piece of cloth 10 after hammering in a peg and filling the hole with dung and soil.

in UMAld.; text as given by Cam., which however omits τῶν; τῶν σπειρομένων Η.; τῶν τοιούτων Vo. Vin.

5 cf. 7. 4. 7. 6 Plin. 19. 120. 7 cf. C.P. 5. 6. 9.

⁸ μεγάλα conj. St.; μεγάλαι Ald. H.

⁹ Made clearer C.P. 5. 6. 7. 10 cf. C.P. 5, 6, 9,

Ενια δε καὶ τοῖς σχήμασιν εξομοιοῦται καὶ τοῖς τόποις· ἡ γὰρ σικύα ὁμοιοσχήμων γίνεται εν & ἄν τεθῆ ἀγγείω.

Καὶ διαφορὰν λαμβάνει κατὰ τοὺς χυμοὺς ἔνια προθεραπευθέντα τῶν σπερμάτων, οἱον τὸ τοῦ σικύου ἐὰν ἐν γάλακτι βρέξαντες σπείρωσιν. ἀλλὰ τὰ μὲν τοιαῦτα ἴσως οἰκειότερα τῆς θεραπείας.

ΙΥ. Γένη δὲ τῶν μέν ἐστι πλείω τῶν δ' οὐκ ἔστιν, οἶον ἀκίμου λαπάθου βλίτου καρδάμου εὐζώμου ἀδραφάξυος κοριάννου ἀνήθου πηγάνου τούτων γὰρ οὔ φασιν εἶναι <γένους διαφοράν.> τῶν δὲ ἔστι, ῥαφανίδος ῥαφάνου τευτλίου σικύου κολοκύντης κυμίνου σκορόδου θριδακίνης. διαιροῦσι δὲ τοῖς τε φύλλοις καὶ ταῖς ῥίζαις καὶ τοῖς χρώμασι καὶ τοῖς χυλοῖς καὶ τοῖς ἄλλοις τοῖς τοιούτοις.

Οῖον τῆς ἡαφανίδος < γένη Κορινθίαν Κλεωναίαν Λειοθασίαν> ἀμωρέαν Βοιωτίαν· εὐαυξεστάτην δὲ τὴν Κορινθίαν, ἡ καὶ τὴν ῥίζαν ἔχει γυμνήν· ἀθεῖται γὰρ εἰς τὸ ἄνω καὶ οὐχ ὡς αὶ ἄλλαι κάτω. τὴν δὲ Λειοθασίαν, ἡν ἔνιοι καλοῦσι Θρακίαν,

 $^{^{1}}$ kal toîs tómois Ald.; katà toùs tómous conj W. cf. C.P. 5. 6. 7.

 ² ἀγγείφ... λαμβάνει om. UMPAld.; διαφορὰν δὲ καὶ Cam.; τόπφ· διαφέρειν δὲ καὶ H.; ἀγγείφ conj. W. from C.P. 5. 6. 7; καὶ διαφορὰν conj. Sch. cf. Geop. 12. 19. 6.
 ³ cf. 7. 1. 6; Geop. 12. 20. 3.

After είναι there is a lacuna in UMAld.; Cam. supplies γένους διαφοράν τῶν δὲ ἀνάπαλιν πλείω γένη; Η. has πλείω γένη 80

ENQUIRY INTO PLANTS, VII. III. 5-IV.

Some things again come to resemble in their shape even the position 1 in which they grow: thus the bottle-gourd becomes like in shape to the vessel 2 in which it has been placed.

Moreover differences in taste are acquired in some cases when the seed has been treated specially beforehand; for instance, the seed of the cucumber produces a fruit with different taste if it is soaked 3 in milk before sowing. But such matters belong perhaps more properly to the subject of cultivation.

Of the various forms of some pot-herbs.

IV. Of some herbs there are several kinds, but of others only one, as basil monk's rhubarb blite cress rocket orach coriander dill rue; of each of these they say that there is 4 but one kind. But of others there is more than one, as radish cabbage beet cucumber gourd cummin garlic lettuce. Differences are marked in the leaves, the root, the colour, the taste, and so forth.

Thus of radish they recognise these various kinds ⁵—the Corinthian, that of Cleonae, the Leiothasian, amorea, the Boeotian. The Corinthian is said to be the strongest in growth, and it has an exposed root; for it pushes upwards, and not downwards like the others. The Leiothasian ⁶ is called by some the

οὐδὲ γένους διαφοράν· τῶν δὲ ἀνάπαλιν πλείω γένη; Plin. 19. 123 rather supports H. ? read as in H.: τῶν δὲ ἐστι is perhaps an attempt to fill the lacuna.

5 cf. Plin. 19. 75 and 76, who gives a kind called viride in place of T.'s ἀμωρέα: see below. After ραφανίδος there is a lacuna in UMAld. (but U has την δὲ μόραν Βοιωτίαν). Text restored from Athen. 2. 48 (cf. Plin. l.c.). Cam. H. Bas. (also Vo.Vin.(?)) give substantially the same.

⁶ The name suggests Thasos, off the Thracian coast.

ἰσχυροτάτην πρὸς τοὺς χειμῶνας. τὴν δὲ Βοι ωτίαν γλυκυτάτην καὶ τῷ σχήματι στρογγύλην, οὐχ ὥσπερ τὴν Κλεωναίαν μακράν. ὅσων δ΄ αν ἢ λεῖα τὰ φύλλα, γλυκύτεραι καὶ ἡδίους, ὅσων δ΄ αν τραχέα, δριμύτεραι. γένος δέ τι παρὰ ταῦτα ἔστιν δ ἔχει τὸ φύλλον εὐζώμῳ ὅμοιον. ῥαφανίδος μὲν οὖν ταῦτα.

Γογγυλίδος δε οί μέν φασιν είναι οί δ' ού φασιν, άλλὰ τῶ ἄρρενι καὶ τῆ θηλεία διαφέρειν, γίνεσθαι δὲ ἐκ τοῦ αὐτοῦ σπέρματος ἄμφω. πρὸς δὲ τὸ ἀποθηλύνεσθαι πηγνύναι δεῖν μανάς. έὰν γὰρ πυκνάς, πάσας ἀπαρρενοῦσθαι, τὸν αὐτὸν δὲ τρόπον κὰν ἐν γἢ μοχθηρὰ σπαρῶσι δι' δ καὶ πρός σπερματισμόν μεταφέροντες φυτεύουσι τάς έκφύσεις καὶ πλατείας. ἔστι δὲ καὶ τὸ σπέρμα τη όψει τὸ γείρον καὶ βέλτιον φανερόν της μέν γάρ χρηστής λεπτὸν τής δὲ μοχθηράς άδρόν. χειμαζομένη δε χαίρει καλ αύτη καλ ή ραφανίς. οἴονται γὰρ ἄμα γλυκαίνεσθαί τε καὶ τὴν αὔξησιν είς την ρίζαν τρέπεσθαι και οὐκ είς τὰ φύλλα. τοις δε νοτίοις και ταις ευδίαις εκκαυλεί ταχύ. τοῦτο μὲν οὖν λόγου δεῖται τῆς ὁμοιώσεως ἐν άμφοῖν εἶναι τὰς διαφοράς.

¹ Diosc. 2. 112 mentions a kind called by the Romans λρμοράκιον. Plin. 19. 82 has armoracia and says that this was called armon in Pontus; Sch. suggests that the latter name may have given rise to both armoracia and λμωρέα.

Plin. 18. 129, cf. 19. 75; Athen. 9. 7.
 πηγνόναι. The verb is used of planting seeds singly; cf. 6. 6. 9; 7. 1. 2; 7. 5. 3.

ENQUIRY INTO PLANTS, VII. IV. 2-3

Thracian radish, and it stands the winter best. The Boeotian is said to be the sweetest and to be round in shape, not of a long shape like that of Cleonae. Those kinds whose leaves are smooth are sweeter and pleasanter to the taste, those whose leaves are rough have a somewhat sharp taste. Besides the above-mentioned kinds ¹ there is yet another, whose leaves resemble those of rocket. These then are the different kinds of radish.

Of the turnip 2 all do not agree that there are several kinds, but some say that the only difference is between the 'male' and the 'female,' and that both forms come from the same seed. In order to produce 'female' plants it is said that the seed should be sown 8 thinly, for that, if it is sown thick, the result is all 'male' plants; and that the same result follows if the seed is sown in poor soil. Wherefore, when they are shifting plants for seeding.4 they plant the seedlings 5 wide apart.6 Good and inferior seed can be easily distinguished by their appearance; the seed of a good plant is fine. that of a poor one coarse. Both this plant and radish like exposure to winter; for it is supposed that this makes them sweeter and that they are thus made to grow roots rather than leaves. a south wind and warm weather they run up quickly. It needs explanation that both plants should thus adapt 7 themselves in special ways.

83

⁴ πρὸς σπερματισμὸν conj. W.; τοὺς σπερματισμοὺς Ald H. cf. 7. 5. 3. ⁵ ἐκφύσεις: cf. 3. 3. 7.

⁶ καl πλατείας corrupt. διεστηκυίας (W.) gives the required sense; but there may be a loss of some words, πλατείας indicating that the object is to produce broader plants. cf. C.P. 5. 6. 9 and Sch.'s note.

⁷ τῆς δμοιώσεως probably corrupt: no correction suggests itself.

4 Τῆς δὲ ἡαφάνου τριχῆ διαιρουμένης, οὐλοφύλλου τε καὶ λειοφύλλου καὶ τρίτης τῆς ἀγρίας, <ἡ ἀγρία> τὸ μὲν φύλλου ἔχει λεῖον μικρὸν δὲ καὶ περιφερές, πολύκλαδος καὶ πολύφυλλος, ἔτι δὲ χυλὸν ἔχουσα δριμὸν καὶ φαρμακώδη, δι' ὁ καὶ πρὸς τὰς κοιλίας αὐτῷ χρῶνται οἱ ἰατροί. ὁμοίως δὲ καὶ ἐν ἐκείναις δοκοῦσι διαφοραὶ καθ' ἑκατέραν ἐπεὶ ἄσπερμόν τι γένος αὐτῶν ἐστιν ἡ κακόσπερμον. τὸ δ' ὅλον ἡ οὕλη τῆς λείας εὐχυλοτέρα καὶ μεγαλοφυλλοτέρα.

Εὐχυλότερον δὲ καὶ τῶν τευτλίων τὸ λευκὸν τοῦ μέλανος καὶ ὀλιγοσπερμότερον, δ καλοῦσί

τινες Σικελικόν.

'Ωσαύτως δὲ καὶ τῆς θριδακίνης· ἡ γὰρ λευκὴ γλυκυτέρα καὶ ἀπαλωτέρα. γένη δὲ αὐτῆς ἐστὶν ἄλλα τρία, τό τε πλατύκαυλον καὶ στρογγυλόκαυλον καὶ τρίτον τὸ Λακωνικόν· αὕτη δὲ τὸ μὲν φύλλον ἔχει σκολυμῶδες, ὀρθὴ δὲ καὶ εὐαυξὴς καὶ ἀπαράβλαστος ἐκ τοῦ καυλοῦ. τῶν δὲ πλατειῶν οὕτω τινὲς πλατύκαυλοι γίνονται ὥστ' ἐνίους φασὶ καὶ θύραις χρῆσθαι κηπουρικαῖς. τὸ δὲ ὁπῶδες σφόδρα καὶ μικρόφυλλον καὶ λευκοκαυλότερον ἔοικεν ἀγρίᾳ.

Τῶν δὲ σελίνων καὶ ἐν τοῖς φύλλοις καὶ ἐν τοῖς καυλοῖς αἱ διαφοραί· τὸ μὲν γὰρ πυκνὸν καὶ οὖλον καὶ δασὰ τὸ φύλλον ἔχει, τὸ δὲ μανότερον καὶ πλατύτερον καυλὸν δὲ μείζω. τούτων δὲ πάλιν τὰ μὲν λευκόκαυλα τὰ δὲ πορφυρόκαυλα ἡ ποικιλόκαυλα· τὸ δ᾽ ὅλον ἄπαν τὸ τοιοῦτον

έμφερέστερον τῷ ἀγρίφ.

¹ Athen. 9. 9; Plin. 19. 80. ² Wild radish. See Index. 84

ENQUIRY INTO PLANTS, VII. IV. 4-6

¹ Of cabbage three kinds are distinguished, the curly-leaved, the smooth-leaved, and thirdly, the wild form.2 The wild form 3 has a small round leaf, it has many branches and many leaves, and further a sharp medicinal taste; wherefore physicians use it for the stomach. Between the other two kinds 4 there seem also to be differences, inasmuch as one of them bears no seed or only inferior seed. In general the curly-leaved kind has a better flavour than the smooth and it has larger leaves.

⁵ So too with beet; the white kind has a better flavour than the black and produces fewer seeds;

some call it 'Sicilian' beet.

So too with lettuce: the white kind is sweeter and tenderer. Of this plant there are three other kinds,6 the flat-stalked, the round-stalked, and the Laconian: the last-named has a leaf like the golden thistle, but is erect and strong-growing and has no side-shoots 8 from the main stem. Of the 'flat' kinds some have such flat stalks that some, they say, use them to make a garden trellis.9 The third kind, which has much milky juice and small leaves and a whiter stem, is like a wild plant.

10 In celery the differences between the various kinds lie in the leaves and stem; one kind is close and curly and has rough leaves, the other is more open in growth and flatter, but has a larger stalk. Again there are kinds with stems white, red or particoloured; and in general all such forms resemble

more the wild kind.

³ ἡ ἀγρία add. W.

⁴ ἐκείναις conj. Sch. from Plin. l.c.; ἐκείνφ Ald. H.

⁵ Athen. 9. 11; Plin. 19. 132. ⁶ Plin. 19. 125.

⁷ Athen. 2. 79. 8 cf. 7. 2. 4.

9 ostiola olitoria Plin. 19. 125. 10 Plin. 19, 124

Σικύου δὲ καὶ κολοκύντης τοῦ μὲν εἶναί φασι γένη τῆς δ' οὐκ εἶναι, καθάπερ τῆς ῥαφανίδος καὶ τῆς γογγυλίδος, ἀλλ' ἐν τῷ αὐτῷ γένει τὰς μὲν βελτίους τὰς δὲ χείρους. τοῦ δὲ σικύου τρία, Λακωνικὸν σκυταλίαν Βοιώτιον τούτων δὲ ὁ μὲν Λακωνικὸς ὑδρευόμενος βελτίων, οἱ δ' ἔτεροι ἀνύδρευτοι.

Διαφέρει δὲ γένει καὶ τὰ κρόμυα καὶ τὰ σκόροδα. πλείω δὲ τοῦ κρομύου τὰ γένη, οίον τὰ κατὰ τὰς χώρας ἐπικαλούμενα Σάρδια Κνίδια Σαμοθράκια, καὶ πάλιν τὰ σητάνια καὶ σγιστὰ καὶ ᾿Ασκαλώνια. τούτων δὲ τὰ μὲν σητάνια μικρά γλυκέα δὲ εὖ μάλα, τὰ δὲ σχιστὰ καὶ άσκαλώνια καὶ ταῖς θεραπείαις διαφέροντα καὶ δηλον ὅτι τῆ φύσει τὸ γὰρ σχιστὸν τῷ μὲν γειμώνι μετά της κόμης έωσιν άργόν, αμα δέ τω ήρι τὰ φύλλα περιαιροῦσι τὰ ἔξω καὶ τὰ ἄλλα θεραπεύουσι περιαιρεθέντων δὲ τῶν φύλλων έτερα βλαστάνει καὶ άμα κάτω σχίζεται, δι' δ καλοῦσι σγιστά, οἱ δὲ καὶ ὅλως φασὶ πάντων δείν, ὅπως ἡ δύναμις εἰς τὸ κάτω καὶ μὴ σπερ-8 μοφυή. των δε 'Ασκαλωνίων ιδία τις ή φύσις. μόνα γὰρ <ού> σχιστὰ καὶ ὥσπερ ἄγονα ἀπὸ τῆς ρίζης, ἔτι δὲ ἐν αὐτοῖς ἀναυξή καὶ ἀνεπίδοτα·

² Plin. 19. 101-104.

4 i.e. making offsets.

¹ Athen. 3, 4; Plin. 19, 68.

³ Σάρδια conj. Meurs. from Plin. l.c.; γάρδια Ald. H.

⁵ 'Ασκαλόνια, whence Eng. shallot; though this name is applied to κ. σχιστόν.
⁶ τὸ add. W.

ENQUIRY INTO PLANTS, VII. IV. 6-8

As to cucumber and gourd, it is said that there are various forms of the former, but of the latter, just as in radish and turnip, the differences are only between better and inferior individuals. Of the cucumber there are three forms, the Laconian the cudgel-shaped and the Boeotian. Of these the Laconian is better with moisture, the others without it.

² There are also various kinds of onion and of garlic; those of the onion are the more numerous, for instance, those called after their localities Sardian.³ Cnidian, Samothracian; and again the 'annual' the 'divided' (shallot) and that of Ascalon.⁵ Of these the annual kind is small but very sweet, while the divided and the Ascalonian differ plainly as to their character as well as in respect of their cultivation. For the 'divided' 6 kind they leave untended in winter with its foliage,7 but in spring they strip off 8 the outside leaves and tend the plant in other ways; when the leaves are stripped off, others grow, and at the same time division takes place under ground, which is the reason of the name 'divided.'9 Some indeed say that all kinds should be thus treated, in order that the force of the plant may be directed downwards and it may not go to seed. Ascalonian kind has a somewhat peculiar character; it is the only kind which does not 10 divide and which does not, as it were, reproduce itself from the root; moreover in the plant 11 itself there is no power of increasing and multiplying; wherefore

κόμης έωσιν conj. Scal.; κοιμησέως UMP2Ald.
περιαιρούσι conj Scal. from Plin. l.c. and G; περιάγουσι
P2Ald. H. ° cf. Pall. 3. 24. 3.
10 οὐ add. Scal. 11 i.e. the part above ground.

87

δι' δ καὶ οὐ πηγνύουσιν ἀλλὰ σπείρουσιν αὐτὰ καὶ σπείρουσιν ὀψὲ πρὸς τὸ ἔαρ, εἶθ' ὅταν βλαστήση μεταφυτεύουσι τελειοῦται δὲ οὕτω ταχέως ὥσθ' ἄμα τοῖς ἄλλοις ἡ καὶ πρότερον ἔξαιρεῖσθαι πλέονα δὲ χρόνον ἐαθέντα ἐν τῆ γῆ σήπεται φυτευθέντα δὲ καυλὸν ἀφίησι καὶ σπέρμα φύει μόνον, εἶτα κενοῦται καὶ αὐαίνεται. τούτων μὲν οὖν τοιαύτη τις ἡ φύσις.

9 Διαφέρει δ' ένια καὶ τοῖς χρώμασιν έν Ἰσσῷ γὰρ τὰ μὲν ἄλλα ὅμοια τοῖς ἄλλοις, λευκὰ δὲ σφόδρα τη χροιά φέρειν δέ φασιν δμοια τοίς Σαρδιανοίς, ίδιωτάτη δὲ ἡ φύσις ἡ τῶν Κρητικῶν, παραπλησία δὲ τρόπου τινὰ τοῖς 'Ασκαλωνίοις, εἰ μη άρα και ή αὐτή. ἐν Κρήτη γάρ ἐστί τι γένος δ σπειρόμενον μεν ρίζαν ποιεί φυτευόμενον δε καυλον καὶ σπέρμα, κεφαλην δὲ οὐκ ἴσχει, γλυκὺ δε τώ χυμώ τοῦτο γαρ οίον ἀνάπαλιν έχει τοῖς 10 άλλοις. άπαντα γάρ πηγνύμενα καὶ βελτίω καλ θάττον παραγίνεται. πάντα δὲ φυτεύεται μετ' 'Αρκτούρον έτι θερμής ούσης τής γής, όπως τὰ ὕδατα πεφυτευμένα καταλαμβάνη. καὶ ὅλα δὲ φυτεύεται καὶ διατεμνόμενα παρά τὴν κεφαλήν. οὐχ ὅμοιαι δὲ αἱ ἐκβλαστήσεις, ἀλλ' ἐκ μέν τοῦ κάτω γίνεται κρόμυον, έκ δὲ τοῦ ἄνω

2 οὕτω conj. Sch. from G; τοῖς ἄλλοις Ald.
 3 i.e. instead of being raised from seed. cf. what is said

7. 2. 2 of the offsets of γήθυον.

 $^{^{1}}$ πηγνύουσι: cf. 7. 4. 3 n. The word evidently has a different sense here; cf. § 10, where πηγνύω and φυτεύω seem to be synonymous.

ENQUIRY INTO PLANTS, VII. IV. 8-10

many do not plant these, but raise them from seed; and the sowing is made late, towards the spring; and then, when the seed has germinated, they transplant. And the plant arrives at maturity so fast that it is taken up with the others or even earlier; whereas, if it is left a longer time in the ground, it rots. If planted on the other hand, it sends up a stem and merely produces seed, and then shrivels up and withers. Such then is the character of these.

Some also shew differences in colour; thus at Issus 5 are found plants which in other respects resemble the others,6 but which are extremely white in colour; and they bear, it is said, onions like those of Sardis. Most distinct however is the character of the Cretan kind, which resembles to some extent that of Ascalon, if indeed it be not the same. For in Crete there is a kind which when sown produces a root, but when planted produces a stem and seed but has no 'head'; 7 and it is sweet in flavour. This kind in fact has just the contrary character to the others; for they all grow better and faster when they are planted. All are planted 8 after the rising of Arcturus while the earth is still warm, so that the rains may come upon them after planting. They are planted 9 either entire or else in sections made by cutting at the 'head.' The growth which results is not uniform; from the lower part comes an onion,

⁴ κενοῦται conj. St. from G exinaniuntur; καινοῦται Ald. ⁵ Ἰσσφ conj. Sch. from G and Plin. l.c.; ἴσφ UM; νήσφ

 ⁶ άλλοις conj. Sch; λευκοῖς Ald.
 ⁷ Sc. bulb.
 ⁸ φυτεύεται conj. Sch; φύεται Ald.
 See next note.

⁹ φυτεύεται M; φύεται Ald. cf. C.P. 1. 4. 5.

χλόη μόνον· ὀρθὸν δὲ διατμηθὲν ὅλως ἀβλαστές ἐστι. τὸ δὲ γήτειον καλούμενον ἀκέφαλόν τι καὶ ὅσπερ αὐχένα μακρὸν ἔχον, ὅθεν καὶ ἡ βλάστησις ἄκρα· καὶ ἐπικείρεται πολλάκις, ὥσπερ τὸ πράσον, δι' ὁ καὶ σπείρουσιν αὐτὸ καὶ οὐ φυτεύουσι. τὰ μὲν οὖν κρόμυα σχεδὸν ταύτας ἔχει τὰς ἰδέας.

Τὸ δὲ σκόροδον φυτεύεται μὲν μικρὸν πρὸ τροπῶν ἡ μετὰ τροπὰς διαιρούμενον κατὰ γέλγεις. 11 διαφορά δέ έστιν αὐτῶν ή τε τῶν ὀψίων πρὸς τὰ πρώϊα γένος γάρ τι τυγχάνει τοιοῦτον δ έν έξήκοντα ήμέραις τελειοῦται, καὶ μεγέθει καὶ μικρότητι. καὶ τῷ μεγέθει γένος τι διάφορόν έστι, μάλιστα δὲ τὸ Κύπριον καλούμενον τοιοῦτον, όπερ ούχ έψοῦσιν άλλὰ πρὸς τοὺς μυττωτοὺς γρώνται, καὶ ἐν τῆ τρίψει θαυμαστὸν ποιεῖ τὸν όγκον εκπνευματούμενον, καὶ έτι τῷ μὴ έχειν ένια τὰς γέλγεις. ἡ δὲ γλυκύτης καὶ ἡ εὐωδία καὶ ή άδρότης σχεδὸν παρὰ τὰς χώρας γίνεται καὶ τὰς θεραπείας, ὥσπερ καὶ τῶν ἄλλων. τελειοῦται δὲ καὶ ἀπὸ σπέρματος ἀλλὰ βραδέως τῷ πρώτω γαρ έτει κεφαλην ηλίκην πράσου λαμβάνει, τῷ δ' ὕστερον γελγιδοῦται, καὶ τῷ τρίτφ τέλειον γίνεται, και οὐδεν χειρον άλλ' ἔνιοί γε 12 καὶ κάλλιόν φασι τοῦ πηκτοῦ. τῆς δὲ ρίζης ἡ γένεσις οὐχ όμοία τοῦ τε σκορόδου καὶ τοῦ κρομύου άλλα του μέν σκορόδου όταν άνοιδήση ή γελγίς κυρτούται πάσα και ένταύθα αὐξηθείσα διαιρείται πάλιν είς τὰς γέλγεις καὶ έξ ένὸς πολλά γίνεται τῷ τελειοῦσθαι τὴν κεφαλήν, τὸ δὲ

¹ i.e. bulb; cf. 9. 11. 6. ² cf. 7. 2. 2. ³ Plin. 19. 111 and 112.

ENQUIRY INTO PLANTS, VII. IV. 10-12

from the upper only foliage; while, if the plant is divided vertically, no growth at all takes place. The kind called horn-onion has no 'head,' 1 but has as it were a long neck, at the top of which comes the new growth; it is often cut, like the leek; wherefore it is raised from seed and not planted. Such then, one may say, are the forms of the onion.

³ Garlic is planted a little before or after the solstice. when it divides into cloves.4 There are different kinds distinguished as late or early, for there is one kind which matures in 5 sixty days. There are also differences as to size. There is one kind which excels in size, especially that variety which is called Cyprian, which is not cooked but used for salads, and, when it is pounded up, it increases wondrously in bulk, making a foaming dressing. There is a further difference, in that some kinds cannot be divided into The sweetness of taste and smell and the vigour depend on the position 6 and on cultivation, as with other herbs. Garlic reaches maturity from seed, but slowly, for in the first year it acquires a 'head' which is only as large as that of the leek, but in the next year it divides into cloves, and in the third is fully grown, and is not inferior, indeed some say it is superior, to the garlic which has been planted.7 The growth of the root in garlic and onion is not the same; in garlic, when the clove has swollen, the whole of it becomes convex 8; then it increases and divides again into the cloves, and becomes several plants instead of one by the maturing of the 'head,'

5 δ έν conj. Sch.; δθεν UMAld.
6 χώρας conj. Dalec : ἄρας UMP-Ald

⁴ γέλγεις conj. Scal. from G (nucleatim divisum); γέτη Ald.

χώραs conj. Dalec.; ωραs UMP₂Ald.
 Sc. not raised from seed.
 So W. renders.

κρόμυον εὐθὺς ἐκ τῆς ῥίζης ἄλλο καὶ ἄλλο παραφίησι, καθάπερ καὶ βολβοὶ καὶ σκίλλα καὶ πάντα τὰ τοιαῦτα, καὶ γὰρ τὰ κρόμυα καὶ τὰ σκόροδα μη ἀναιρούντων ἀλλ' ἐώντων πολλά γίνεται. φέρειν δέ φασι καὶ τὸ σκόροδον ἐπὶ τῆς φύσιγγος σκόροδα καὶ τὸ κρόμυον κρόμυα περὶ μὲν οὖν τῶν γενέσεων ίκανῶς εἰρήσθω.

V. Φίλυδρα δὲ πάντα τὰ ἄλλα λάγανα καὶ φιλόκοπρα πλην πηγάνου, τοῦτο δὲ ηκιστα φιλόκοπρον. τὰ χειμερινὰ δὲ οὐχ ήττον τῶν θερινῶν καὶ τὰ ἐπίκηρα τῶν ἰσχυρῶν. κόπρον δὲ μάλιστα έπαινούσι την συρματίτιν, την δέ των υποζυγίων μοχθηραν δια το μάλιστα έξικμάζεσθαι. ζητοῦσι δὲ τὴν κόπρον ἄμα τῷ σπόρω μάλιστα συναναμιγθείσαν οί δὲ καὶ σπείροντες ἐπιβάλλουσι χρώνται δὲ καὶ τῆ ἀνθρωπίνη ὤμῆ πρὸς τὴν χύλωσιν. φιλυδρότερα δὲ τὰ χειμερινὰ τῶν θερινών καὶ τὰ ἀσθενή των ἰσχυρών, έτι δὲ τὰ πλείστης δεόμενα τροφής. φίλυδρα καὶ τὸ κρόμυον καὶ τὸ γήθυον καίτοι φασί τινες οὐ ζητείν, εάν τὸ πρώτον επιγένηται δὶς ἡ τρίς. 2 τῶν δὲ ὑδάτων ἄριστα τὰ πότιμα καὶ τὰ ψυχρά, γείριστα δὲ τὰ άλυκὰ καὶ δυσμανῆ, δι' δ καὶ έκ των όχετων ου χρηστά συμπεριφέρει γαρ σπέρματα πόας. ἀγαθὰ δὲ τὰ ἐκ διός ταῦτα

¹ cf. 7. 2. 2 and 3.

² φύσιγγος conj. Casaub. on Athen. 2. 78; σφύριγγος UM Ald. See LS φύσιγξ.

³ και τὸ κρόμυον κρόμυα conj. Sch.; και τὰ κρόμμυα UMAld. 4 Plin. 19, 156.

ENQUIRY INTO PLANTS, VII. IV. 12-V. 2

while the onion puts out another and another growth straight from the root, as do purse-tassels 1 and squill and all such plants. For both onions and garlic multiply if they are not removed but left alone. They say also that garlic produces garlic heads on the stalk, 2 and that the onion in like manner produces onions. 3 Let this suffice for an account of their ways of growth.

Of the cultivation of pot-herbs; manure and water.

V. 4 All the pot-herbs are lovers of water and of dung, except rue, which does not at all like dung; this is true of the winter no less than of the summer herbs, and of the tender no less than of the strong ones. The dung which is most commended is that which is mixed with litter, while that of beasts of burden is held to be bad, because it is most apt to lose its moisture. Dung which is mixed with the seed is most in request, but some cast the manure on while they are sowing, and they also use fresh human dung as a liquid manure.5 The winter crops like moisture more than the summer ones, and the weak more than the strong, as well as those which specially need feeding. Onion and long onion also love moisture, though some say that they do not require it, if at the outset it has been applied twice or thrice. 6 Fresh cold water is the best, and the worst is that which is brackish and thick:7 wherefore the water from irrigation ditches is not good, for it brings with it seeds of weeds. Rain

⁵ Lit. 'for their liquid-manuring.' cf. C.P. 3. 9. 2, where χύλωσιs must have the same sense.

⁶ Plin. 12. 182 and 183,

^{&#}x27; δυσμανή UMAld.; δυσμενή Η.

γὰρ δοκεῖ καὶ φθείρειν τὰ θηρία [γινόμενα] τὰ γόνιμα κατεσθίοντα. φασὶ δέ τινες οὔτε τοῖς σικύοις συμφέρειν οὔτε τοῖς κρομύοις. ἀρδεύουσι δὲ τὰ μὲν ἄλλα πρωὶ ἡ πρὸς ἐσπέραν, ὅπως μὴ καθέψηται, τὸ δὲ ὤκιμον καὶ μεσημβρίας· καὶ γὰρ διαβλαστάνειν θᾶττόν φασι θερμῷ τὸ πρῶτον ἀρδευόμενον. τὸ δὲ πολὺ λίαν ὕδωρ δοκεῖ συμφέρειν ἄλλως τε καὶ ἐὰν [μὴ] ἔχη κόπρον· πολλάκις γὰρ πεινῆν τὰ λάχανά φασι, καὶ ταῦτα γνωρίζειν τοὺς ἐμπείρους τῶν κηπουρῶν.

Μεταφυτευόμενα δὲ πάντα καλλίω καὶ μείζω γίνεται· καὶ γὰρ τὰ τῶν πράσων μεγέθη καὶ τὰ τῶν ῥαφανίδων ἐκ μεταφυτείας. μάλιστα δὲ μεταφυτεύουσι πρὸς τοὺς σπερματισμούς· καὶ τὰ μὲν ἄλλα ὑπομένει, οἶον γήθυον πράσον ῥάφανος σίκυος σέλινον γογγυλὶς θρίδαξ, <τὰ δὲ> γλίσχρως. ἄπαντα δ' εὐαυξέστερα καὶ μείζω πηγνυμένων τῶν σπερμάτων ἡ σπειρομένων.

Θηρία δὲ γίνεται ταῖς μὲν ἡαφανῖσι ψύλλαι, τἢ δὲ ἡαφάνω κάμπαι καὶ σκώληκες, καὶ ἐν τἢ θριδακίνη καὶ ἐν τοῖς πράσοις καὶ ἐν ἄλλοις δὲ πλείοσιν αἱ πρασοκουρίδες. ταύτας μὲν οὖν ἡ κράστις ἀθροισθεῖσα ἀπόλλυσι καὶ ὅταν κόπρος

⁴ Plin. 19. 183.

 $^{^1}$ γινόμενα τὰ γόνιμα H.; γινόμενα γόνιμα UMAld.; ? τὰ τὰ γόνιμα. Either γινόμενα or γόνιμα seems to be due to dittography. For γόνιμα cf. C.P. 1. 15. 1: τὰς γονίμους ἀρχάς. 2 καθέψηται conj. Sch. after Plin. l.c.; καθάψηται P_2 Ald.

³ έχη κόπρου conj. Dalec.; μη έχη κ. Ald.; μετέχη κόπρου conj. W. cf. 7. 5. 1, χύλωσιν; C. P. 3. 9. 2.

ENQUIRY INTO PLANTS, VII. v. 2-4

water is good, for it also appears to destroy the pests which devour the young plants. Some however say that rain-water is not good for melons nor for onions. Most herbs are watered in early morning or at evening, so that they may not be dried up 2; but basil is watered even at noon, for it is said that it grows more quickly if it is watered at first with warm water. In general water seems to be extremely beneficial, especially if it is mixed with dung 3; for, they say, pot-herbs often are hungry, and experienced gardeners can recognise when this is so.

⁴ All herbs grow finer and larger if transplanted; for even the size of leeks and radishes depends on transplantation. Transplanting is done especially in view of collecting seed ⁵: and, while most herbs bear it well, as long onion leek cabbage cucumber celery turnip lettuce, others bear it less well.⁶ All however make better growth and are larger if the

seed is planted 7 rather than scattered.

Of the pests which infest pot-herbs.

⁸ As for pests,—radish is attacked by spiders,⁹ cabbage by caterpillars and grubs, while in lettuce, leek, and many other herbs occur 'leek-cutters.' ¹⁰ These are destroyed by collecting green fodder,¹¹ or when they have been caught somewhere in a mass

⁷ πηγνυμένων: cf. 6. 6. 9; 7. 4. 3. ⁸ Plin. 19. 177.

⁹ ψύλλαι: cf. Arist. H.A. 9. 39. 1.

11 κράστις conj. R. Const.; κρᾶσις Ald.

⁵ σπερματισμούς conj. Scal.; σπερματικούς UMAld. cf. 7. 4. 3.
⁶ τὰ δὲ γλίσχρως conj. Sch., adding τὰ δὲ; γλίσχρως U; γλίσχρος M; γλίσχρος Ald.; γλήχων conj. Scal. Sch. also conjectures τὰ λίσχρα: see LS. s.v.

¹⁰ πρασοκουρίδες: ? leaf-maggots. cf. Arist. H.A. 5. 19. 20; Geop. 12. 9.

άθρόα που καταλάβη φιλόκοπρον δ' δν τὸ θηρίον ἀναδύεται καὶ ἐνδῦσα κοιμᾶται ἐν τῆ κόπρω, δι' δ δὴ ράδιον θηρεύειν ἄλλως δ' οὐκ ἔστι. ταῖς δὲ ραφανῖσι πρὸς τὰς ψύλλας πρόσφορον τὸ ἐπισπείρειν ὀρόβους. πρὸς δὲ τὸ μὴ γίνεσθαι ψύλλας οὕ φασιν εἶναι φάρμακον οὐδέν. ὑπὸ δὲ τὸ ἄστρον ὥκιμον μὲν λευκαίνεται κορίαννον δὲ άλμậ. τὰ μὲν οὖν συμβαίνοντα διὰ τούτων θεωρητέον.

Τῶν δὲ σπερμάτων τὰ μέν ἐστιν ἰσχυρότερα τὰ δὲ ἀσθενέστερα πρὸς διαμονήν ἰσχυρότερα μὲν οἰον κορίαννον τεύτλιον πράσον κάρδαμον νᾶπυ εὕζωμον θύμβρα, ἀπλῶς τὰ δριμέα πάντα ἀσθενέστερα δὲ γήθυον, τοῦτο γὰρ οὐκ ἐθέλει μένειν, ἀδράφαξυς ὤκιμον κολοκύντη σίκυος, ἀπλῶς τὰ θερινὰ τῶν χειμερινῶν μᾶλλον. διαμένει δὲ οὐδὲν πλέον τεττάρων ἐτῶν ὤστε ἔτι χρήσιμον εἶναι πρὸς τοὺς σπόρους ἀλλὰ διένα μὲν βελτίω, τὰ δὲ τριένα οὐδὲν χείρω, τὸ δ᾽ ὑπερτεῖνον ἤδη χεῖρον.

Πρὸς δὲ τὴν μαγειρικὴν χρείαν ἐπὶ πλείω δια-

1 κόπρος άθρόα που καταλάβη Ald.; κόπρον άθρόαν πού τις καταβάλη conj. W. after Sch.; κόπρον άθρόαν conj. Scal.

3 πρός τὰς ψύλλας πρόσφορον το mBas.; ψύλλας πρός το Ald. Η.; πρός τὰς ψύλλας ἀρκεῖ το conj. W.

4 ψύλλας Ald.; καμπάς conj. Sch. followed by W.

⁵ cf. Geop. 12. 7; Pall. 1. 35. 8; Plin. l.c.

96

² φιλόκοπρου δ' δυ τὸ θήριου ἀναδύεται καὶ ἐνδῦσα conj. W.; φιλόπονου τὸ θήριου ἀναδεύεται καὶ ἐν αἰς κοιμᾶται UMAld.; φίλυπνου conj. R. Const., but W.'s conj. is confirmed by Geop. l.c. The change of gender in ἐνδῦσα is strange.

ENQUIRY INTO PLANTS, VII. v. 4-6

of dung, the pest being fond of dung emerges, and, having entered the heap, remains dormant there 2; wherefore it is then easy to catch, which otherwise it is not. To protect 3 radishes against spiders 4 it is of use to sow vetch 5 among the crop; to prevent the spiders from being engendered they say that there is no specific. 6 Basil turns pale about the rising of the dog-star, and coriander becomes mildewed. In these instances we may observe the accidents which occur to pot-herbs.

Of the time for which seed of pot-herbs can be kept.

⁸ Of seeds some have more vitality than others as to keeping; among the more vigorous ones are coriander beet leek cress mustard rocket savory, and in general ⁹ those of pungent taste; among the less vigorous are long onion—which will not keep—orach basil gourd cucumber; and in general the summer herbs keep less well than the winter ones. No seed will keep more than four years so as still to be of use for sowing; though it is better in the second year, ¹⁰ in some cases it does not deteriorate in three years, ¹¹ but after that time ¹² deterioration begins.

However for cooking purposes seed will keep a

97

VOL. II.



н

⁶ Plin. 19, 176.

⁷ άλμα conj. W.; άλμαι MAld.; άλμαίνεται Vo. Vin.; άλμαται mBas. cf. 8. 10. 1; C.P. 6. 10. 5. In all three places W. introduces this word, comparing ψωριῶν ἐρυσιβῶν, etc.

⁸ Plin. 19. 181.

^{*} άπλῶs conj. St. from G; ἄλλωs Ald.; ἄλωs U.

διένα conj. Scal.; δι' ενα UMAld.H.
 τὰ δὲ τριένα conj.W.; διὰ δὲ τρεῖς UMAld.H.

¹² ύπερτείνου conj. Scal.: cf. 8. 11. 5; ὑπὲρ γαῖον UMAld.; ὑπερβαῖνου Η.

μένει, πλην ἀσθενέστερα ταῦτα ἀναγκαῖον εἶναι διά την άναπνοην και την σκωλήκωσιν. Φθορά δὲ μάλιστα μὲν ὑπὸ τῶν θποίων γίγνεται γὰρ ἐν άπασι καὶ τοῖς δριμέσιν, ἥκιστα δὲ ἐν τῶ σικυῶνι· ού μην άλλα και έξικμαζόμενα πικρά γίνεται τη γεύσει, δι' δ καὶ πρὸς τὴν χρείαν χείρω. περί μεν των σπερμάτων και άπλως των κηπευομένων ίκανῶς εἰρήσθω.

VI. Περὶ δὲ τῶν ἀγρίων καὶ τῶν καλουμένων άρουραίων πειρατέον όμοίως είπειν. τυγχάνει δέ τὰ μὲν ὁμώνυμα τοῖς ἡμέροις ἄπαντα γάρ ἐστι τὰ γένη ταῦτα καὶ ἄγρια, καὶ σχεδὸν τά γε πολλὰ παραπλησίαν έχοντα την όψιν τοις ημέροις, πλην τοίς γε φύλλοις έλάττω ταῦτα καὶ τραγύτερα καὶ τοῖς καυλοῖς καὶ μάλιστα τοῖς γυλοῖς δριμύτερα καὶ ἰσχυρότερα, καθάπερ ή τε θύμβρα καὶ ή ορίγανος ή τε ράφανος καὶ τὸ πήγανον ἐπεὶ καὶ τὸ λάπαθον ἄγριον, καίπερ εὐστομώτερον τοῦ ημέρου ὄν, τὸν χυλὸν ὅμως ὀξύτερον ἔχει καὶ τούτω μάλιστα διαφέρει. πάντα δὲ καὶ Επρότερα τῶν ἡμέρων, καὶ ἴσως αὐτῷ τούτῳ τά γε πολλά καὶ δριμύτερα καὶ ἰσχυρότερα.

Ίδίως δὲ ἡ ράφανος ἔχει παρὰ τὰ ἄλλα τοὺς καυλούς περιφερεστέρους και λειοτέρους της ημέ-

98

i.e. drying-up; cf. Plat. Tim. 85 A.
 σκωλήκωσιν conj. Sch.; κώλυσιν Ald.; σκωλήκησιν conj. R. Const.

³ σικυῶνι Ald.: perhaps here a general term for cucumbers, gourds, etc.; σικυῶν M; σικύφ conj. W.
⁴ Plin. 19. 185.

ENQUIRY INTO PLANTS, VII. v. 6-vi.

longer time, except that such seed must necessarily become less vigorous by reason of 'evaporation' and destruction by worms. The chief cause of loss is vermin; for vermin occur in all the seeds, even those which are pungent, though least in the gourd tribe; such seeds however, as they lose their moisture, become bitter in taste and inferior for use. Let this suffice for an account of the seeds and in general of herbs cultivated in gardens.

Of uncultivated herbs: the wild forms of pot-herbs.

VI. ⁴ We must now endeavour to speak in the same way of the wild kinds and of those which are called uncultivated herbs. Some of these have the same names as the cultivated ⁵ kinds; for all these kinds exist also in a wild form, and most of them resemble the cultivated kinds in appearance, except that in the wild forms the leaves and also the stalks are smaller and rougher, and in particular these forms are more pungent and stronger in taste, for instance, savory ⁶ marjoram cabbage and rue; the wild monk's rhubarb (dock) indeed, though it has a pleasanter taste than the cultivated, yet has ⁷ a sharper flavour; and this is the chief difference. Moreover all the wild kinds are less juicy than the cultivated, and perhaps this is the very reason why most ⁵ of them are more pungent and stronger.

A peculiarity of 'wild cabbage' as compared with the others is that its stems are rounder and smoother

⁵ ἡμέροις conj. Sch.; εἰρημένοις Ald. The correction would seem unnecessary but that Ald. gives εἰρημένοις in § 4 where ἡμέρους is required.
⁶ cf. Diosc. 3. 37.

⁷ όν, τὸν Ι conj.; τὸν δὲ MSS.W.

⁸ γε conj. Sch.; τε UMAld.

ρου, καλ τὴν τοῦ φύλλου πρόσθεσιν ἐκείνη μὲν ἔχει πλατεῖαν αὕτη δὲ περιφερεστέραν, καὶ αὐτὸ δὲ τὸ φύλλον ἀγωνότερον ἐπεὶ τά γε ἄλλα τραχύτερα καὶ τοῖς καυλοῖς καὶ τοῖς φύλλοις.

Η δε γογγυλίς και την ρίζαν έχει μακράν και

ραφανιδώδη καὶ τὸν καυλὸν βραχύν.

Θριδακίνη δὲ τό τε φύλλον βραχύτερον τῆς ήμέρου, καὶ τελεουμένης ἀκανθοῦται, καὶ τὸν καυλὸν ὁμοίως, τὸν ὀπὸν δὲ δριμὰν καὶ φαρμακώδη. φύεται δ' ἐν ταῖς ἀρούραις ἀπίζουσι δ' αὐτὴν ὑπὸ πυραμητόν, καί φασι καθαίρειν ὕδρωπα καὶ ἀχλὰν ἀπ' ὀφθαλμῶν ἀπάγειν καὶ ἄργεμα

άφαιρείν έν γάλακτι γυναικείφ.

Τὸ δ' ἰπποσέλινον καὶ ἐλειοσέλινον καὶ ὀρεοσέλινον καὶ πρὸς ἐαυτὰ διαφορὰν ἔχει καὶ πρὸς τὸ ἡμερον· τὸ μὲν γὰρ ἐλειοσέλινον τὸ παρὰ τοὺς ὀχετοὺς καὶ ἐν τοῖς ἔλεσι φυόμενον μανόφυλλόν τε καὶ οὐ δασὺ γίνεται, προσεμφερὲς δέ πως τῷ σελίνῳ καὶ τῷ ὀσήματι. τὸ δ' ἱπποσέλινον φύλλον μὲν ἐμφερὲς τῷ ἐλειοσελίνῳ, δασὺ δὲ καὶ μεγαλόκαυλον καὶ τὴν ρίζαν ὥσπερ ῥαφανὶς ἔχει τὸ πάχος μέλαιναν· μέλας δὲ καὶ ὁ καρπός, μέγεθος δὲ μεῖζον ὀρόβου. χρήσιμα δ' ἄμφω φασὶ πρὸς στραγγουρίαν εἰναι ἐν οἴνῳ γλυκεῖ λευκῷ καὶ τοῖς λιθιῶσι· φύεται δὲ

² Plin. 20. 20; Diosc. 2. 110.

¹ τραχύτερα conj. Sch.; τραχύτερον Ald., which contradicts what has just been said.

^{*} τελεουμένης conj. W.; τελεούμενος U; τελειούμενον $P_2Ald.$ cf. C.P. 4. 3. 5.

ENQUIRY INTO PLANTS, VII. vi. 2-3

than in the cultivated kind, and, while in the latter the attachment of the leaf is flat, in the wild kind it is rounder, and the leaf itself has less angles; in other cases the wild form is the rougher 1 both in stem and leaf.

²The wild turnip has a long root, like that of the

radish, and a short stem.

The wild lettuce has a shorter leaf than the cultivated kind, and, as the plant matures,³ it becomes spinous; the stem is also shorter, while the juice is pungent and medicinal. It grows in fields; they extract its juice at the time of wheat-harvest, and it is said that it purges away dropsy and takes away dimness of sight and removes ulcers⁴ on the eye; for which purpose it is administered in human milk.

5 'Horse-celery' (alexanders) 'marsh-celery' and 'mountain-celery' (parsley) differ both from one another and from the cultivated kind; 'marsh-celery,' which grows by irrigation-ditches and in marshes, has scanty leaves,6 and is not of close habit, 'yet it somewhat resembles the cultivated kind in smell taste and appearance. 'Horse-celery' has a leaf like that of the marsh kind, but is of close habit and has a big stalk, and its root is as thick as a radish and black; 8 the fruit is also black, and in size is larger than the seed of a vetch. They say that both kinds are serviceable in cases of strangury and for those suffering from stone, being administered in sweet white wine. Both kinds grow equally 9

⁹ δμοίως conj. Sch.; δμως Ald.

^{4 9. 9. 5;} Plin. 20. 58; Diosc. 2. 136.

⁵ Plin. 19. 124.

 $^{^{6}}$ μανόφυλλον: Plin. l.c. seems to have read μονόφυλλον.

⁷ Diosc. 3. 64. ⁸ Diosc. 3. 67.

όμοίως πανταχοῦ· γίνεται δὲ καί τι δάκρυον ἐξ αὐτοῦ ὅμοιον τἢ μύρρα· οἱ δέ φασιν ὅλως μύρραν.

Τὸ δὲ ὀρεοσέλινον μείζους ἔτι διαφορὰς ἔχει· τὸ μὲν γὰρ φύλλον ἔοικε κωνείω, ρίζα δὲ λεπτή, τὸν δὲ καρπὸν ἔχει καθάπερ ἄνηθον πλὴν ἐλάττω· διδόασι δὲ τοῦτον ἐν οἴνω αὐστηρῷ τῶν γυναικείων χάριν.

Ένια δὲ ὅλως ἀσύμβλητα τοῖς ἡμέροις ἐστὶ κατά γε τοὺς χυλοὺς καὶ τὰς δυνάμεις, ὥσπερ σίκυος ὅ τε ἄγριος καὶ ὁ ἥμερος, ἀλλ' ἐκ τῆς προσόψεως ἔχει τὴν ὁμοιότητα, καθάπερ καὶ ἐν τοῖς στεφανώμασιν ἡ ἰωνία· τὸ γὰρ φύλλον ἔχει παρόμοιον. τούτων μὲν οὖν ἐν τοῖς εἰρημένοις αἱ διαφοραί.

VII. Τῶν δὲ ἀρουραίων λεγομένων μετὰ ταῦτα ρητέον, καὶ ὅλως εἴ τι ποιῶδές ἐστιν ὁ μὴ τυγχάνει βρωτόν. καλοῦμεν γὰρ λάχανα τὰ πρὸς τὴν ἡμετέραν χρείαν ἐν δὲ τῷ καθ' ὅλου κἀκεῖνα περιέχεται, δι' ὁ καὶ περὶ ἐκείνων λεκτέον. λάχανα μὲν δὴ καὶ τὰ τοιαῦτα καλεῖται, κιχόρη ἀπάπη χόνδρυλλα ὑποχοιρὶς ἡριγέρων, καὶ ὅλως

¹ τι conj. Sch.; τδ Ald. cf. 9. 1. 4.

δλως P₂Bas; ὅλως Ald.; ² ἀπλῶς W.
 κωνείφ conj. Sch.; κονίφ Ald. cf 1. 5. 3 n.

⁴ καρπόν conj. Cornarius on Diosc. 3. 67. and Dalec.; καυλόν UMAld. cf. Diosc. l.c.

⁵ ἡμέροις Η.; είρημένοις UMAld. cf. 7. 6. 1 n.

⁶ See Index, σίκυος.

⁷ i.e. which gives them a common name.

 $^{^{8}}$ εἴ τι ποιωδές ἐστιν Η.; ἐγγειποιῶδες U; ἐγγειποιῶδές ἐστι MAld.

ENQUIRY INTO PLANTS, VII. vi. 3-vii. i

everywhere. There is also a sort 1 of gum which exudes from the plant, like myrrh, and some say that

it is 2 myrrh.

'Mountain-celery' (parsley) exhibits even greater differences; its leaf is like that of hemlock,³ the root is slender, and the fruit ⁴ like that of dill, but smaller; it is given in dry wine for diseases of women.

In some cases however the wild kinds are not in the least like the cultivated ⁵ in taste and properties; thus the wild and the cultivated cucumber ⁶ are quite different, and their resemblance ⁷ is due only to their general look, as, among coronary plants, there is resemblance between the wild and the cultivated kinds of gilliflower; for the leaves are alike. We have then described the differences which these plants present.

Of other uncultivated herbs, which may be classed with pot-herbs.

VII. Next we must speak of the differences found in the herbs called 'uncultivated,' and in general in any herbaceous plants 8 which are not edible. For we give the name of 'pot-herbs' to those which are cultivated for our own use, but in a wider sense the term includes these also; wherefore we must speak of them too. 9 Under the name 'pot-herbs' are included also 10 such plants as chicory dandelion 11 khondrylla 12 cat's ear groundsel, and in general all

⁹ Plin. 21. 89. ¹⁰ καl add. Scal.

12 χόνδρυλλα conj. Salm. from Plin. l.c., cf. 7. 11. 4 n.; αν-

δρύαλα Ald.G. cf. Plin. 21. 105; Diosc. 2. 133.

¹¹ ἀπάπη (or ἀπάτη) conj. Sch.; ἀφάκη Ald. The latter is a leguminous plant mentioned 8. 5. 3, etc.: for ἀπάπη cf. 6. 4. 8; 7. 8. 3; 7. 11. 3; for spelling see notes on the last two passages.

όσα κιχοριώδη καλείται διὰ τὴν ὁμοιότητα τῶν φύλλων πάντα γάρ πως ἐμφερῆ ἔχει τῷ κιχορίῳ πάλιν καυκαλὶς ἔνθρυσκον ἡδύοσμον. οἱ δὲ μυρία ἄλλα καλοῦσιν, σκάνδιξ καὶ ὅσα ἄλλα τοιαῦτα σκανδικώδη, τραγοπώγων, οἱ δὲ κόμην καλοῦσιν, ὁ τὴν μὲν ῥίζαν ἔχει μακρὰν καὶ γλυκείαν τὰ δὲ φύλλα τῷ κρόκῳ ὅμοια πλὴν μακρότερα, τὸν καυλὸν δὲ βραχύν, ἐφ' οὖ τὴν κάλυκα μεγάλην καὶ ἐξ ἄκρου μέγαν τὸν πάππον πολιόν, ἀφ' οὖ

καλείται τραγοπώγων.

Όμοίως δὲ καὶ ὅσα ἄλλα τοιαύτας μὲν ἰδέας ἔχει τοὺς δὲ χυλοὺς ἐδωδίμους ἡ ὡμοὺς ἡ ἑφθούς ἔνια γὰρ δεῖται πυρώσεως, ὥσπερ μαλάχη καὶ τευτλὶς καὶ τὸ λάπαθον καὶ ἡ ἀκαλύφη καὶ τὸ παρθένιον τὸν δὲ στρύχνον καὶ ἀμὸν ἐσθίουσιν, δν καὶ εὐκήπευτόν τινες πρότερον καὶ ἔτερα δὲ πλείω τούτων, ἐν οῖς καὶ ὁ παροιμιαζόμενός ἐστι διὰ πικρότητα κόρχορος ἔχων τὸ φύλλον ἀκιμῶδες. πάντα δὲ τὰ μὲν ἐπέτεια τὰ δὲ ἐπετειόκαυλα τυγχάνει τὰ μὲν γὰρ ἐξαυαίνονται τῶν δὲ διαμένουσιν εἰς πλείω χρόνον αὶ ρίζαι σχεδὸν δὲ οὐκ ἐλάττω τὰ τοιαῦτά ἐστι.

3 Φύεται δὲ τὰ μὲν καὶ ἀπὸ τῶν ριζῶν καὶ ἀπὸ τῶν σπερμάτων, τὰ δὲ ἔτερα μόνον ἀπὸ σπέρ-

1 καλείται conj. Sch.; ταῦτα Ald.

Plin. 27. 142; Diosc. 2. 138.
 κάλυξ: cf. 8. 2. 4; 8. 4. 3.

5 πάππον conj. W.; παγητὸν UMAld.; πώγωνα Η. cf. Diosc. l.c., where Saracenus corrects καρπὸς to πάππος.

6 τοιαύτας (sc. herbaceous) PmBas.; τοιαύτα τὰς Ald.; τὰς αὐτὰς conj. W.

² ξυθρυσκον: Šch. conjectured ξυθρυσκον, form corrected by L.Dindorf; ξυθουσικόν Ald.G. cf. Plin. 22. 81.

ENQUIRY INTO PLANTS, VII. vII. 1-3

those that are called 1 'chicory-like' because of the resemblance in the leaves; for to a certain extent the leaves of all these are like those of chicory; and we may add kaukalis chervil 2 green mint. Some include under the name countless others, as wild chervil and all plants that resemble it, and goat's beard,3 which some call kome ('hair'), which has a long sweet root and leaves like those of the crocus. but longer, and a short stem, on which is set the sheath 4; this is large, and on the top is the large mass of grey pappus,5 from which it gets its name of 'goat's beard.

In like manner all those may be included which have a similar 6 appearance, but juices suitable for food whether raw or cooked; for some need the action of fire, as malakhe (cheese-flower) beet monk's rhubarb nettle and bachelor's buttons; while garden nightshade 7 is also eaten raw, and some in former times 8 considered it worth growing in gardens. There are also many more, including the plant which has become proverbial of for its bitterness, blue pimpernel, which has a leaf like basil. All these are either annual or have annual stems; for some of them wither away altogether in one season, while of others the roots persist for a longer time, and to this

Some of these plants grow from roots and also from seed-unless in some cases they come up

8 πρότερον Ald.; *πρότερον Bas.; ωνύμασαν conj. W. Text

probably defective.

class belong the majority.

⁷ i.e. στρύχνος δ εδώδιμος: cf. 7. 15. 4. The American 'wonder-berry.'

⁹ κόρχορος εν λαχάνοις is the proverb. cf. Ar. Vesp. 239, Schol.; Plin. 21. 183. (= 'Is Saul also among the prophets?')

ματος, εἰ μή τι καὶ αὐτόματον. ἡ δὲ βλάστησις καὶ τούτων καὶ τῶν ἄλλων τῶν μὲν ἄμα τοῖς πρώτοις ὑετοῖς ἐστι μετ' ἰσημερίαν, οἶον ἀπάπης καὶ τοῦ κύνωπος καὶ ἡν καλοῦσί τινες βούπρηστιν, τῶν δὲ μετὰ Πλειάδα, καθάπερ καὶ κιχορίου καὶ σχεδὸν τῶν ἄλλων τῶν κιχοριωδῶν. καὶ τὰ μὲν εὐθὺς ἄμα τῆ βλαστήσει τὸ ἄνθος ἀφίησι, καθάπερ ἡ ἀφία, τὰ δὲ ὕστερον οὐ πολλῷ, καθάπερ ἡ ἀνεμώνη, τὰ δὲ ἄμα τῷ ἡρι καὶ ἐκκαυλεῖ καὶ ἀνθεῖ, καθάπερ τὸ κιχόριον καὶ τὰ κιχοριώδη καὶ τῶν ἀκανθικῶν ὅσα λαχανώδη.

Διαφορὰ δὲ τῶν ἀνθῶν πολλή, περὶ ἡς ἐν τοῖς πρότερον εἴρηται· σχεδὸν γάρ ἐστι κοινὸν ἀπάντων· ἔνια δὲ καὶ ὅλως ἀνανθῆ, καθάπερ καὶ τὸ ἐπίπετρον. συμβαίνει δὲ τοῖς ἄμα τῷ καυλῷ τὸ ἄνθος ἀφιεῖσι ταχεῖαν εἶναι τὴν ἀπάνθησιν· πλὴν ἡ μὲν ἀπάπη γηράσαντος τοῦ πρώτου πάλιν ἄλλο καὶ ἄλλο παραφύει, καὶ τοῦτο ποιεῖ παρ' ὅλον τὸν χειμῶνα καὶ τὸ ἔαρ ἄχρι τοῦ θέρους. πολὺν δὲ χρόνον καὶ ὁ ἡριγέρων. τὰ δὲ ἄλλα οὐ ποιεῖ τοῦτο, καθάπερ οὐδὲ ὁ κρόκος οὔτε ὁ εὔοσμος οὔθ' ὁ λευκὸς οὐθ' ὁ ἀκανθώδης· οὖτοι δὲ ἄοσμοι.

VIII. Κοινή δὲ διαφορὰ πάντων τῶν ποιωδῶν ή τοιάδε· τὰ μὲν γάρ ἐστιν ὀρθόκαυλα καὶ νευρό-

¹ δè after τούτων om. W.

² ἀπάπης (or ἀπάτης) conj. Sch.; ἀφάκης U; ἀφάκεις MAld. cf. 7. 7. 1 n. Plin. l.c., however, has aphace.

³ ἐπίπετρον conj. Scal. from Plin. l.c.; ἐπίμετρον UMAld.G. cf. Hesych. 4 καυλώ conj. Sch.: καρπώ UMAld.G.

ENQUIRY INTO PLANTS, VII. vii. 3-viii. 1

spontaneously. The growth alike of these ¹ and of others takes place in some cases with the first rains after the equinox, for instance, dandelion ² ribgrass and the plant which some call buprestis; in other cases after the rising of the Pleiad, for instance, chicory and most of the plants of that class. Some produce their flower immediately at the time of making growth, as lesser celandine, some not long after, as anemone, while some as soon as spring comes send up both their stems and flower, as chicory and the plants which resemble it, and those spinous plants which come under the head of pot-herbs.

There is much difference in the flowers, of which we have spoken already; for such difference is a thing common to all; and some are altogether flowerless, as stonecrop.³ Those which produce their flower with the stem ⁴ quickly shed the flower; except that dandelion,⁵ when the first flower is past its prime, produces another and yet another, and continues to do so right through the winter and spring up to the summer. Groundsel ⁶ also blooms for a long time; the others however do not do this; for instance the crocus does not, neither the scented (saffron crocus) nor the white nor the spinous kind,⁷—which last are scentless.

Of the differences in stem and leaf found in all herbaceous plants.

VIII. A distinction which is found in all herbaceous plants alike is the following:—some have straight

107

δ ἀπάπη γηρήσαντος conj. W.; ἀπηγηράσαντος U; ἀπογηράσαντος MAld.; ἀφάκη ἀπογηράσαντος H. cf. Plin. l.c.; 7. 7. 1 n.
 cf. C.P. 1. 22. 4; Plin. 25. 106.

⁷ See Index. This plant can only have been called κρόκοs because it produced a yellow dye.

καυλα, τὰ δὲ ἐπιγειόκαυλα, καθάπερ μαλάχη σκάνδιξ σίκυος ἄγριος τὸ δὲ ἡλιοτρόπιον ἔτι μάλλον ώς είπειν τοιούτον, ώσπερ και έν τοις . ἀκανθώδεσιν οὖσιν τρίβολος καὶ ἡ κάππαρις καὶ άλλα πλείω καὶ γὰρ ἐκείνων ἡ διαφορὰ πλείων. ένια δὲ περιαλλόκαυλα, μὴ ἔχοντα δὲ ποῦ προσπέσωσιν επιγειόκαυλα, καθάπερ επετίνη καὶ ἀπαρίνη καὶ άπλῶς ὧν ὁ καυλὸς λεπτὸς καὶ μαλακὸς καὶ μακρός, δι' δ καὶ φύονται ταῦτα ώς ἐπὶ τὸ παν έν άλλοις κοινή δή και αυτη ή διαφορά πάντων οὐ μόνον τῶν ποιωδῶν καὶ φρυγανικῶν άλλὰ καὶ τῶν θαμνωδῶν καὶ γὰρ ἡ ἔλιξ καὶ ἔτι μαλλον ή σμίλαξ περιαλλόκαυλον.

Έτι δὲ καὶ τῶν ποιωδῶν τὰ μὲν πολύκαυλα τὰ δέ μονόκαυλα· καὶ τῶν μονοκαύλων τὰ μὲν ἀπαράβλαστα κατά τὸν καυλὸν τὰ δὲ παραβλαστικά, καθάπερ καὶ ἐν τοῖς ἡμέροις ἥ τε ῥαφανὶς καὶ άλλ' άττα. πολύκαυλα δε ώς άπλως είπειν τὰ έπιγειόκαυλα, μονόκαυλα δὲ καὶ ὀλιγόκαυλα τὰ ορθόκαυλα. τούτων δὲ ἀπαράβλαστα τὰ λειόκαυλα κρόμυον πράσον σκόροδον, ώσπερ καὶ ἐν τοις ήμέροις και τὰ μὲν εὐθύκαυλα τὰ δὲ σκολιόκαυλα καὶ τούτων [τοῖς ἡμέροις] ὑπάρχει.

Διαφορά δέ τις καὶ τοιάδε τῶν ποιωδῶν ἐστι· τὰ μὲν γὰρ ἐπιγειόφυλλα τὰ δ' ἐπικαυλόφυλλα τυγγάνει τὰ δ' ἀμφοτέρως. ἐπιγειόφυλλα μὲν

¹ επιγειδκαυλα conj. Cornarius; επετειδκαυλα Ald H. ² cf. 7. 15. 1; Diosc. 4. 190 and 191; Plin. 22. 57.

ENQUIRY INTO PLANTS, VII. viii. 1-3

and fibrous stems, some prostrate stems, ¹ as malakhe (cheese-flower) wild chervil 'wild cucumber' (squirting cucumber); while heliotropion ² has this character ³ to an even greater extent, and so, among spinous plants, have caltrop caper and several others; for in these too the above-mentioned distinction is even more marked. Some again have clasping stems, but if they have nothing on which to throw themselves, their stems become prostrate, as epetine bedstraw and in general those which have a slender soft long stem; wherefore these in general grow in the midst of ⁴ other plants. This point of difference too is common not only to all herbaceous plants and under-shrubs, but also to shrubby ones; for helix (ivy) has a clasping stem, and, still more, smilax.

Again of herbaceous plants too some have several stems, some only one; and of the latter some have no side-shoots along the stem, while others have side-shoots, for instance, among cultivated plants radish and some others. Those with prostrate stems have generally more than one, while those with erect stems have but one or a few. Of these those with smooth stems have no side-shoots, as onion leek garlic—the wild, as well as the cultivated forms; and of these 5 again some have straight, some crooked

stems.

There is also the following point of difference in herbaceous plants:—some have their leaves on the ground, some on the stem, some have both characters. The following have ground leaves—crowsfoot ⁶ the

4 ἐν; G seems to have read ἐπ'.

6 cf. C.P. 2. 5. 4; Plin. 22. 48; Diosc. 2. 130.

³ τοιούτον conj. Sch. from G; τούτων Ald.

⁵ τοις ήμέροις probably repeated by mistake from above.

κορωνόπους ἄνθεμον ἀφύλλανθες ἄγχουσα πόα ανεμώνη απαργία αρνόγλωσσον απάπη έπικαυλόφυλλα δὲ κρηπὶς ἄνθεμον τὸ φυλλῶδες λωτὸς λευκόϊον άμφοτέρως δὲ τὸ κιγόριον καὶ γὰρ ἐπὶ τῶν καυλῶν ἄμα ταῖς ἐκφύσεσι ταῖς ἀκρεμονικαῖς έκφύει τι καὶ ἄνθος καὶ τῶν φυλλακάνθων ἔνια, πλην άκανθώδεσι κομιδή, καθάπερ ο σόγκος.

ΙΧ. "Εστι δὲ καὶ τὰ μὲν ἄκαρπα τὰ δὲ κάρπιμα. καὶ ὅλως τῶν ποιωδῶν τὰ μὲν ἄγρι τῶν φύλλων ἀφικνεῖται, τὰ δὲ καυλὸν ἔχει καὶ ἄνθος καρπον δε ού. τὰ δε καὶ καρπον ώσπερ τελειοτάτην Φύσιν, εἰ μή τι καὶ ἄνευ τοῦ ἄνθους καρποφόρον, ώσπερ έπλ τῶν δένδρων.

Διαφέρει δὲ καὶ τὰ φύλλα σχεδὸν οὐκ ἐλάττοσιν άλλα πλείοσι διαφοραίς ή τα των δένδρων. καὶ πρὸς αὐτὰ δὲ ἐκεῖνα διαφοράς ἔγει μεγίστην μεν ώς είπειν ότι τὰ μεν ἀπὸ μίσχου προσπέφυκε, τὰ δὲ αὐτὰ μὲν ὡς ἀπλώς, τὰ δὲ καυλική τινι προσφύσει. και των μεν έν τη βλαστήσει προτερεί < ὁ καυλός>, τῶν δὲ πλείστων τὰ φύλλα, καὶ σχεδὸν ἐν τῆ ἀρχῆ μέγιστα γίνονται καὶ μάλιστα έδώδιμα τὰ δὲ ἐκ τῶν δένδρων προωθεῖ τινα καυλόν.

017

¹ ἀφύλλανθες placed after ἄνθεμον by Sch.; in Ald. placed after ἀνεμώνη. cf. 7. 14. 2; Plin. 21. 56.
2 ἀπάπη U; ἀπάτη Ald. cf. 7. 7. 1 n.
3 cf. 1. 13. 1. and Index.

⁴ έκφύει τι και MSS .; ? έκφύει φύλλον τε και W.

ENQUIRY INTO PLANTS, VII. vIII. 3-1X. I

anthemon whose flowers have no petals 1 (wild camomile) alkanet grass anemone hawk's beard plantain dandelion 2; the following have leaves on the stem—ox-tongue the anthemon which has petalled flowers 3 trefoil gilliflower; while chicory has both kinds of leaves; for this plant produces, 4 as well as leaves, a certain number of flowers on the stems at the points where the side-shoots are attached. Similar too are some of the plants with spinous leaves, but not those that are altogether spinous, as sow-thistle.

Of other differences seen in herbaceous plants in general, as compared with one another and with trees.

IX. Again some are barren, while others bear fruit, and, speaking generally, of herbaceous plants some get as far as producing leaves only, others have a stem and flower, but no fruit; some again have fruit as the completion of their development, while some bear fruit even though they have no flower, as is the case with some trees.

⁶ The leaves of herbaceous plants again differ in hardly fewer, nay, even in more, ways than those of trees, and further, they present differences as compared with these, the chief being perhaps that some are attached by a leaf-stalk, some are attached directly, some attached with cauline appendages.⁶ And in some herbaceous plants the stalk ⁷ is the first part to grow, but in most the leaves, which almost at the outset grow to their largest and are best for eating; whereas the leaves of trees always push out first a sort of stalk.

⁵ Plin. 21. 100.

⁷ δ καυλός add. Sch. from G.

⁶ i.e. petiolate, sessile, and decurrent respectively.

2 Διαφέρουσι δὲ καὶ τοῖς ἄνθεσι πολύ· ἐν μὲν γὰρ τοῖς δένδρεσι τά γε πλεῖστα λευκά, τὰ δὲ μικρὸν ἐπιπορφυρίζοντα, τὰ δὲ ποώδη καὶ χλοώδη, κεχρωσμένον δὲ ἀνθινῷ < οὐδέν· ἐν δὲ τοῖς ποιώ-δεσι τῶν ἀνθῶν > πολλαὶ καὶ παντοδαπαὶ χροιαὶ καὶ ἄκρατοι καὶ μεμιγμέναι καὶ εὕοσμοι δὴ καὶ ἄσσμοί εἰσιν. καὶ τὰ μὲν δένδρα τὴν ἄνθησιν ἀθρόαν ποιεῖται, τούτων δ' ἔνια κατὰ μέρος, ὥσπερ ἔλέχθη καὶ περὶ τοῦ ἀκίμου, δι' δ καὶ πολὺν χρόνον ἀνθεῖ, καθάπερ ἄλλα τε πολλὰ καὶ τὸ ἡλιοτρόπιον καὶ τὸ κιχόριον.

Πολλαί δὲ καὶ τῶν ριζῶν διαφοραί καὶ τρόπον τινὰ αἱ τούτων φανερώτεραι· εἰσὶ γὰρ αἱ μὲν ξυλώδεις αἱ δὲ σαρκώδεις καὶ ἰνώδεις, ὥσπερ καὶ τῶν ἡμέρων, καθάπερ αἵ τε τοῦ σίτου καὶ τῆς πόας τῆς πλείστης. αὐτῶν δὲ τούτων ἔκασται πλείστας ἔχουσι διαφορὰς χρώμασιν ὀσμαῖς χυμοῖς μεγέθεσιν· αἱ μὲν γὰρ λευκαὶ αἱ δὲ μέλαιναι αἱ δ' ἐρυθραί, καθάπερ ἤ τε τῆς ἀγχούσης καὶ τοῦ ἐρευθεδάνου· αἱ δ' ὥσπερ ξανθαὶ καὶ ξυλοειδεῖς· καὶ γλυκεῖαι δὲ καὶ πικραὶ καὶ δριμεῖαι καὶ εὐώδεις καὶ κακώδεις, καὶ ἔνιαι φαρμακώδεις, ὡς

Διαφοραί δε και των σαρκωδων αι μεν γαρ στρογγύλαι αι δε προμήκεις και βαλανώδεις, ωσπερ ἀσφοδέλου και κρόκου και αι μεν λεπυριώδεις, ωσπερ ή τοῦ βολβοῦ και τῆς σκίλλης και ὅσαι βολβώδεις και κρομύου δε και γηθύου και

έν ἄλλοις εἴρηται.

¹ cf. 1. 13. 1.

² οὐδὲν... ἀνθῶν add. Scal. from G (κεχρωσμένων δὲ ἀνθικῷ πολλαὶ UMAld.); ἀνθινῷ for ἀνθικῷ conj. W., who also added τῶν ἀνθῶν. See LS. ἀνθινός.

ENQUIRY INTO PLANTS, VII. 1X. 2-4

There is also much difference as to the flowers between herbaceous plants and trees; for in trees 1 most of the flowers are white, while some are slightly reddish, others are greenish or greenish-vellow, but none of them 2 have distinct gay colours; while in herbaceous plants the flowers shew many and various colours, both simple and in combination, and further, some of them are scented, others not. Again 3 trees produce all their bloom at once, while some herbaceous plants have a succession of flowers, as we said 4 of basil: wherefore it is in flower for a long period, as are many other herbs, such as heliotropion and chicory.

There are also many differences in the roots, and, in a way, the differences in these are more obvious; some are woody, some fleshy and fibrous, just as in the cultivated kinds, as are those of corn and most kinds of grass. Again the roots themselves exhibit in each case very many differences in colour smell taste and size; some are white, some black, some red, as those of alkanet and madder; some are vellowish,5 or the colour of wood.6 Again there are roots which are sweet, bitter, pungent, fragrant, evil-smelling; and some are medicinal, as has been said elsewhere.

There are also differences between those with fleshy roots: the roots of some are round, of some oblong and acorn-shaped, as those of asphodel and crocus; 7 some consist of several layers, as those of purse-tassels squill and others which belong to that class, onion long onion and others like

113

VOL. II.

I

³ ἄοσμοι είσιν. καὶ τὰ μέν conj. W.; ἄοσμοι. καὶ ἐνίων τὰ μέν 4 7. 3. 1.

⁵ δσπερ MSS.; πλείσται conj. W.

⁷ cf. 1. 6. 7; 6. 6. 10. 6 ξυλοειδείs: cf. 7. 3. 2.

όσα τούτοις όμοια. αί δὲ όμαλεῖς καὶ ψαθυραὶ καὶ μαλακαὶ δι' ὅλου καὶ ισπερ ἄφλοιοι, καθάπερ τοῦ ἄρου αί δὲ φλοιὸν ἔχουσι πρὸς τῆ σαρκί, καθάπερ ή τοῦ κυκλαμίνου καὶ τῆς γογγυλίδος. ούν άπασαι δ' αί εὐώδεις ή γλυκείαι ή εὔστομοι καὶ ἐδώδιμοι, οὐδ' αἱ πικραὶ ἄβρωτοι ἀλλ' ὅσαι άβλαβεῖς εἰσι τῶ σώματι μετὰ τὴν προσφοράν. ένιαι γάρ γλυκείαι μεν θανάσιμοι δε καί νοσώδεις, 5 αί δὲ πικραί μὲν ἡ κακώδεις ωφέλιμοι δέ. τὸν αὐτὸν δὲ τρόπον καὶ φύλλα καὶ καυλοί, καθάπερ τοῦ ἀψινθίου καὶ τοῦ κενταυρίου. διαφορά δὲ καί κατά την βλάστησιν καί κατά την άνθησιν, οίον ἀργομένου γειμώνος και μεσούντος και πάλιν ήρος ή θέρους ή μετοπώρου. καὶ ἐπὶ τῶν καρπών δε όμοίως τῷ βρωτούς είναι καὶ έγχύλους ένίοις καὶ φύλλα καὶ σπέρματα καὶ ρίζας καὶ έν αὐτοῖς τούτοις κατά τοὺς χυλούς, οἶον ὀξύτητι καί δριμύτητι καί γλυκύτητι καί αὐστηρότητι καὶ ταῖς ἄλλαις ταῖς τοιαύταις άπλῶς τε καὶ κατὰ τὸ μᾶλλον. τας μέν ουν διαφοράς έν τούτοις ληπτέον.

Χ. Διηρημένων δὲ κατὰ τὰς ὥρας ἑκάστων πρός τε τὰς βλαστήσεις καὶ κατὰ τὰς ἀνθήσεις καὶ τελειώσεις τῶν καρπῶν, οὐδὲν ἀναβλαστάνει πρὸ τῆς οἰκείας ὥρας οὕτε τῶν ῥιζοφυῶν οὕτε τῶν σπερμοφυῶν, ἀλλ' ἔκαστον ἀναμένει τὴν

1 καl conj. W.; A Ald.

² τῷ βρωτοὺς εἶναι καὶ ἐγχύλους ἐνίοις conj. W.; τὸ βρωτὰ εἶναι καὶ καυλοὸς εἶναι Ald. H. Text probably defective.

ENQUIRY INTO PLANTS, VII. 1x. 4-x. 1

these. Some are smooth loose and soft throughout, and, as it were, without 'bark,' as those of cuckoo-pint, while some have a 'bark' attached to the fleshy part, as those of cyclamen and turnip. And not all those that are fragrant or sweet or pleasant to the taste are also 1 edible, any more than all those that are bitter are uneatable; any (whether sweet or bitter) that are harmless to the body after being eaten are edible; for some that are sweet are deadly and dangerous to health, while some are beneficial even if they are bitter or have an evil smell. The same may be said of the leaves and stalks. as in the case of wormwood and centaury. There are also differences in the time of growth and of flowering, the season being variously the beginning or middle of winter, or again spring, summer, or autumn. So too is there in like manner a difference in the fruits, which 2 in some of these plants are edible and juicy, as well as 3 the leaves seeds and roots. And in these cases there are further differences in the taste (of those which are edible and juicy), which may be sharp, pungent, sweet, dry, or exhibit other similar differences, either altogether or in degree. These are examples of the differences which we find.

Of the seasons at which herbs grow and flower.

X. Each plant having its proper season for growth, flowering and maturing of the fruit, nothing grows ⁴ before its proper season either of those grown from a root or of those grown from seed, but each awaits

³ καὶ βίζας seems irrelevant.

⁴ ἀναβλαστάνει conj. W.; ἀναβλαστεῖ καὶ Ald. H.

οικείαν οὐδ' ὑπὸ τῶν ὑδάτων οὐδὲν πάσχον ἔνια γαρ θερινά κομιδή καὶ τή βλαστήσει καὶ τή ανθήσει, καθάπερ ο τε σκόλυμος καὶ ο σίκυος άνριος, ώσπερ καλ περλ των φρυγανικών έλέχθη περί κονύζης τε καί καππάριδος καί των άλλων. οὐδεν γὰρ οὐδε εκείνων ἀνθεί καὶ βλαστάνει πρὸ 2 τῆς οἰκείας ὥρας. δι' δ κᾶν ταύτη δόξαιεν ᾶν διαφέρειν των δένδρων. των μεν γάρ αμα πως πάντων ή έγγυς ή βλάστησις, εί δὲ μη κατά μίαν γε ώραν ώς είπειν τούτων δε έν πολλαίς μάλλον δε εν απάσαις ή βλάστησις και έτι μαλλον ή ανθησις, ώστε εί τις εθέλει κατανοείν σχεδον συνεγής γίνεται καθ' όλον ένιαυτον και ή βλάστησις καὶ ή ἄνθησις αἰεὶ γὰρ ἔτερον έξ ἐτέρου διαδεγόμενον πάσας καταλαμβάνει τὰς ώρας. οίον μετά την απάπην κρόκος έσται και ανεμώνη καὶ ὁ ἢριγέρων καὶ τὰ ἄλλα χειμερινά, μετὰ δὲ ταθτα τὰ ἦρινὰ <καὶ θερινὰ> καὶ μετοπωρινά. 3 πολλά δέ, ώσπερ έλέγθη, διά τὸ κατά μέρος ἀνθεῖν έπιτείνεται ταις ώραις ένια γάρ ούτως άνθει, καθάπερ ή τε ἀπάπη και τὸ ὀνοχειλές και τὸ κιχόριον καὶ τὸ ἀρνόγλωσσον καὶ ἄλλα· διὰ δὲ την συνέχειαν και την περικατάληψιν την ύπ' άλλήλων οὐ φαίνεται ράδιον ἐν ἐνίοις οὐδ' δρίσαι ποία πρώτα βλαστάνει καὶ ποία όψιβλαστή. πλην εί τις υποθοίτο του έτους την άρχην τινα

³ ἀπάπην conj. W.; ἀφάκην UMAld. cf. 7. 7. 1 n.

⁵ και θερινά seems to have dropped out.

116

¹ οὐδ' conj. W.; οὕθ' Ald. ² Reference not discoverable.

⁴ κρόκος conj. Sch. (adding δ); ήρος U (corrected); κρος M; om. Ald.: το κρόκον mBas.

⁶ διὰ τό κατά μέρος άνθεῖν conj. W.; καὶ τῶν κατὰ μέρος άνθέων Ald.

ENQUIRY INTO PLANTS, VII. x. 1-3

its proper season and is not 1 affected even by rain. For some are plants which belong properly to summer as to their growth and their flowering, as golden thistle and squirting cucumber, as was said of shrubby plants and of konyza caper and the rest; for of these too none blooms or grows before its proper season. Wherefore in this respect too these plants would seem to differ from trees. For trees make their growth all at once or nearly so, or at all events we may say that they do so all at one season; but the plants of which we are now speaking have their times of growing and still more of flowering at many or rather at all seasons; so that, if one will consider it, both the growing and the flowering are almost continuous throughout the year; for one continually succeeds to another, so that all seasons are covered; thus after the dandelion 8 will come the crocus 4 anemone groundsel and the other plants of winter, and after these those of spring summer 5 and autumn. Some again, as was said, because they do not produce all their bloom at once,6 cover a longer season; for there are some that thus bloom, for instance dandelion bugloss chicory plantain, and others; but because of this continuity and overlapping it does not seem easy in some cases to define which first make growth and which are late in growing,unless one were to lay down that the 'year'

⁷ ἀπάπη conj. W.; ἀφάκη Ald. cf. 7. 7. 1 n.

8 dvoxeiles conj. Sch. from Plin. 21. 100; Diosc. 4. 24; dvo-

κίχλης UMAld.

[§] i.e. unless one has a fixed starting point. τινα ῶρας τινος ἀρχῆ conj. W.; τινα πρὸς τῆ Γνα ζῆ(?) U; text defective in MAld., but both give Γνα ζῆ; W. conjectures also τροπὰς τὰς χειμερινάς. ? εἶναι ὥρας τινος ἀρχήν or εἶναι ὥραν τινα (omitting εἶ ζῆ as a trace of a lost sentence).

- 4 ὥρας τινὸς ἀρχῷ. καὶ αὐτῶν δὲ τούτων τὰς γενέσεις καὶ τὰς ὥρας ὅταν τελειωθέντων τῶν καρπῶν πάλιν ἄλλας ἀρχὰς ἐνίστωνται τῆς γενέσεως ὅπερ μάλιστα δοκεῖ συμβαίνειν μετ ἀσημερίαν μετοπωρινήν τότε γὰρ ἤδη τὰ σπέρματα πλεῖστα τετελείωται καὶ τῶν δενδρικῶν καρπῶν οἱ πολλοί, καὶ ἄμα μεταβολή τις αὐτοῦ τούτου προσγίνεται καὶ τῆς ὥρας ὅσα δὲ ἀτελῆ καὶ ἄπεπτα περικαταλαμβάνεται, τούτοις κατὰ λόγον ἐκ περιόδου καὶ ἡ βλάστησις γίνεται καὶ ἡ ἄνθησις καὶ ἡ τελείωσις δι' ὁ συμβαίνει τὰ μὲν ὑπὸ τροπὰς ἀνθεῖν τὰ δ' ὑπὸ Κύνα τὰ δὲ καὶ μετὰ ᾿Αρκτοῦρον καὶ ἰσημερίαν μετοπωρινήν.
 - 'Αλλά ταῦτα μὲν ἔοικε κοινοτέραν ἔχειν σκέψιν εἰς ἀφορισμὸν ἀρχῆς. ὅτι δὲ αί διαφοραὶ πλείους ἡ οὐκ ἐλάττους ἐν τούτοις φανερόν. ἐπεὶ καὶ ἀείφυλλα τῶν τοιούτων ἐστὶν ἔνια, καθάπερ τὸ πόλιον καὶ ἡλιοτρόπιον καὶ τὸ ἀδίαντον.

ΧΙ. 'Αφωρισμένων οὖν τούτων περὶ τὰς διαφορὰς ἐν οἶς γίνονται καὶ πῶς λεκτέον ἤδη τὰς καθ' ἔκαστον ἱστορίας . . . ὅσα μὴ κατὰ τὴν

⁴ Plin. 21. 100.

¹ i.e. to fix the date of the beginning and end of the cycle of the plant's life.

 ² αὐτοῦ τούτου: ?the plant itself. αὐτοῦ τοῦ ἔτους conj. Sch.
 ³ i.e. according as the seed ripened last year or this year.
 In this rather obscure section I follow W.'s explanations.

ENQUIRY INTO PLANTS, VII. x. 3-xi. 1

begins when a certain season begins. Further in these plants it is not easy to define 1 in each case the time of first growth and the season when, the fruits being matured, it makes a fresh start in reproduction. This seems chiefly to occur after the autumnal equinox; for by that time most of the seeds are matured, like most of the fruits of trees: moreover a change then takes place in the seed 2 itself as well as in the season. But in the case of any seeds which are still immature and unripe and so are overtaken by winter, the period of first growth, the flowering of the new plant, and the period of maturity are proportionally later. Wherefore it comes to pass that some bloom at the solstice, some at the rising of the dog-star, and some after the rising of Arcturus and the autumnal equinox.8

But these matters seem to require a wider investigation in order to determine when the process begins. However it is clear from what has been said that these plants present at least as many differences as trees. ⁴ For some again of this class are evergreen, as hulwort heliotropion and maidenhair

Of the classes into which herbaceous plants may be divided, as those having a spike and chicory-like plants.

XI. Having then made these explanations we must now give a separate account of each plant, discussing the differences (in those plants in which they occur) and saying how they arise 5 except those peculiarities which belong to the



⁵ Lacuna in UMAld.Cam.Bas., leaving the connexion of the next clause obscure.

ίδίαν έκάστου φύσιν. λέγω δὲ οἶον τὰ σταχυώδη καὶ τὰ σκανδικώδη καὶ μονοφυή, κὰν εἴ τι ἔτερόν ἐστι τοιοῦτον κοινὸν ἐπὶ πάντων λαβεῖν, ὁ τῷ αἰσθήσει γνώριμον ἡ φύλλοις ἡ ἄνθεσιν ἡ ῥίζαις ἡ καρποῖς, ἐκ γὰρ τῶν φανερῶν ὁ μερισμὸς

ωσπερ καὶ ἐκ τῶν ῥιζῶν.

Σταχυώδη μὲν οὖν ἐστιν ὅ τε κύνωψ ὑπό τινων καλούμενος πλείους ἔχων ἰδέας ἐν ἑαυτῷ· καὶ ὁ ἀλωπέκουρος καὶ ὁ στελέφουρος ὑπ' ἐνίων δὲ ἀρνόγλωσσον τῶν δὲ ὅρτυξ καλούμενος· παρόμοιον δὲ τούτῳ τρόπον τινὰ καὶ ἡ θρυαλλίς. ἀπλᾶ δὲ καὶ μονοειδῆ τρόπον τινὰ ταῦτα καὶ στάχυν οὐκ όξὺν οὐδ' ἀθερώδη ἔχοντα· ὁ δ ἀλωπέκουρος μαλακὸν καὶ χνοωδέστερον, ὅτι καὶ ὅμοιον ταῖς τῶν ἀλωπέκων οὐραῖς, ὅθεν καὶ τοὕνομα μετείληφεν. ὅμοιος δὲ τούτῳ καὶ ὁ στελέφουρος, πλὴν οὐχ ὥσπερ ἐκεῖνος ἀνθεῖ κατὰ μέρος ἀλλὰ δι' ὅλου τοῦ στάχυος ὥσπερ ὁ πυρός. ἡ δὲ ἄνθησις ἀμφοῦν χνοώδης, καθάπερ καὶ τοῦ σίτου· παρόμοιον δὲ τῆ ὅλη μορφῆ τῷ πυρῷ πλὴν πλατυφυλλότερον. ὡσαύτως δὲ τούτοις καὶ τὰ ἄλλα τὰ σταχυώδη λεκτέον.

Τὰ δὲ κιχοριώδη πάντα μὲν ἐπετειόφυλλα καὶ ῥιζόφυλλα, βλαστάνει δὲ μετὰ Πλειάδα πλὴν τῆς ἀπάπης, τοῖς δὲ καυλοῖς καὶ ταῖς ῥίζαις

1 i.e. spicate.

² σκανδικώδη: i.e. umbellate. One would expect κιχοριώδη, to correspond with § 3; but the three classes mentioned seem to be merely 'samples' of classification: of the three only one (τὰ σταχυώδη) is described below, and other classes are added.

 $^{^3}$ μονοφυή I conj.: i.e. those which have a scape: cf. 2. 6. 9; C.P. 1. 1. 3. μυδφαα U; μυσφῶα MAld.; om, G.

ENQUIRY INTO PLANTS, VII. xi. 1-3

character of individual kinds. I mean for instance the plants which have a spike,1 those which may be classed with wild chervil,2 and those which have a single stem,3 . . . or any other such class in which one can find some such general characteristics obvious to the senses either in leaves flowers roots or fruits; (for the classification is to be made

by the visible parts, as well as by the roots).4

⁵ An example of the plants which have a spike is the plant which some call 'dog's eye 6' (rib-grass), which comprises several forms; we have also 'foxbrush,' stelephuros (plantain), which some call 'lamb's tongue' and some 'quail-plant'; and somewhat similar to this is thryallis. These are simple plants and uniform in character, having a spike which is not pointed nor bearded; while in 'fox-brush' it is soft and somewhat downy, in that it actually resembles the brush of a fox, whence also it has obtained its name. Similar to this is stelephuros (plantain), except that it does not, like that plant, flower here and there, but all up the spike like wheat. The bloom of both is downy like that of corn, and the plants in their general appearance resemble wheat, but have broader leaves. Of the other plants which have a spike a similar account may be given.

⁷ The chicory-like plants all have annual leaves and have root-leaves, and they begin to grow after the Pleiad, except dandelion 8: but in their stems

⁵ Plin. 21. 101.

⁴ Roots being the basis of classification in xii. below.

⁶ κύνωψ conj. Sch.; ἀχύνωψ UAld.; Plin. l.c. has cynops (cf. 7. 7. 3); oculus caninus G. 7 i.e. composites. Plin. l.c.

⁸ ἀπάπης Û; ἀπάτης MAld. cf. 7. 7. 1 n.; 7. 8. 3 n.

μεγάλας ἔχουσι διαφοράς· οἱ μὲν γὰρ τῶν ἄλλων ἀπλούστεροι καὶ ἐλάττους, ὁ δὲ τοῦ κιχορίου μέγας καὶ ἀποφύσεις ἔχων πολλάς, ἔτι δὲ γλίσχρος καὶ δυσδιαίρετος, δι' ὁ καὶ δεσμῷ χρῶνται· παραβλαστητικὸν δὲ καὶ τῆ ῥίζη καὶ ἄλλως μακρόρριζον, δι' ὁ καὶ δυσώλεθρον· ὅταν γὰρ ἐκλαχανίσωνται, πάλιν τὸ ὑπόλοιπον ἀρχὴν λαμβάνει γενέσεως. συμβαίνει δὲ καὶ παρανθεῖν αὐτοῦ μέρος ἄλλο καὶ ἄλλο, καὶ τοῦτο ἄχρι τοῦ μετοπώρου, σκληροῦ δοκοῦντος εἶναι τοῦ καυλοῦ. φέρει δὲ καὶ λοβὸν ἐν ῷ τὸ σπέρμα περὶ τὰ ἄκρα τῶν καυλῶν.

'Η δὲ ὑποχοιρὶς λειοτέρα καὶ ἡμερωτέρα τῷ προσόψει καὶ γλυκυτέρα καὶ οὐχ ὥσπερ ἡ χόν-δρυλλα· τὸ γὰρ ὅλον οὐκ ἐδώδιμος αὕτη καὶ ἄβρωτος καὶ ἐν τῷ ῥίζῃ δριμὸν ὀπὸν ἔχει καὶ πολύν.

"Αβρωτος δὲ καὶ πικρὰ ἡ ἀπάπη· πρωϊανθὴς δὲ καὶ ταχὺ γηράσκει καὶ ἀποπαπποῦται, εἶτ' ἄλλο φύεται πάλιν καὶ ἄλλο καὶ τοῦτο παρ' ὅλον ποιεῖ τὸν χειμῶνα καὶ τὸ ἔαρ ἄχρι τοῦ θέρους· τὸ δ' ἄνθος μηλινοειδές.

'Ωσαύτως δὲ καὶ ἡ πικρίς καὶ γὰρ αὕτη τῷ ἡρι ἀνθεῖ, καὶ παραπλησία δι' ὅλου τοῦ χειμῶνος καὶ τοῦ θέρους παρανθεῖ τῆ γεύσει δὲ πικρά, δι' ὁ καὶ τοὕνομα εἴληφε. ταῦτα μὲν οὖν ἐν ταύ-

² Plin. 21. 88. ³ τὸ conj. Sch.; τὴν Ald.

¹ ελάττους conj. Scal.; θάττους UMAld.

⁴ Meaning not obvious; σκληροῦ is perhaps corrupt.

ὁποχοιρίς conj. Scal.; ὑποχώρησις UMAld. cf. 7. 7. 1 n.
 οὐχ ἄσπερ: an adjective has perhaps dropped out between these words; ? πικρά (amara Plin. l.c.).

ENQUIRY INTO PLANTS, VII. xi. 3-4

and roots they exhibit great differences; for in some these are simpler and fewer,¹ but the stem of chicory is large and has many side-shoots; also it is tough and hard to break, wherefore it is used for withes²; it makes side-growths from the root, and also has long roots, wherefore it is hard to kill; for, when the top is taken off to use as a vegetable, what remains³ starts growing again. Moreover different parts of it flower at different times, and the flowering goes on till autumn, since the stem appears to be hard.⁴ Also it bears a pod, which contains the seed, at the top of the stem.

Cat's ear⁵ is smoother and has a more cultivated appearance, and is also sweeter and not like ⁶ khondrylla⁷; for the latter is altogether uneatable and ⁸ unfit for food, and its root contains a quantity of

pungent juice.

Dandelion 9 is also unfit for food and bitter: it flowers early and quickly waxes old and the flower turns to pappus; but then another flower forms, and yet another, and this goes on right through the winter and spring up to the summer; and the flower is vellow.

10 The like may be said of *pikris*: for this plant too blooms in spring, and like dandelion it flowers throughout the winter, and it flowers also to some extent in summer; in taste it is bitter, whence its name. These are the special points of difference

10 Plin. 22. 66.

⁷ χόνδρυλλα conj. St.; χανδρυ αλλα U; χανδρὸ αλλα M; χανδρὰς άλλὰ Ald. H.; cadryalia G (Tarv.); candralia G (Bas. Par.).

 ⁸ τὸ γὰρ ὅλον οὐκ ἐδώδιμος αὅτη καὶ conj. W.; τὸ ὅλον οὐκ ἐδ.
 αὅτη γὰρ Ald.; αὅτη γὰρ ἄβρωτος conj. Sch. from Plin. 21. 105.
 ⁹ ἀπάπη conj. W.; ἀφάκη Ald. of. 7. 7. 1 n.

ταις ταῖς διαφοραῖς. πειρατέον δέ, ὡς ἐλέχθη, καὶ τῶν ἄλλων λαμβάνειν ὁμοίως.

ΧΙΙ. Πολὺ δέ τι γένος ἐστὶ καὶ τῶν σαρκορρίζων ἡ κεφαλορρίζων, α καὶ πρὸς τὰ ἄλλα καὶ καθ' αὐτὰ τὰς διαφορὰς ἔχει ρίζαις τε καὶ φύλλοις καὶ καυλοῖς καὶ ταῖς ἄλλαις μορφαῖς. τῶν γὰρ ρίζῶν, ὥσπερ εἴρηται πρότερον, αἱ μὲν λεπυριώδεις αἱ δὲ σαρκώδεις, καὶ αἱ μὲν ἔχουσαι φλοιὸν αἱ δ᾽ ἄφλοιοι, ἔτι δὲ αἱ μὲν στρογγύλαι αἱ δὲ προμήκεις καὶ αἱ μὲν ἐδώδιμοι αἱ δ᾽ ἄβρωτοι. ἐδώδιμοι μὲν γὰρ οὐ μόνον βολβοὶ καὶ τὰ ὅμοια τούτοις, ἀλλὰ καὶ ἡ τοῦ ἀσφοδέλου ρίζα καὶ ἡ τῆς σκίλλης, πλὴν οὐ πάσης ἀλλὰ τῆς Ἐπιμενιδείου καλουμένης, ἡ ἀπὸ τῆς χρήσεως ἔχει τὴν προσηγορίαν αὕτη δὲ στενοφυλλοτέρα τε καὶ λειοτέρα τῶν λοιπῶν ἐστιν.

'Εδώδιμος δὲ καὶ ἡ τοῦ ἄρου καὶ αὐτὴ καὶ τὰ φύλλα προαφεψηθέντα ἐν ὅξει καί ἐστιν ἡδεῖά τε καὶ πρὸς τὰ ῥήγματα ἀγαθή. πρὸς δὲ τὴν αὔξησιν αὐτῆς, ὅταν ἀποφυλλίσωσιν, ἔχει δὲ μέγα σφόδρα τὸ φύλλον, ἀνορύξαντες στρέφουσιν, ὅπως ἂν μὴ διαβλαστάνῃ ἀλλὰ πᾶσαν ἕλκῃ τὴν τροφὴν εἰς ἑαυτήν, δ καὶ ἐπὶ τῶν βολβῶν τινες

¹ of. 7. 11. 2 ad fin.

² Plin. 19. 93 and 94.

^{*} $\mu\rho\rho\rho\alpha\hat{a}$ s: cf. 8. 4. 2. 4 7. 9. 4. 5 The legends about Epimenides suggest that the 'use' was possibly in magic: cf. what is said of $\sigma\kappa\ell\lambda\lambda\alpha$ 7. 13. 4. cf. Plin. l.c.

ENQUIRY INTO PLANTS, VII. xi. 4-xii. 2

about these plants; now we must endeavour, as was said, to set forth the special points of the other classes in like manner.

Of herbs which have fleshy or bulbous roots.

XII. ² There is a large class of these which have fleshy or bulbous roots: these exhibit differences both as compared with other plants and with one another both in roots leaves stems and their other prominent features.³ Of the roots, as has been said ⁴ already, some are in layers, some fleshy, some have a 'bark,' some not; and again some are round, some oblong, some edible and some not fit for food. Among edible roots are not only purse-tassels and others which resemble them, but also the roots of asphodel and squill, though not of all kinds of the latter, but only of the kind called 'Epimenides' squill (French sparrow-grass) which gets its name from its use ⁵; this kind has narrower leaves and is smoother than the others.

⁶ The root of cuckoo-pint is also edible, and so are the leaves, if they are first boiled down in vinegar; they are sweet, and are good for fractures. To increase the root, having first stripped ⁷ off the leaves (and the leaf is very large), they dig ⁸ it up and invert ⁹ it in order that it may not shoot, ¹⁰ but may draw all the nourishment into itself. This some

10 διαβλαστάνη: cf. C.P. 4, 8. 1.

125

⁶ Plin. 19. 96; 24. 162.

⁷ ἀποφυλλίσωσιν conj. Sch.; ἀποφυλάσσωσιν U; ἀποφυλλάσσωσιν MAld.

⁸ ἀνορύξαντες conj. St.; δ ἀνορύξαντες Ald.

⁹ cf. 1. 6. 10; Plin. 19. 94 and 97, who seems to have read κατορύξαντες: so also G. ? 'they plunge it in a pit.'

ποιοῦσι συντιθέντες· ἡ δὲ τοῦ δρακοντίου, καλοῦσι γάρ τι δρακόντιον ἄρον διὰ τὸ τὸν καυλὸν ἔχειν

τινα ποικιλίαν, άβρωτος και φαρμακώδης.

'Αλλὰ ή τοῦ φασγάνου καλουμένου γλυκεῖά τε έψηθεῖσα, καὶ τριφθεῖσα μιγνυμένη τῷ ἀλεύρω ποιεῖ τὸν ἄρτον γλυκὺν καὶ ἀσινῆ· στρογγύλη δέ ἐστι καὶ ἄφλοιος καὶ ἀποφύσεις ἔχουσα μικράς, ὥσπερ τὸ γήθυον· πολλὰς δὲ εὐρίσκουσιν ἐν ταῖς σκαλοπιαῖς· χαίρει γὰρ καὶ συλλέγει τὸ ζῶον.

'Η δὲ τοῦ θησείου τῆ μὲν γεύσει πικρά, τριβομένη δὲ κοιλίαν ὑποκαθαίρει. φαρμακώδεις δέ τινές εἰσι καὶ ἔτεραι, πολλών δὲ οὔτε φαρμακώδεις οὔτε ἐδώδιμοι. καὶ αὖται μὲν ἐν ταῖς ῥίζαις

αί διαφοραί.

ΧΙΙΙ. Κατὰ δὲ τὰ φύλλα τοῖς τε μεγέθεσιν καὶ τοῖς σχήμασιν. ὁ μὲν ἀσφόδελος μακρὸν καὶ στενότερον καὶ ὑπόγλισχρον ἔχει τὸ φύλλον, ἡ δὲ σκίλλα πλατὰ καὶ εὐδιαίρετον, τὸ δὲ φάσγανον ὑπό τινων δὲ καλούμενον ξίφος ξιφοειδές, ὅθεν ἔσχε καὶ τοὔνομα, ἡ δὲ ἶρις καλαμωδέστερον· τὸ δὲ τοῦ ἄρου πρὸς τῆ πλατύτητι καὶ ἔγκοιλον καὶ σικυῶδές ἐστιν· ὁ δὲ νάρκισσος στενὸν καὶ πολὰ καὶ λιπαρόν· βολβὸς δὲ καὶ τὰ βολβώδη παντελῶς στενὰ καὶ τοῦ κρόκου δ᾽ ἔτι στενότερον.

Καυλὸν δὲ τὰ μὲν οὐκ ἔχει τὸ ὅλον οὐδ' ἄνθος, ὅσπερ τὸ ἄρον τὸ ἐδώδιμον τὰ δὲ τὸν τοῦ ἄνθους μόνου, ισπερ ὁ νάρκισσος καὶ ὁ κρόκος ἔνια δὲ ἔχει, καθάπερ ἡ σκίλλα καὶ ὁ βολβὸς καὶ ἡ Ἰρις καὶ τὸ Είφιον μέγιστον δὲ πάντων ἀσφό-

² cf. 9. 20. 3; Plin. 24. 142; Diosc. 2. 166.

126

 $^{^1}$ συντιθέντες: sense doubtful. Sch. and W. mark the word as corrupt.

ENQUIRY INTO PLANTS, VII. xII. 2-XIII. 2

do also with purse-tassels, when they lay them by.¹
² However the root of edder-wort (for a kind of cuckoo-pint is so called because of its variegated stem)

is not good for food, but is used for drugs.

But the root of the plant called corn-flag is sweet, and, if cooked and pounded up and mixed with the flour, makes the bread sweet and wholesome. It is round and without 'bark,' and has small offsets like the long onion. Many of them are found in moles' runs ⁸; for this animal likes them and collects them.

⁴ The root of theseion is bitter to the taste, but when pounded purges the bowels. There are also certain others of these roots which have medicinal properties, but of many the roots are neither medicinal nor edible. Such are the differences in

the roots.

XIII. ⁵ In the leaves the differences are in size and shape. Asphodel has a long leaf, which is somewhat narrow and tough, while that of squill is broad and tears easily; corn-flag, which is called by some xiphos ('sword'), has a sword-like leaf, whence its name, and iris one more like a reed. That of cuckoopint, in addition to being broad, is concave and like that of cucumber; that of the narcissus is narrow substantial and glossy, those of purse-tassels and plants of that character are quite narrow, and that of crocus narrower still.

⁶ Some have not a stem at all, nor a flower, as the edible cuckoo-pint; some have only the flower-stem, as narcissus and crocus; some however have a stem, as squill purse-tassels iris and corn-flag; but asphodel

⁶ Plin. 21. 108 and 109.

παρὰ ταῖς σκαλοπιαῖς conj. Sch.; ἐν ταῖς σκ. conj. W.; ταῖς σκολοπίαις UMAld.
 Plin. 22. 66.
 Plin. 21. 108.

δελος ό γὰρ ἀνθέρικος μέγιστος ὁ δὲ τῆς ἴριδος ἐλάττων μὲν σκληρότερος δὲ τὸ δὲ ὅλον ἀνθερικώδης. ἔστι δὲ καὶ πολύκαρπος ὁ ἀσφόδελος, καὶ ὁ καρπὸς αὐτοῦ ξυλώδης τῆ μὲν μορφῆ τρίγωνος τῷ δὲ χρώματι μέλας γίνεται δὲ ἐν τῷ στρογγύλῳ τῷ ὑποκάτω τοῦ ἄνθους, ἐκπίπτει δὲ τοῦ βέρους, ὅταν τοῦτο διαχάνη. τὴν ἄνθησιν ποιεῖται κατὰ μέρος, ὥσπερ καὶ ἐπὶ τῆς σκίλλης, ἄρχεται δὲ πρῶτον ἀπὸ τῶν κάτωθεν. ἐν δὲ τῷ ἀνθερίκῳ συνίσταται σκώληξ, δς εἰς ἄλλο μεταβάλλει ζῶον ἀνθρηνοειδές, εἶθ' ὅταν ὁ ἀνθέρικος αὐανθῆ διεσθίον ἐκπέταται. δοκεῖ δὲ ἴδιον ἔχειν πρὸς τὰ ἄλλα τὰ λειόκαυλα, διότι στενὸς ὧν ἀποφύσεις ἄνωθεν ἔχει. πολλὰ δὲ εἰς τροφὴν παρέχεται χρήσιμα· καὶ γὰρ ὁ ἀνθέρικος ἐδώδιμος σταθευόμενος καὶ τὸ σπέρμα φρυγόμενον καὶ πάντων δὲ μάλιστα ἡ ῥίζα κοπτομένη μετὰ σύκου καὶ πλείστην ὄνησιν ἔχει καθ' Ἡσίοδον.

"Απαντα μέν οὖν φιλόζωα τὰ κεφαλόρριζα μάλιστα δ' ἡ σκίλλα· καὶ γὰρ κρεμαννυμένη ζῆ καὶ πλεῖστόν γε χρόνον διαμένει· δύναται δὲ καὶ τὰ θησαυριζόμενα σώζειν, ὥσπερ τὴν ῥόαν ἐμπηγνυμένου τοῦ μίσχου, καὶ τῶν φυτευομένων δ' ἔνια βλαστάνει θᾶττον ἐν αὐτῆ· λέγεται δὲ καὶ πρὸ τῶν θυρῶν τῆς εἰσόδου φυτευθεῖσαν ἀλεξητήριον εἶναι τῆς ἐπιφερομένης δηλήσεως. πάντα δὲ ταῦτα ἀθρόα φύεται, καθάπερ καὶ τὰ κρόμυα καὶ τὰ σκόροδα· παραβλαστάνουσι γὰρ ἀπὸ τῆς

1 ἐκπίπτει conj. W.; ἐκπίπτων Ald.

 $^{^2}$ ἄνθησιν conj. Scal.; ἄναυσιν corr. to αδανσιν U; ἄναυσιν M; αδανσιν Ald.; floret per partes G.

ENQUIRY INTO PLANTS, VII. xIII. 2-4

has the tallest of all-for the antherikos (asphodelstalk) is very tall: that of iris is smaller, but tougher, though in general it is like the asphodel-stalk. Asphodel also produces much fruit, and its fruit is woody: in shape it is triangular and in colour black: it is found in the round vessel which is below the flower, and it falls out 1 in summer when this splits open. It does not produce all its flowers 2 at once; in which respect it resembles squill, but the flowering begins at the bottom. In the stalk of asphodel forms a grub which changes into another creature like a hornet, and then, when the stem withers, eats its wav out and flies away. A peculiarity of the plant as compared with others which have a smooth stem appears to be that, though it is slender, it has outgrowths at the top. It provides many things useful for food: the stalk is edible when fried, the seed when roasted, and above all the root 8 when cut up with figs; in fact, as Hesiod says,4 the plant is extremely profitable.

Now all bulbous plants are tenacious of life, but especially squill; for this even lives when hung up and continues to do so for a very long time; it is even able to keep other things that are stored, for instance the pomegranate, if the stalk of the fruit is set in it; and some cuttings 5 strike more quickly if set in it; and it is said that, if planted before the entrance door of a house, it wards off mischief 6 which threatens it. All these bulbs grow in masses, as do onions and garlic; for they make offsets from the root, and some

129

ĸ

VOL. II.

³ cf. 7. 9. 4; 9. 9. 6.

⁴ Hes. Op. 41.

⁵ cf 2. 5. 5; C.P. 5. 6. 10.

⁶ Sc. witchcraft. veneficiorum noxam Plin. 21. 108.

ρίζης· ἔνια δὲ καὶ ἀπὸ τοῦ σπέρματος φανερῶς, οἷον ὅ τε ἀνθέρικος καὶ τὸ λείριον καὶ τὸ φάσγανον καὶ ὁ βολβός.

- ΄Αλλ΄ ίδιον τοῦτο τοῦ βολβοῦ λέγεται, τὸ μὴ ἀπὸ πάντων βλαστάνειν ἄμα τῶν σπερμάτων, ἀλλὰ τοῦ μὲν αὐτοετὲς τοῦ δ΄ εἰς νέωτα, καθάπερ τὸν αἰγίλωπά φασι καὶ τὸν λωτόν. τοῦτο μὲν οὖν εἴπερ ἀληθὲς κοινὸν ἑτέρων. κοινὸν δὲ ἴσως καὶ τὸ μέλλον λέγεσθαι, πλὴν οὖ πολλῶν, θαυμαστὸν δὲ ἐπὶ πάντων, ὅπερ ἐπί τε τῆς σκίλλης καὶ τοῦ ναρκίσσου συμβαίνει τῶν μὲν γὰρ ἄλλων καὶ τῶν έξ ἀρχῆς φυτευομένων καὶ τῶν βλαστανόντων καθ΄ ὥραν ἔτους τὸ φύλλον ἀνατέλλει πρῶτον, εἰθ' ὕστερον ὁ καυλός ἐπὶ δὲ τούτων ὁ καυλὸς πρότερον.
- Τοῦ ναρκίσσου δὲ ὁ τοῦ ἄνθους μόνον εὐθὺ προωθῶν τὸ ἄνθος· τῆς δὲ σκίλλης καθ' αὐτόν, εἰς ὕστερον ἐπὶ τούτῷ τὸ ἄνθος ἀνίσχον προσκαθήμενον· ποιεῖται δὲ τὰς ἀνθήσεις τρεῖς, ὧν ἡ μὲν πρώτη δοκεῖ σημαίνειν τὸν πρῶτον ἄροτον, ἡ δὲ δευτέρα τὸν μέσον, ἡ δὲ τρίτη τὸν ἔσχατον· ὡς γὰρ ᾶν αὖται γένωνται καὶ οἱ ἄροτοι σχεδὸν οὕτως ἐκβαίνουσιν· ὅταν δὲ οὖτος ἀπογηράση, τότε ἡ τῶν φύλλων βλάστησις πολλαῖς ἡμέραις ὕστερον· ὡσαύτως δὲ καὶ ἐπὶ τοῦ ναρκίσσου, πλὴν οὕτε καυλὸν ἔτερον ἔχει παρὰ τὸν τοῦ ἄνθους, ὅσπερ εἰπομεν, οῦτε καρπὸν φανερόν, ἀλλ

¹ αἰγίλωπα conj. Sch. from Plin. 21. 103; γίλωπα UMAld.
² εὐθὸ προωθῶν conj. W.; εὐθυπρόωρον Ald. H. cf. Plin. 21.
66, where however the statement is transferred to the crocus.

ENQUIRY INTO PLANTS, VII. xIII. 4-6

plainly are also increased by seed, as the asphodel polyanthus narcissus corn-flag and purse-tassels.

However it is said to be a peculiarity of pursetassels that all the seeds do not germinate at once, but some in the same year, some the next year; a like account is given of aigilops 1 and trefoil. If then this is true, it is not peculiar to this plant. Nor perhaps is the following characteristic, which is not found in many plants and is marvellous wherever it does occur—and it is found in squill and narcissus: namely that, whereas in most plants, whether those originally planted or those which are produced from them in season, the leaf comes up first and then presently the stem, in these plants the stem comes up first.

In the case of narcissus it is only the flower-stem which comes up, and it immediately pushes up ² the flower. But in squill it is the stem ³ proper which thus appears, and presently the flower appears emerging ⁴ from and sitting on it. And it makes three flowerings, ⁵ of which the first appears to mark the first seed-time, the second the middle one, and the third the last one; for, according as these flowerings have occurred, ⁶ so the crops usually turn out. But, when the flower-stem ⁷ has waxed old, then the growth of the leaves follows many days later. So also is it with narcissus, except that it has no second stem besides the flower-stem, as we said, nor any visible fruit; but the flower itself

³ i.e. the whole 'bud.'

⁴ ἀνίσχον Ald.; ἀνίσχων conj. Sch. followed by W.

⁵ Plin. l.c.; cf. 18. 237.

⁶ i.e. the flowering is the sign when to sow. The same is said of the fruiting of $\sigma \chi \hat{v}$ vos de signis 55.

⁷ οὖτος conj. Sch.; οὕτως Ald.

αὐτὸ τὸ ἄνθος ἄμα τῷ καυλῷ καταφθίνει καὶ

όταν αὐανθη τότε τὰ φύλλα ἀνατέλλει.

Πρὸς μὲν οὖν τὰ ἄλλα τὰ συνάμφω ταῦτα ἴδια· πρὸς δὲ τὰ προανθοῦντα τῶν Φύλλων καὶ τῶν καυλών, όπερ δοκεί ποιείν το τίφυον και έτερα των ανθικών, έτι τε των δένδρων ή αμυγδαλή μάλιστα ή μόνον, δτι ταῦτα μὲν ἄμα τῷ ἄνθει προφαίνει τὸ φύλλον ἡ εὐθὺς κατόπιν, ὥστε καὶ διαζητεισθαι περί τινων, έπὶ δὲ τούτων οίον ἀφ' έτέρας άρχης φαίνεται καὶ διὰ τὸ πληθος τών ήμερῶν καί διὰ τὸ μὴ πρότερον βλαστάνειν πρὶν τοῦ μεν τὸ ἄνθος τοῦ δὲ καὶ ὁ καυλὸς ὅλος ἀπογηράση. ή δὲ βλάστησις προτέρα μὲν τῆς σκίλλης, ύστέρα δὲ τοῦ ναρκίσσου πολύ δὲ πλέον τὸ φύλλον οὖτος ἀφίησι, καί ἐστιν ἡ ῥίζα αὐτὴ μικρὰ καὶ οὐ μεγάλη, προσεμφερής δὲ κατὰ τὸ σχήμα τῶ βολβῶ, πλὴν <οὐ> λεπυριώδης, ταῦτα μὲν οὖν ἔγει σκέψιν.

Τῶν δὲ βολβῶν ὅτι πλείω γένη φανερόν, καὶ γὰρ τῷ μεγέθει καὶ τῆ χρόα καὶ τοῖς σχήμασι διαφέρουσι καὶ τοῖς χυλοῖς ἐνιαχοῦ γὰρ οὕτω γλυκεῖς ιστε καὶ ἀμοὺς ἐσθίεσθαι, καθάπερ ἐν Χερρονήσω τῆ Ταυρικῆ. μεγίστη δὲ καὶ ἰδιωτάτη διαφορὰ τῶν ἐριοφόρων ἔστι γάρ τι γένος τοιοῦτον, ὁ φύεται μὲν ἐν αἰγιαλοῖς ἔχει δὲ τὸ ἔριον ὑπὸ τοὺς πρώτους χιτῶνας, ιστε ἀνὰ μέσον εἰναι

 $^{^1}$ T(quov Ald., cf. C.P. 1. 10. 5; 7' Iquov conj.W.; iphyum GBas.Par. cf. 6. 6. 11. 2 7 add. Sch.

³ êπl conj. H.; περl UMAld. 4 W. adds av.

⁵ αὐτη : sc. apart from offsets.
⁶ μικρὰ conj. Sch.; οὐ μικρὰ Ald.

ENQUIRY INTO PLANTS, VII. xIII. 6-8

perishes with the stem, and when it has withered,

then the plant puts up its leaves.

These two plants then, as compared with the other bulbous plants are peculiar; and, as compared with those which bloom before the leaves and stems appear (as the autumn squill 1 seems to do, and other plants with conspicuous flowers, as well as, among trees, the almond especially, if not alone), there is the distinction that, while these two put forth their leaves along with the flowers or 2 immediately afterwards (so that about some the matter is uncertain) in 3 the case of these two the flower appears, as it were, from a different starting-point, there being a considerable number of days in between, and the growth of the leaves not beginning till,4 in the case of one of them, the flower, and in the case of the other, the whole stem has withered. Squill produces its leaves before the flower, narcissus afterwards: but the latter produces much more abundant foliage. and the individual 5 root is small 6 rather than large, resembling purse-tassels in shape, except that it is not formed of scales.7 About these matters then there is doubt.

Of purse-tassels it is plain that there are several kinds; for they differ in size colour shape and taste. ⁸ In some places they are so sweet as to be eaten raw, as in the Tauric Chersonese. But the greatest and most distinct difference is shown by the 'wool-bearing' purse-tassels; for there is such a kind, and it grows on ¹⁰ the sea-shore, and has the wool beneath the outer tunic, so that it is between

⁸ Plin. 19. 95; Athen. 2. 64.

 $^{^{7}}$ οὐ λεπυριώδης conj. Sch. from G, non squamata; οὐδὲ πυρώδη UMAld; οὐ λεπυρώδης Η.

Plin. 19. 32. See Index. 10 $\ell \nu$ after $\mu \hat{\epsilon} \nu$ add. W.

τοῦ τε ἐδωδίμου τοῦ ἐντὸς καὶ τοῦ ἔξω· ὑφαίνεται δὲ ἐξ αὐτοῦ καὶ πόδεια καὶ ἄλλα ἰμάτια· δι' δ καὶ ἐριῶδες τοῦτο καὶ οὐχ ὥσπερ τὸ ἐν Ἰνδοῖς τριχῶδες.

Πλείω δὲ καὶ τὰ βολβώδη καὶ ἐλάττω. ταῦτα δὲ . . . καθάπερ τὸ λευκόῖον καὶ βολβίνη καὶ ὀπιτίων καὶ κύϊξ καὶ τρόπον τινὰ τὸ σισυρίγχιον. βολβώδη δὲ ταῦτα ὅτι στρογγύλα ταῖς ῥίζαις ἐπεὶ τοῖς γε χρώμασι λευκὰ καὶ οὐ λεπυριώδη. ἴδιον δὲ τοῦ σισυριγχίου τὸ τῆς ῥίζης αὐξάνεσθαι τὸ κάτω πρῶτον, ὁ καλοῦσι . . . χειμῶνα, τοῦ δ' ἤρος ὑποφαίνοντος τοῦτο μὲν ταπεινοῦσθαι τὸ δ' ἄνω τὸ ἐδώδιμον αὐξάνεσθαι. καὶ τὰ μὲν τοιαύτας ἔχει τὰς διαφοράς.

ΧΙΥ. "Ιδια δὲ καὶ ταῦτα ἐν τοῖς ποιώδεσιν, οἶον τό [τε] ἐπὶ τοῦ ἀδιάντου συμβαῖνον οὐδὲ γὰρ ὑγραίνεται τὸ φύλλον βρεχόμενον οὐδὶ ἐπίδροσόν ἐστι διὰ τὸ μὴ τὴν νοτίαν ἐπιμένειν, ὅθεν καὶ ἡ προσηγορία. γένη δὲ αὐτοῦ δύο, τὸ μὲν λευκὸν τὸ δὲ μέλαν, χρήσιμα δ' ἀμφότερα πρὸς ἔκρυσιν κεφαλῆς τριχῶν ἐν ἐλαίφ τριβόμενα. φύεται δὲ

¹ δι' δ καὶ έρ. τοῦτο: text probably defective. ? δι' δ καὶ $< \chi \rho \eta \sigma \iota \mu \delta \nu \ \tau \delta > \epsilon \rho \iota \hat{\omega} \delta \epsilon s \ \tau οῦτο:$ 'wherefore this woolly kind is serviceable, which the Indian hairy kind is not.'

² Plin. 19. 95.

³ καὶ ἐλάττω· ταῦτα δὲ: text corrupt and defective.

^{*} ὀπιτίων Η.; ὁ πιτίων Ald.; pithyon Plin. l.c.; ὀπιτίων and κύτξ were possibly earth-nuts.

⁵ γε conj. Sch ; τε Ald

ENQUIRY INTO PLANTS, VII. xiii. 8-xiv. 1

the edible inside and the outside: of it are woven felt shoes and other articles of apparel. Wherefore ¹ this kind is woolly and distinct from the Indian kind,

which is hairy.

² There are also several kinds of plants of the same class as purse-tassels ³ such as snowdrop star-flower opition ⁴ kyix, and to a certain extent Barbary nut. These belong to this class only in having round roots; for in colour ⁵ they are white, and the bulbs are not formed of scales. A peculiarity of Barbary nut is that the lower end of the root grows first, and this is called; it grows ⁶ during winter, but, when spring appears, it decreases, while the upper part, which is edible, grows. Such are the differences in these plants.

Of certain properties and habits peculiar to certain herbaceous plants.

XIV: There are also the following peculiarities in herbaceous plants, for instance that ⁷ which we find in 'wet-proof' (maidenhair); ⁸ the leaf does not even get wet when it is watered, nor does it catch the dew, ⁹ because the dew does not ¹⁰ rest on it; whence its name. ¹¹ There are two kinds, the white 'wet-proof' (English maidenhair), and the black (maidenhair); and both are useful to prevent the falling off of the hair of the head, for which purpose they are pounded up and mixed with olive-oil. They grow

⁷ I have bracketed τε. 8 Plin. 22. 62-65.

10 μη before την add. W.

^{6 &#}x27;Grows' supplied from G and Plin. l.c., who have no trace of δ καλοῦσι.

 $^{^{\}circ}$ ἐπίδροσον conj. W.; ἐπίδηλον UP2MAld.; nec quicquam adhaesisse humoris constat G.

¹¹ Plin. l.c.; 27. 138; 25. 132.

μαλιστα πρὸς τὰ ὑδρηλά. ὡς δὲ οἴονταί τινες, καὶ πρὸς στραγγουρίαν τὸ τριχομανὲς ποιεῖ· ἔχει δὲ τὸν καυλὸν ὅμοιον τῷ ἀδιάντῷ τῷ μέλανι, φύλλα δὲ μικρὰ σφόδρα καὶ πυκνὰ καὶ πεφυκότα καταντικρὺ ἀλλήλων, ῥίζα δὲ οὐχ ὕπεστι· χωρία δὲ φιλεῖ σκιερά.

Των δε κατα μέρος ανθούντων ίδιον το περί το ανθεμον, ότι των μεν άλλων πάντων τα κάτω πρώτον άπανθει τούτου δε τα άνω· τυγχάνει δ' αὐτοῦ τὸ μεν κύκλω τὸ λευκὸν ἄνθος τὸ δε ἐν τῷ μέσῷ τὸ χλωρόν· καὶ καρπὸς δς ἐκπίπτει, καθάπερ τοις ἀκανθώδεσι, καταλιπων τὴν πρόσφυσιν κενήν· εἴδη δ' αὐτοῦ πλείω.

*Ίδιον δὲ καὶ τὸ περὶ τὴν ἀπαρίνην, ἢ καὶ τῶν ἱματίων ἀντέχεται διὰ τὴν τραχύτητα καί ἐστι δυσαφαίρετον· ἐν τούτω γὰρ ἐγγίνεται τῷ τραχεῖ τὸ ἄνθος οὐ προϊὸν οὐδὲ ἐκφαῖνον ἀλλ' ἐν ἑαυτῷ πεττόμενον καὶ σπερμογονοῦν· ὥστε παρόμοιον εἶναι τὸ συμβαῖνον ὥσπερ ἐπὶ τῶν γαλεῶν καὶ ῥινῶν· ἐκεῖνά τε γὰρ ἐν ἑαυτοῖς ὡστοκήσαντα ζωογονεῖ, καὶ αὕτη τὸ ἄνθος ἐν ἑαυτῆ κατέχουσα καὶ πέττουσα καρποτοκεῖ.

XV. "Όσα δὲ τὰς ἀνθήσεις λαμβάνουσι ἀκολουθοῦντα τοῖς ἄστροις, οἶον τὸ ἡλιοτρόπιον καλούμενον καὶ ὁ σκόλυμος, ἄμα γὰρ ταῖς τροπαῖς καὶ οὖτος, ἔτι δὲ τὸ χελιδόνιον, καὶ γὰρ τοῦτο ἄμα τῷ χελιδονία ἀνθεῖ, ταῦτα δὲ δόξειεν ἄν τῆ μὲν φυσικὴν ἔχειν τὴν αἰτίαν τῆ δὲ συμπτωματικήν.

¹ i.e. the white kind. Sch. followed by G adds τδ καλ τριχομανές καλούμενον after τδ μέν λευκόν above.

ENQUIRY INTO PLANTS, VII. xiv. 1-xv. 1

especially in damp places. Some think that trikhomanes 1 (English maidenhair) is also useful in cases of strangury. Its stem is like that of the black kind. but it has small leaves, which are close set and grow in opposite pairs; there is no root below, and the plant loves shady places.

Of those plants which do not flower all at once anthemon has the peculiarity that, while in all others 2 the lower part flowers first, in this plant it is the upper part which does so; the outer circle of the flower is white.8 and the centre green4; and the fruit falls off, as in spinous plants, leaving the attach-

There are several forms of it. ment bare.

⁵ Bedstraw has the peculiarity that it sticks to clothes owing to its roughness, and it is hard to pull away: indeed it is in this rough part that the flower is contained: it does not project nor show, but matures within itself and produces seed; so that its habit is like that of weasels and sharks; for, as these animals 6 likewise produce eggs in themselves and then bear their young alive, so this plant keeps its flower within itself, matures it and produces fruit.

XV. 7 As to these plants whose flowering time is dependent on the heavenly bodies,8 as the plant called heliotropion, golden thistle (for this also blooms at the solstice), and also 'swallow-plant' (greater celandine)—for this blooms when the 9 Swallow-wind blows—the reason in these cases would seem to be

partly in their nature and partly accidental.

 $^{\circ}$ $\tau \hat{\varphi}$ conj. Sch.; $\tau \hat{\eta}$ MAld. cf. Plin. 2. 122.

² Ότον after πάντων om. W. after Sch.

^{4 ?} om. τδ before χλωρόν. 3 το λευκον: ? λευκον το.

Plin. 21. 104.
 cf. Arist. H.A. 6. 11.
 Athen. 15. 32.
 Δατροις conj. St.; ἀγρίοις Ald.

Πολλά δὲ τοιαῦτά ἐστι καὶ ἐν ἑτέροις ἴδια· οἰον καὶ ἡ τοῦ ἀειζώου φύσις τὸ διαμένειν ὑγρὸν ἀεὶ καὶ χλωρόν, φύλλον σαρκῶδες ἔχον καὶ λεῖον καὶ πρόμηκες. φύεται δὲ ἔν τε τοῖς άλιπέδοις τοῖς τε ἐπὶ τῶν τειχῶν ἀνδήροις καὶ οὐχ ἡκιστα ἐπὶ τῶν κεράμων, ὅταν ἐπιγένηται γῆς τις ἀμμώ-

δης συρροή.

Πολλά δ' ἄν τις ἴσως λάβοι καὶ ἔτερα περιττά. χρὴ δέ, ὥσπερ πολλάκις εἴρηται, τὰς ἰδιότητας θεωρεῖν καὶ τὰς διαφορὰς πρὸς τὰ ἄλλα. τὰ μὲν ἐν πλείοσιν ἰδέαις ἐστὶ καὶ σχεδὸν οἶον ὁμωνύμοις, ὥσπερ ὁ λωτός· τούτου γὰρ εἴδη πολλὰ διαφέροντα καὶ φύλλοις καὶ καυλοῖς καὶ ἄνθεσι καὶ καρποῖς, ἐν οἶς καὶ ὁ μελίλωτος καλούμενος· καὶ δυνάμει δὲ τῆ κατὰ τὴν προσφοράν, ἔτι τε τῷ μὴ τοὺς αὐτοὺς τόπους ζητεῖν. ὁμοίως δὲ καὶ ἔτερα πλείω.

Τὰ δὲ ἐν ἐλάττοσιν, ὅσπερ ὁ στρύχνος ὁμωνυμία τινὶ παντελῶς εἰλημμένος · ὁ μὲν γὰρ ἐδώδιμος καὶ ὅσπερ ἤμερον, καρπὸν ἔχων ῥαγώδη, ἔτεροι δὲ δύο εἰσίν, ὧν ὁ μὲν ὕπνον ὁ δὲ μανίαν ἐμποιεῖν δύναται, πλείων δ' ἔτι δοθεὶς καὶ κτείνει. ὁμοίως δὲ τοῦτο καὶ ἐφ' ἔτέρων ἐστὶ λαβεῖν, ἃ πολλὴν ἔχει διάστασιν. περὶ μὲν οὖν τῶν ἄλλων τῶν ποιωδῶν ἱκανῶς εἴρηται. περὶ δὲ τοῦ σίτου καὶ τῶν σιτωδῶν μετὰ ταῦτα λεκτέον τοῦτο γὰρ ἔτι κατάλοιπον ἦν.

¹ olov conj. W.; διὸ Ald.

² άλιπέδοις conj. Sch.; άληπέδοις U; άληπέδοις M; άλοπέδοις Ald. cf. Xen. Hell. 2. 4. 30.

³ τε after τοιs add. W. after Lobeck.

⁴ e.g. λωτός and μελίλωτος. See Index, λωτός.

⁵ μελίλωτος conj. Bod.; μελίας σίτος Ald.

ENQUIRY INTO PLANTS, VII. xv. 2-4

Such peculiarities are common in other plants also; thus ¹ it is the nature of the house-leek to remain always moist and green, its leaf being fleshy smooth and oblong. It grows on flat shores,² on the ³ earthy tops of walls, and especially on tiled roofs, when there is on them a sandy accumulation of earth.

Possibly one might mention many other eccentricities. But, as has been repeatedly said, we must only observe the peculiarities and differences which one plant has as compared with others. Some plants are found in several forms which have almost 4 the same name, for instance the lotos; for of this there are many forms differing in leaves stems flowers and fruit, including the plant called melilotos 5; there are also forms differing in the virtues for which 6 they are used as food, and again in their fondness for different localities. So too is it with many other plants.

plants.

Others are found in fewer forms, as strykhnos,7 which is a general name covering plants that are quite distinct; one is edible and like a cultivated plant, having a berry-like fruit, and there are two others,8 of which the one is said to induce sleep, the other to cause madness, or, if it is administered in a larger dose, death. The same thing may be observed in other plants which are widely different. Now about the other herbaceous plants enough has been said; but concerning corn and corn-like plants we must speak next; for this subject still lies before us.

6 δὲ τῆ conj. Sch.; διτταὶ UM; διτταῖs Ald.

⁷ cf. 7. 7. 2; Plin 21. 177-179; Diosc. 4. 70-73; Index. 8 In 9. 11. 5 these two plants are said to be συνώνυμοι, i.e. different forms of the same plant, whereas the 'edible' στούχνος is the same only in name (δμωνυμία). cf. 9. 12. 5.

BOOK VIII

Ι. Περὶ μὲν οὖν τῶν ἄλλων ποιωδῶν ἱκανῶς εἰρήσθω· περὶ δὲ σίτου καὶ τῶν σιτωδῶν λέγωμεν ὁμοίως τοῖς πρότερον· τοῦτο γὰρ κατάλοιπον ἦν τῶν ποιωδῶν.

Δύο δὲ αὐτοῦ γένη τὰ μέγιστα τυγχάνει· τὰ μὲν γὰρ σιτώδη, οἶον πυροὶ κριθαὶ τίφαι ζειαὶ τὰ ἄλλα τὰ ὁμοιόπυρα ἡ ὁμοιόκριθα· τὰ δὲ χεδροπά, οἶον κύαμος ἐρέβινθος πισὸς καὶ ὅλως τὰ ὅσπρια προσαγορευόμενα· τρίτον δὲ παρ' αὐτὰ κέγχρος ἔλυμος σήσαμον καὶ ἀπλῶς τὰ ἐν τοῖς θερινοῖς ἀρότοις ἀνώνυμα κοινῆ προσηγορίą.

2 Έστι δὲ ἡ μὲν γένεσις αὐτῶν μία καὶ ἁπλη· φύεται γὰρ ἀπὸ σπέρματος, ἐὰν μή τι σπάνιον καὶ ὀλίγον ἀπὸ τῆς ῥίζης. ὡραι δὲ τοῦ σπόρου τῶν πλείστων δύο· πρώτη μὲν καὶ μάλιστα ἡ περὶ Πλειάδος δύσιν, ἢ καὶ Ἡσίοδος ἡκολούθηκε

¹ τῶν before ποιωδῶν om. Sch.

² Plin. 18. 48-80.

³ ξλυμος: μελίνη appears to be the Attic name for this plant. Sch. would restore it for ξλυμος here and 4. 4. 10; 8, 11, 1.

¹⁴²

BOOK VIII

OF HERBACEOUS PLANTS: CEREALS, Pulses, and 'Summer Crops.'

Of the three classes and the times of sowing and of germination.

I. Let the above suffice for an account of the other herbaceous 1 plants; let us now discuss corn and corn-like plants in the same manner as those already treated; for this class of herbaceous plants we reserved.

² There are two principal classes; there are the corn-like plants such as wheat barley one-seeded wheat rice-wheat and the others which resemble either of the first two; and again there are the leguminous plants, as bean chick-pea pea, and in general those to which the name of pulses is given. Besides these there is a third class, which includes millet ³ Italian millet, sesame and in general the plants which belong to the summer seed-time, ⁴ which lack any common designation.

There is only one single way of propagating these; they grow from seed, except that some may grow rarely and scantily from a root. There are two seasons for sowing most of them; the first and most important is about the setting of the Pleiad 5; this rule we find even Hesiod 6 following with

⁶ Hes. Op. 383.

⁴ cf. 8. 7. 3.

⁵ Πλειάδος conj. Sch.; πλειάδας U; πλειάδων Ald.

καὶ σχεδὸν οἱ πλεῖστοι, δι' δ καὶ καλοῦσί τινες αὐτὴν ἄροτον ἄλλη δ' ἀρχομένου τοῦ ἦρος μετὰ τὰς τροπὰς τοῦ χειμῶνος. οὐ τῶν αὐτῶν δὲ ἐκατέρα. τὰ μὲν γὰρ αὐτῶν φιλεῖ πρωϊσπορεῖσθαι, τὰ δὲ ὀψὲ διὰ τὸ μὴ δύνασθαι φέρειν τοὺς χειμῶνας, τὰ δὲ πρὸς ἀμφοτέρας τὰς ὥρας οὐ κακῶς ἔχει, καὶ πρὸς χειμῶνα καὶ πρὸς ἔαρ.

3 Πρωΐσπορα μὲν οὖν έστι πυρὸς κριθή, καὶ τούτων ἡ κριθὴ πρωϊσπορώτερον· ἔτι δὲ ζειὰ τίφη ὀλύρα καὶ εἴ τι ἔτερον ὁμοιόπυρον· ἀπάντων γὰρ σχεδὸν ὁ αὐτὸς χρόνος τῆς σπορᾶς· τῶν δὲ χεδροπῶν μάλιστα ὡς εἰπεῖν κύαμος καὶ ὧχρος· ταῦτα γὰρ διὰ τὴν ἀσθένειαν προλαβεῖν τῆ ῥιζώσει βούλεται τοὺς χειμῶνας· πρωΐσπορον δὲ καὶ ὁ θέρμος· ἀπὸ τῆς ἄλω γάρ φασι δεῖν καταβάλλειν εὐθύς.

'Οψίσπορα δὲ τούτων γε αὐτῶν ὅσα διαφέρει τοῖς γένεσιν, οἶον πυρῶν τέ τι γένος καὶ κριθῶν ὁ καλοῦσι τρίμηνον διὰ τὸ ἐν τοσούτῷ τελειοῦσθαι· καὶ τῶν χεδροπῶν τὰ τοιάδε, φακὸς ἀφάκη πισός. ἐν ἀμφοτέραις δὲ ταῖς ὥραις τῶν χεδροπῶν, καθάπερ ὅροβος ἐρέβινθος· οἱ δὲ καὶ τὸν κύαμον ὀψὲ σπείρουσιν, ἐὰν ὑστερήσωσι τῶν πρώτων ἀρότων. ἀπλῶς δὲ πρωϊσποροῦσι τὰ μὲν δι' ἰσχὸν ὡς δυνάμενα φέρειν τοὺς χειμῶνας, τὰ δὲ δι ἀσθένειαν, ὅπως προλάβωσι ταῖς εὐδίαις τὴν αὕξησιν. δύο μὲν οὖν αὖται. τρίτη δὲ τῶν θερινῶν ἡν

² τῶν δέ γε Ald.; γε om. Sch.

¹ A cultural variety of Geid. cf. 8. 9. 2.

⁸ δχρος conj. W.; κέγχρος Ald.; om. G. cf. 8. 3. 1 and 2. 4 cf. 8. 11. 8.

¹⁴⁴

ENQUIRY INTO PLANTS, VIII. 1. 2-4

most authorities; wherefore some call it simply 'the seed-time.' Another time is at the beginning of spring after the winter equinox. However different crops are sown at the two seasons. For some of them love to be sown early, some late because they cannot bear the winters, while some will do not amiss at either season, both towards winter and towards spring.

Crops sown early are wheat and barley, and of these the latter is sown the earlier; also rice-wheat one-seeded wheat olyra, and others which resemble wheat. For all of these the time of sowing is about the same. Of leguminous plants 2 bean and okhros,8 it may be said, are specially sown at this time; for these on account of their weakness like to be well rooted before the winter. Lupin is also sown early; in fact they say it should be sown straight from the threshing-floor.4

Those which are sown late are certain special varieties 5 of these very kinds, as a certain kind of wheat, and of barley the kind which is called 'three months barley' because it takes that time to mature; and among leguminous plants lentil tare pea. However some of these plants are sown at both seasons, as vetch and chick-pea; some also sow beans late, if they have missed the first seed-time. To speak generally, some crops are sown early because of their robustness, since they can stand the winters, some because of their weakness, so that their growth may be secured 6 in the fine weather. These then are the two seasons; the third is that of the summer crops of which we

145

VOL, II.

⁵ τοις γένεσιν: τῆ γενέσει W. i.e. 'certain kinds which differ as to their germination.' 6 cf. C.P. 4, 7, 2.

εἴπομεν, ἐν ἡ κέγχρος σπείρεται καὶ μέλινος καὶ σήσαμον, ἔτι δ' ἐρύσιμον καὶ ὅρμινον. χρόνοι μὲν οὖν ἑκάστων οὖτοι.

Βλαστάνει δὲ τὸ μὲν θᾶττον τὸ δὲ βραδύτερον καὶ κριθὴ μὲν καὶ πυρὸς ἐβδομαῖα μάλιστα· προτερεῖ δὲ ἡ κριθὴ μᾶλλον· τὰ δ' ὄσπρια τεταρταῖα ἡ πεμπταῖα πλὴν κυάμων· κύαμος δὲ καὶ τῶν σιτωδῶν ἔνια πλείοσιν· ἐνιαχοῦ γὰρ καὶ πεντεκαιδεκαταῖος, ὁτὲ δὲ καὶ εἰκοσταῖος· δυσφυέστατον γὰρ τοῦτο πάντων, ἐὰν δὲ δὴ καὶ σπαρέντος ἐπὶ πλέον ὕδωρ ἐπιγένηται, καὶ παντελῶς. εἰ δὲ τῶν ἐν τοῖς ἠρινοῖς ἀρότοις θᾶττον ἡ ἔκφυσις διὰ

την ώραν σκεπτέον.

Χρή δὲ τὰς ἀναβλαστήσεις καὶ τὰς διαφύσεις ταύτας ὡς ἐπὶ τὸ πᾶν διαλαβεῖν· ἐνίστε γὰρ ἐνιαχοῦ καὶ ἐν ἐλάττοσιν ἡμέραις, καθάπερ ἐν Αἰγύπτω <κριθή>· τριταίαν γάρ φασι καὶ τεταρταίαν ἀνατέλλειν· παρ' ἄλλοις δὲ ἐν πλείοσι τῶν εἰρημένων, ὅπερ καὶ οὐκ ἄλογον, ὅταν καὶ χώρα καὶ ἀὴρ διαφέρη καὶ πρωϊαίτερον ἡ ὀψιαίτερον ἀρόση καὶ τὰ ἐπιγινόμενα ἀνόμοια τυγχάνη. ἡ μὲν γὰρ μανὴ καὶ κούφη καὶ εὐκράτω ἀέρι ταχὸ καὶ ἡαδίως ἀναδίδωσιν, ἡ δὲ γλίσχρα καὶ βαρεῖα βραδέως, ἡ δὲ τοῖς τόποις αὐχμωδεστέρα βραδύτερον.

΄ Έτι δὲ ἄν χειμῶνες ἐπιγένωνται καὶ αὐχμοὶ καὶ εὐδίαι καὶ πάλιν ὕδατα· καὶ γὰρ ἐν τούτοις πολὺ παραλλάττουσιν. ὡσαύτως δὲ καὶ ἐὰν ἡ

8 cf. 7. 1. 4.

¹ žvia conj. W.; žv Ald.

² The reason is given C.P. 4. 8. 2.

ENQUIRY INTO PLANTS, VIII. 1. 4-7

spoke, in which are sown millet Italian millet sesame, and also erysimon and horminon. Such then are the times for each.

Some are quicker in coming up, some slower. Barley and wheat generally come up on the seventh day, but barley is the earlier. Pulses take four or five days, except beans; for they, like some kinds 1 of corn, require a longer time; in some places they take as much as fifteen days, or even twenty. This crop indeed is the slowest to start of all, and if after the sowing there is a long spell of wet weather, it is extremely slow.2 Whether the sprouting 3 of crops sown at the spring seed-time is quicker because of the season is matter for enquiry.

These times of sprouting or germination must be taken generally; for at some times and places germination takes fewer days, as with barley 4 in Egypt, where it is said to come up on the third or fourth day: while elsewhere it takes longer than the period mentioned, which is not surprising when both soil and climate are different, when one makes the sowing earlier or later, and when the crop is subjected to different influences afterwards. open light soil with a favourable 5 climate produces quick and easy growth, while soil that is sticky and heavy tends to slow growth, and that of a specially dry district to slower growth still.

Moreover the time of growth is affected, according as storms supervene, or droughts, or fine weather or again rain; for these conditions make wide differences. So too it makes a difference if the

4 κριθή add. W.

⁵ εὐκράτω conj. Scal. from G (benigno caelo); εὐκάρπω Ald.

γη προειργασμένη καὶ κόπρον ἔχουσα τυγχάνη, καὶ ἐὰν μηδὲν τούτων· ἐπεὶ καὶ περὶ τὸ πρωϊσπορεῖν ἔκαστα καὶ ὀψισπορεῖν αὶ χῶραι διαφέρουσιν. ἔνιοι δὲ καὶ περὶ τὴν Ἑλλάδα πάντα πρωϊσπορεῖν εἰώθασι διὰ ψυχρότητα τῆς χώρας, ὅσπερ οἱ Φωκεῖς, ὅπως ἂν οἱ χειμῶνες μὴ νήπια καταλαμβάνωσιν.

ΙΙ. Βλαστάνει δὲ τὰ μὲν ἐκ τοῦ αὐτοῦ τὴν ρίζαν ἀφιέντα καὶ τὸ φύλλον, τὰ δὲ ἐκάτερον ἐξ ἐκατέρου τοῦ ἄκρου. πυρὸς μὲν οὖν καὶ κριθὴ καὶ τίφη καὶ ὅλως ὅσα σιτώδη πάντα ἐξ ἐκατέρου ὅσπερ ἐν τῷ στάχυϊ πέφυκεν, ἀπὸ μὲν τοῦ κάτω τοῦ παχέος τὴν ρίζαν ἀπὸ δὲ τοῦ ἄνω τὸν βλαστόν· ἐν δέ τι καὶ συνεχὲς γίνεται τὸ ἀμφοῦν τῆς τε ρίζης καὶ τοῦ καυλοῦ. κύαμος δὲ καὶ τὰ ἄλλα χεδροπὰ οὐχ ὁμοίως, ἀλλ' ἐκ τοῦ αὐτοῦ τὴν ρίζαν καὶ τὸν καυλόν, καθ' δ καὶ ἡ πρόσφυσις αὐτῶν ἐστι πρὸς τὸν λοβόν, ἐν ῷ καὶ ἔχουσιν οἶον ἀρχήν τινα φανεράν· ἐπ' ἐνίων δὲ καὶ αἰδοιῶδες φαίνεται, καθάπερ ἐπὶ τῶν κυάμων καὶ τῶν ἐρεβίνθων καὶ μάλιστα τῶν θέρμων· ἐκ τούτου γὰρ ἡ μὲν ρίζα κάτω τὸ δὲ φύλλον καὶ ὁ καυλὸς ἄνω χωρεῦ.

Ταύτη μὲν οὖν πη διαφέρει. τῆ δὲ ὁμοίως ἔχει τῷ πάντα κατὰ τὴν πρόσφυσιν τοῦ λοβοῦ καὶ τοῦ στάχυος ἀφιέναι τὴν ῥίζαν καὶ μὴ

ώς προείργ. Ald.H.; ώς om, Sch. from G.
 ² cf. 8. 8. 2.
 ³ ωσπερ conj. Scal.; πάντα Ald. (? repeated by mistake). cf. C.P. 4. 7. 4.

¹⁴⁸

ENQUIRY INTO PLANTS, VIII. 1. 7-11. 2

ground has been well tilled ¹ and given dung, or if neither of these things has been done: for the soil makes a difference even as to the early or late germination of each crop. In Hellas some are used to sow everything earlier because of the coldness of the soil, for instance the Phocians; ² the object being that the winter may not overtake the crop while it is still tender.

Of differences in the mode of germination and of subsequent development.

II. In germinating some of these plants produce their root and their leaves from the same point, some separately, from either end of the seed. Wheat barley one-seeded wheat, and in general all the cereals produce them from either end, in a manner corresponding to 3 the position of the seed in the ear, the root growing from the stout lower part, the shoot from the upper part; but the part corresponding to the root and that corresponding to the stem form a single continuous whole. Beans and other leguminous plants do not grow in the same manner, but they produce the root and the stem from the same point, namely the point at which the seed is attached to the pod, which, it is plain, is a sort of starting point of fresh growth. In some cases there is also a formation resembling the penis, as in beans chick-peas and especially in lupins; from this 4 the root grows downwards, the leaf and the stem upwards.

There are then these different ways of germinating; but a point ⁵ in which all these plants agree is that they all send out their roots at the place where

⁴ τούτου conj. Sch.; τούτων Ald. cf. C.P. 4. 7. 4. ⁵ cf. C.P. 4. 7. 7.

καθάπερ έν τοις δενδρικοίς τισιν ανάπαλιν, οίον άμυγδαλή καρύφ βαλάνφ τοις τοιούτοις. ἐν ἄπασι δὲ ἡ ῥίζα μικρῷ πρότερον ἐκφύεται τοῦ καυλοῦ· συμβαίνει δὲ ἔν γέ τισι τῶν δένδρων ώστε τὸν μὲν βλαστὸν ἐν αὐτῷ τῷ σπέρματι βλαστάνειν πρώτον, αὐξανομένου δὲ διίστασθαι τὰ σπέρματα—πάντα γάρ πως καὶ ταῦτα διμερῆ, τὰ δὲ δὴ χεδροπὰ φανερῶς πάντα δίθυρα καὶ σύνθετα την δε ρίζαν εὐθὺς έξω προωθεῖσθαι. έν δὲ τοῖς σιτηροῖς διὰ τὸ καθ' εν αὐτὰ είναι τοῦτο μὲν οὐ συμβαίνει, προτερεῖ δὲ ἡ ρίζα μικρόν.

Αναφύεται δε ή μεν κριθή και ο πυρος μονόφυλλα, ό δὲ πισὸς καὶ ὁ κύαμος καὶ ὁ ἐρέβινθος πολύφυλλα. ρίζαν δὲ ἔχει τὰ μὲν χεδροπὰ πάντα ξυλώδη καὶ μίαν ἀπὸ δὲ ταύτης καὶ ἀποφύσεις λεπτάς. βαθυρριζότατον δὲ ὡς εἰπεῖν τούτων ό ἐρέβινθος, ἐνίστε δὲ καὶ παρακαθίησιν άλλ' ό πυρὸς καὶ ἡ κριθὴ καὶ τὰ ἄλλα τὰ σιτώδη πολύρριζα καὶ λεπτόρριζα, δι' δ καὶ ταρρώδη. καὶ πολύκλαδα καὶ πολύκαυλα πάντα τὰ τοιαῦτα. σχεδον δε και εναντίωσίς τις εκατέρων εστί τά μεν γαρ χεδροπα μονόρριζα όντα πολλας άνωθεν άπὸ τῶν καυλῶν ἀποφύσεις ἔχει πλὴν κυάμου. τὰ δὲ σιτηρὰ πολύρριζα πολλούς μὲν ἀνίησι

βαλάνφ: διοσβαλάνφ Sch. from niBod.
 τισι τῶν δένδρων conj. W.; σιτώδεσιν UMAld.; τοῖς δενδρικοῖς conj. Sch. This and W.'s other conjectures in this section are rather desperate, but are accepted provisionally as at least restoring a satisfactory sense. The passage looks as if it had been deliberately tampered with by someone who misunderstood it.

ENQUIRY INTO PLANTS, VIII, II, 2-3

the seed is attached to the pod or ear, whereas the contrary is the case with the seeds of certain trees, as almond hazel acorn 1 and the like. And in all these plants the root begins to grow a little before the stem; whereas in certain trees 2 the bud first begins to grow within the seed itself, and, as it increases in size, the seeds split-for all such seeds are in a manner in two halves, and those of leguminous plants again all plainly have two valves and are double-and then the root is immediately thrust out; but in cereals,3 since the seeds are in one piece,4 this does not 5 occur, but the root grows a little before the bud.

Barley and wheat come up with a single leaf, but peas beans and chick-peas with several. 6 All the leguminous plants have a single woody root, and also slender side-roots springing from this. The chick-pea is about the deepest rooting of these, and sometimes it has side-roots; but wheat barley and the other cereals have a number of fine roots, wherefore they are matted together.8 Again all such plants have many branches and many stems. And there is a sort of contrast between these two classes; the leguminous plants, which have a single root, have many side-growths above from the stemall except beans; while the cereals, which have many roots, send up many shoots, but these have

³ σιτηροίε conj. W.; χεδροποίε UMAld.

⁴ καθ' ἐν αὐτὰ conj. W.; κατὰ τὸ αὐτὸ UMAld.

⁵ où conj. Scal. from G; our UMAld.

⁶ Plin. 18. 51.

⁷ Newrds conj. St.; Newral Ald. H.

^{*} ταρρώδη: cf. 6. 7. 4.

⁹ μèν conj. Sch.; γὰρ Ald. H.

βλαστούς, ἀπαράβλαστοι δὲ οὖτοι, πλὴν εἴ τι γένος πυρῶν τοιοῦτον, οὖς καλοῦσι σιτανίας καὶ

κριθανίας.

Τὸν μὲν οὖν χειμῶνα ἐν τῆ χλόη μένει τὰ σιτώδη, διαγελώσης δὲ τῆς ὥρας καυλὸν ἀφίησιν ἐκ τοῦ μέσου καὶ γονατοῦται. συμβαίνει δ' εὐθὺς ἐν τῷ τρίτῷ γόνατι, τοῖς δὲ ἐν τῷ τετάρτῷ, καὶ τὸν στάχυν ἔχειν ἀλλ' οὐ φανερὸν ἐν τῷ ὄγκῷ·—γίνεται δὲ ἐν τῷ ὅλῷ καλάμῷ πλείω τούτων·—ὥστε σχεδὸν ἄμα τῷ καλαμοῦσθαι συνίστασθαι <ἡ> μικρὸν ὕστερον· ἀλλ' οὐ πρότερον φανερὸς γίνεται πρὶν ἄν προαυξηθεὶς ἐν τῆ κάλυκι γένηται, τότε δὲ ἡ κύησις φανερὰ διὰ τὸν ὅγκον.

'Απολυθεὶς δ' εὐθὺς ἀνθεῖ μεθ' ἡμέρας τέτταρας ἡ πέντε καὶ πυρὸς καὶ κριθὴ καὶ ἀνθεῖ σχεδὸν τὰς ἴσας, οἱ δὲ τὰς πλείστας λέγοντες ἐν ταῖς ἑπτά φασιν ἀπανθεῖν. ἀλλὰ τῶν χεδροπῶν χρόνιος ἡ ἄνθησις· χρονιωτάτη δὲ τῶν μὲν ἄλλων ὀρόβου καὶ ἐρεβίνθου, τούτων δ' ἀπάντων τοῦ κυάμου καὶ ἐν μεγίστη διαφορᾶ· τετταράκοντα γὰρ ἡμερῶν ἀνθεῖν λέγουσι· πλὴν οἱ μὲν ἀεὶ παρανθοῦντος ἐτέρου καὶ ἐτέρου λέγουσι, κατὰ μέρος γὰρ ἀνθεῖν, οἱ δὲ ἀπλῶς. ἡ γὰρ ἄνθησις τῶν μὲν σταχυηρῶν ἀθρόως τῶν δὲ ἐλλοβωδῶν καὶ χεδροπῶν πάντων κατὰ μέρος· πρῶτα γὰρ ἀνθεῖ τὰ κάτω, καὶ ὅταν ταῦτα ἀπανθήση τὰ ἐχόμενα, καὶ οὕτως αἰεὶ βαδίζει πρὸς τὰ ἄνω.

¹ Plin. 18. 52. ² Plin. 18. 56. ³ cf. 7. 7. 1; 8. 4. 3.

ENQUIRY INTO PLANTS, VIII. 11. 3-5

no side-shoots—except such sorts of wheat as are called *sitanias* and *krithanias* ('barley-wheat').

¹ During winter cereals remain in the blade, but, as the season begins to smile, they send up a stem from the midst and it becomes jointed. ² And it comes to pass that the ear also at once appears in the third, or in some cases in the fourth joint, though it is not distinctly seen in the mass of growth (the whole stem contains more joints than three or four), so that it must be formed at the same time that the straw grows or but a little later; though it does not become conspicuous till it has first swollen and formed in the sheath,³ and by that time its size makes its development visible.

Four or five days after being set free 4 wheat and barley flower and remain in bloom for a like number of days; those who put the period at the longest say that the bloom is shed in seven days. 5 On the other hand the flowering period of leguminous plants lasts a long time; that of vetch and chick-pea is longer than that of most, but that of the bean is far longer than that of any of them; they say that it is in bloom for forty days; some however give this period absolutely, others say that at different times different parts are in flower, 6 since the whole plant does not flower at once. For plants with an ear bloom all at once, but plants with pods and all leguminous plants bloom part at a time; the lower part blooms first, and, when this bloom has fallen, the part next above it, and so on up to the top.

⁵ Plin. 18. 59.

⁴ Sc. from the sheath. ἀπολυθείς Ald.H.; ἀποχυθείς conj. Sch. followed by W. cf. ἀπόχυσις 8. 3. 4.

⁶ παρανθοῦντος conj. Η.; παρανθοῦντες Ald,

δι' δ πολλά των δρόβων τίλλεται τὰ μεν κάτω κατερρυηκότα τὰ δ' ἄνω χλωρὰ πάμπαν.

- Μετὰ δὲ τὴν ἀπάνθησιν άδρύνονται καὶ τελειοῦνται πυρὸς μὲν καὶ κριθὴ τετταρακοσταῖα μάλιστα παραπλησίως δὲ καὶ τίφη καὶ τἄλλα τὰ τοιαῦτα. τετταρακοσταῖον δέ φασι καὶ τὸν κύαμον, ὥστε ἐν ἴσαις ἀνθεῖν καὶ τελειοῦσθαι τὰ δ' ἄλλα ἐν ἐλάττοσιν ἐλαχίσταις δὲ ὁ ἐρέβινθος, εἴπερ ἀπὸ τῆς σπορᾶς ἐν τετταράκοντα τελειοῦται ταῖς ἀπάσαις ὥσπερ τινές φασιν ἐπεὶ τό γ' ὅλον ὅτι τάχιστα φανερόν. οἱ δὲ κέγχροι καὶ τὰ σήσαμα καὶ οἱ μέλινοι καὶ ὅλως τὰ θερινὰ σχεδὸν ὁμολογεῖται τὰς τετταράκονθ' ἡμέρας λαμβάνειν οἱ δὲ φασι καὶ ἐλάττους.
- Διαφέρει δὲ καὶ πρὸς τὴν τελείωσιν χώρα τε χώρας καὶ ἀὴρ ἀέρος· ἐν ἐλάττοσι γὰρ ἔνιαι δοκοῦσιν ἐκφέρειν, ὥσπερ ἄλλαι τε καὶ μάλιστα ἐπιδήλως Αἴγυπτος· ἐκεῖ γὰρ κριθαὶ μὲν ἐν ἐξαμήνω πυροὶ δὲ ἐν τῷ ἑβδόμω θερίζονται· περὶ δὲ τὴν Ἑλλάδα κριθαὶ μὲν ἐν τῷ ἑβδόμω παρὰ δὲ τοῖς πλείστοις ὀγδόω, πυροὶ δὲ ἔτι προσεπιλαμβάνουσιν. οὐ μὴν οὐδὲ ἐκεῖ τό γε πᾶν πλῆθος οὕτως, ἀλλ' ὅσον εἰς ἀπαρχήν· κομίζεται γὰρ πρὸς ἱερῶν τινῶν χρείαν ἄλφιτα νέα τῷ ἔκτω μηνὶ καὶ ταῦτα ἐκ τῶν ἄνω τόπων ὑπὲρ Μέμφιν.

Β Λέγεται δε καὶ ἐν Σικελία τῆς Μεσσηνίας ἐν

¹ μέλινοι Ald, Η,; έλυμοι Vo, Vin, cf. 8. 1. 1 n.

ENQUIRY INTO PLANTS, VIII. 11. 5-8

Wherefore, at the time when some of the vetches are gathered, the lower seeds have already fallen,

while the upper ones are still quite green.

After the flowering is over wheat and barley develop and mature in about forty days; one-seeded wheat and other such plants take about the same time. So too, they say, does the bean, which blooms and matures in a like number of days: but the others take fewer, and fewest of all the chick-pea, since, as some say, it takes only forty days from the time when it is sown to that when it is mature; and in any case it is clear that the plant as a whole develops very rapidly. Millet sesame Italian millet 1 and the summer crops in general, it is fairly well agreed, take the same number of days, that is, forty; though some say that they take less.

Of differences in development due to soil or climate.

² Again as to the development of the plant there are differences according to soil and climate. Some soils seem to produce the crop in fewer days; for instance, Egypt may be given as a specially conspicuous example; in that country barley is reaped in six months and wheat in seven: while in Hellas the barley ⁸ harvest is in the seventh month, or in most parts in the eighth, and wheat requires an even longer time. However even in Egypt the whole harvest is not gathered at such an early date, but only what is required for the first-fruits; for they gather new grain for the meal required in certain sacrifices in the sixth month, and that too in the regions high up the Nile, above Memphis.

It is said also that in the Messenian district in

² Plin. 18. 49. ³ κριθαί conj. Sch.; πυροί UMAld.

ταῖς καλουμέναις Μύλαις ταχειάν τινα γίνεσθαι τὴν τελείωσιν τῶν ὀψίων τὸν τῶν ὀσπρίων μὲν γὰρ σπορητὸν ἔξ μῆνας, τὸν δὲ τῷ ὑστάτῷ σπείραντα θερίζειν ἄμα τοῖς πρώτοις ἀγαθὴν δὲ διαφερόντως εἶναι τὴν χώραν, ὅστε τριακοντάχοα ποιεῖν, ἔχειν δὲ καὶ νομὰς θαυμαστὰς καὶ ὕλην. ἐν Μήλῷ δέ τι θαυμασιώτερον λέγουσιν ἐν γὰρ τριάκοντα ἡ τετταράκοντα ἡμέραις σπαρέντα θερίζουσι, δι' δ καὶ λέγειν αὐτοὺς ὅτι μέχρι τούτου δεῖ σπείρειν ἔως ἃν ἴδῃ τις δράγμα γίνεσθαι δὲ οὕτε ὅσπρια τοιαῦτα οὕτε πολλὰ παρ' αὐτοῖς. δεινὴν δέ τινα διαδοῦναι τὴν χώραν τροφήν καὶ γὰρ εἶναι σιτοφόρον μὲν καὶ ἐλαιοφόρον ἀγαθὴν ἀμπελοφόρον δὲ μετρίαν.

Υπερβάλλον δ' έτι τούτου καὶ πάντων θαυμασιώτερον τὸ περὶ Χαλκίαν τὴν νῆσον τὴν 'Ροδίων γινόμενον' ἐκεῖ γάρ φασιν εἶναί τινα τόπον πρώϊον οὕτω καὶ εὕφορον ὡς σπαρεισῶν κριθῶν ἄμα ταῖς ἄλλαις θερίσαντες ταύτας σπείρουσιν εἶτα πάλιν, εἶτα θερίζουσιν ἄμα τοῖς λοιποῖς μεγίστη μὲν οὖν, εἴπερ ἀληθής, αὕτη διαφορά. τὸ γὰρ εἰς ἐτέραν χώραν μετενεχθέντα διαφέρειν, ὅσπερ ἐκ Κιλικίας φασὶν εἰς Καππαδοκίαν καὶ ὅλως τὴν ἐπέκεινα τοῦ Ταύρου, ἦττον ἄτοπον

φανερά γὰρ ή τῶν τόπων διάστασις.

¹ τριακοντάχοα conj. Sch.; τριάκοντα χοὰs Ald. ² cf. C.P. 4. 11. 8.

ENQUIRY INTO PLANTS, VIII. 11. 8-9

Sicily at the place called Mylae the late sown crops mature rapidly; thus the sowing of pulses goes on for six months, but he that made the last sowing gathers his crop at the same time as the first: also that the soil is exceedingly good, so that it yields thirty-fold 1; and there are also wonderful pastures and forest-land. They tell of an even more wonderful thing in Melos 2; there they reap thirty or forty days after sowing; wherefore it is a saying of the islanders that "one should continue sowing till one sees a swathe." However it is said that pulses 3 in their country do not grow like this, nor are they abundant. Yet they say that the soil is wonderfully productive; for it is good both for corn and olives, and fairly good for vines.

However what occurs in Chalkia,⁴ an island belonging to the Rhodians, goes even beyond this and is more extraordinary than all the instances given; there they say that there is a place which is so early and so fertile that, when the barley is sown after reaping the crop with the other crops, they then sow again, and then reap the crop thus sown at the same time as the remaining crops; this then, if it be true, marks a difference greater than we find anywhere else. For it is less surprising that there should be a difference in crops transferred ⁵ to another region, as they say occurs when they are transferred from Cilicia to Cappadocia or in general beyond the Taurus; for these regions are obviously very dissimilar.

4 cf. Thuc. 8. 41 foll.

 $^{^3}$ δσπρια τοιαθτα I conj. ; δψιμα ταθτα UAld.; ξψιμα ταθτα M.G ; P omits ταθτα.

 $^{^{5}}$ μετενεχθέντα διαφέρειν conj. Sch. and W. from G; μετεγκόντας σπείρειν Ald.

10 Τὸ δὲ τὴν αὐτὴν διφορεῖν, ἐν ῷπερ γε ἄπαξ αἱ ἄλλαι, σύνορον οὖσαν καὶ μίαν θαυμασιώτατον αὕτη μὲν οὖν ἐν μεγίστη διαφορᾳ̂.

Τὰ δὲ κατὰ τὰς ἄλλας χώρας οὐ πολλὴν ἡ οὐδεμίαν ὡς εἰπεῖν τοῦ γε χρόνου λαμβάνει διάστασιν προτερεῖ γὰρ ταῖς ὥραις τὰ ᾿Αθήνησι τῶν περὶ Ἑλλήσπουτον ἡμέραις τριάκοντα μάλιστα ἡ οὐ πολλῷ πλείοσιν εἰ μὲν οὖν καὶ ὁ σπορητὸς πρότερον, μετάθεσις ἂν εἴη τῆς ὥρας εἰ δ᾽ ἄμα, δῆλον ὅτι πλείων ἂν ὁ χρόνος.

11 Ο υ μικράν δε ποιουσι διαφοράν ουδε οι τόποι, καίπερ ενιοι συνεγγυς όντες τὰ γὰρ εν Σαλαμινι προτερεί πολυ τῶν ἄλλων τῶν εν τῷ 'Αττικῷ καὶ ὅλως τὰ ἐπιθαλάττια καὶ εἰς ταῦτα καὶ εἰς τοὺς ἄλλους καρπούς, ὡς τὰ περὶ τὴν 'Ακτὴν καλουμένην τῆς Πελοποννήσου καὶ τὰ ἐν Φαλύκω τῆς Μεγαρίδος πλὴν ἐνταῦθά γε συμβάλλεται καὶ τὸ λεπτόγεων εἶναι καὶ ψαφαρὰν τὴν χώραν. καὶ τὰ μὲν περὶ τὴν γένεσιν καὶ τὴν τελείωσιν οὕτως ἔγει.

ΙΙΙ. Διαφέρει δὲ καθ' ὅλα τὰ γένη τὰ διηρημένα τῶν γενῶν, οἶον σῖτος χεδροπὰ τὰ θερινά,
καὶ καθ' ἔκαστον γένος τὰ ὁμογενῆ. τὰ μὲν γὰρ
σιτώδη τὸ φύλλον ἔχει καλάμου, τῶν δὲ χεδρο-

 $^{^1}$ i.e. and so in part account for the difference. Ein $\tau \hat{\eta}$ s whas conj. Sch.; $\hat{\eta}$ $\tau \hat{\eta}$ s xwas MP; $\epsilon \hat{\eta}$ n $\tau \hat{\eta}$ s xwas Ald. H.

² i.e. we cannot say how far the difference is due to climate without knowing whether the seed-time at either place is the same.

ENQUIRY INTO PLANTS, VIII. II. 10-III. I

But that one particular land should produce two crops in the time that other lands to which it is close take to produce one, is very remarkable; wherefore

Chalkia exhibits the greatest difference.

The crops grown in other regions show not much, if any, dissimilarity as to time; those grown at Athens are only about thirty days or not much more before those of the Hellespont region. Now, if the sowing should turn out to be also earlier, that would shift the season back 1; if it is at the same time, it is plain that the difference of time

would be greater.2

Again the particular district makes a considerable difference, even as between places which are not far apart; thus the crops of Salamis are far earlier than those of the rest of Attica, and so in general are those of places by the sea; and this applies to other fruits as well as these: for instance, those of the place called Akte in the Peloponnese and of Phalykos ³ in the Megarid are early; but here something is contributed by the fact that the soil is light and crumbling. Such are the facts in regard to growth and development.

Of differences between the parts of cereals, pulses, and summer crops respectively.

III. There are also differences between 4 the whole classes which we have mentioned, namely cereals leguminous plants 5 and summer crops, as well as between the several members 6 of the same class. Cereals have the leaf of a reed, while of

⁴ καθ' conj. Sch.; καὶ Ald.H. ⁵ cf. 8. 1. 1.

6 δμογενή conj. Sch.; δμοιογενή Ald.

 $^{^3}$ έν Φαλύκφ I conj.: cf. 2. 8. 1; έν Φαλήκφ conj. W.; έφαλύκφ U; έκ φαλήκφ M; έκ φαλήκον Ald.

πῶν τὰ μὲν περιφερές, οἶον ὁ κύαμος καὶ σχεδὸν τὰ πλεῖστα, τὰ δὲ προμηκέστερον, οἶον ὁ πισὸς καὶ ὁ λάθυρος καὶ ὁ ὁ ἄχρος καὶ τὰ τοιαῦτα. καὶ τὰ μὲν ἰνώδη τὰ δ' ἄφλεβα καὶ ἄϊνα. τὸ δὲ σήσαμον καὶ τὸ ἐρύσιμον ἰδιώτερα παρὰ ταῦτα.

2 Πάλιν ὁ καυλὸς τῶν μὲν γονατώδης καὶ κοῖλος, δι' ὁ καὶ καλεῖται κάλαμος ὁ δὲ τοῦ κυάμου κοῖλος, τῶν δ' ἄλλων χεδροπῶν ξυλωδέστερος, ξυλωδέστατος δὲ ὁ ἐρέβινθος τῶν δὲ θερινῶν κέγχρου μὲν καὶ μελίνου καλαμώδης, σησάμου δὲ καὶ ἐρυσίμου ναρθηκώδης μᾶλλον. καὶ τὰ μέν ἐστιν ὀρθόκαυλα, καθάπερ πυρὸς καὶ κριθὴ καὶ ὅλως τὰ σιτώδη καὶ θερινά, τὰ δὲ πλαγιόκαυλα μᾶλλον, οἶον ἐρέβινθος ὄροβος φακός, τὰ δ' ἐπιγειόκαυλα, καθάπερ ἄχρος πισὸς λάθυρος ὁ δὲ δόλιχος, ἐὰν παρακαταπήξη τις ξύλα μακρά, ἀναβαίνει καὶ γίνεται κάρπιμος, εἰ δὲ μή, φαῦλος καὶ ἐρυσιβώδης μόνος δ' ἡ μάλιστα τῶν χεδροπῶν ὀρθόκαυλος ὁ κύαμος.

Έχει δὲ καὶ τὰ ἄνθη διαφορὰν καὶ τῆ φύσει καὶ τῆ θέσει, περὶ ὧν σχεδὸν ἐν τοῖς καθ' ὅλου διείλομεν, ὅτι τὰ μὲν χνοώδη, καθάπερ σίτου καὶ παντὸς τοῦ σταχυώδους τὰ δὲ φυλλώδη, καθάπερ τῶν χεδροπῶν, καὶ τῶν πλείστων κολοβά·

Plin. 18. 58.
 i.e. 'summer crops'; cf. 8. 1. 1.
 Sc. but not jointed. W. suggests that the original text may have been τῶν δὲ διόλου κοῖλος οδον ὁ τοῦ κυάμου.

⁴ μελίνου Ald. H.; ἐλύμου V; ἐλύμου Vin. cf. 8. 1. 1; 8. 1. 6. ⁵ ħ add. St.; om. Ald. H.G. ⁶ 1. 13. 1.(?)

⁷ χνοώδη. No rendering seems quite satisfactory: the 160

ENQUIRY INTO PLANTS, VIII. III. 1-3

leguminous plants some have a round leaf, as beans and most others, some a more oblong leaf, as pea lathyros okhros and the like. ¹Some again have fibrous leaves, others leaves without veins and fibres. Again sesame ² and erysimon ² have leaves quite distinct from these.

Again the stem of cereals is jointed and hollow; wherefore it is called the 'reed,' while that of the bean is hollow,3 and that of the other leguminous plants is more woody, that of chick-pea woodiest of all; of the summer crops that of millet and Italian millet 4 is reed-like, that of sesame and erysimon is more like the stem of ferula. again have erect stems, as wheat barley and in general the cereals and summer crops; some have rather a crooked stem, as chick-pea vetch lentil; some a creeping stem as okhros pea lathyros; while calavance, if long stakes are set by it, climbs them and becomes fruitful, whereas otherwise the plant is unhealthy and liable to rust; the bean, most of all leguminous plants, if not 5 alone among them, has an erect stem.

The flowers also shew differences in character and in position (of which matters we have to some extent treated in our general account) 6; thus some are 'downy,' 7 as those of corn 8 and of any plant that has an 'ear'; others are 'leafy,' 9 as those of leguminous plants, and in most cases they are irregular 10 flowers; for most of these have

meaning is that such flowers may be classed with those distinguished by this term in 1. 13. 1, as not being petaloid.

16 cf. 6. 5. 3. i.e. they depart from radial symmetry.

161

VOL. II.

M

^{*} σίτου και παντός τοῦ σταχυώδους conj. Sch. from G, ut omnium fere gerentium spicam; που και παντός τοῦ χυλώδους UMAld.

* Sc. petaloid.

τὰ γὰρ πολλὰ κολοβανθη χνοῶδες δὲ καὶ τὸ τοῦ κέγχρου καὶ μελίνου τοῦ δὲ σησάμου καὶ τοῦ ἐρυσίμου φυλλῶδες. καὶ ὅτι δὴ τὰ μὲν ἔχει περὶ αὐτὸν τὸν καρπόν, οἶον τὰ σιτώδη καὶ κεγχρώδη περὶ τὸν στάχυν τὰ δὲ χεδροπὰ ἐξ αὐτοῦ πως τοῦ ἄνθους ἡ ἀπό γε τῆς αὐτῆς ἀρχῆς γίνεται. καὶ τὴν ἄνθησιν, ὅτι τὰ μὲν ἀθρόαν τὰ δὲ κατὰ μέρος ποιεῖται καὶ τἆλλα δὲ τὰ παραπλήσια τούτοις.

'Όμοίως δὲ καὶ τὰ κατὰ τοὺς καρπούς, ὅτι τὰ μὲν ἔχει στάχυν, τὰ δὲ χεδροπὰ λοβόν, τὰ δὲ κεγχρώδη φόβην ἡ δὲ καλαμώδης ἀπόχυσις φόβη. τὸ δ' ὅλον ἐναγγειόσπερμα, τὰ δὲ ἐνυμενόσπερμα, τὰ δὲ ἀνυμενόσπερμα καὶ ἔτι τὰ μὲν ἀκρόκαρπα, τὰ δὲ πλαγιόκαρπα, καὶ ὅσα δὴ ἄλλα ταύτης ἔχεται τῆς θεωρίας.

"Όλως δὲ πολυκαρπότερα καὶ πολυχούστερα τὰ χεδροπά, τούτων δ' ἔτι μᾶλλον τὰ θερινὰ κέγχρος καὶ σήσαμον, αὐτῶν δὲ τῶν χεδροπῶν 5 μάλιστα φακός. ἀπλῶς δὲ τὰ μικροσπερμότερα μᾶλλον ὡς εἰπεῖν, ὥσπερ καὶ τῶν λαχανωδῶν κύμινον ἀπάντων ὄντων πολυσπέρμων. ἰσχυρότερα δὲ πρὸς μὲν τὸν χειμῶνα καὶ ὅλως τὰ τοῦ ἀέρος τὰ σιτηρά, πρὸς δὲ τὴν τροφὴν τὰ

5 τὸ δ' δλον: ? τὰ δ' οἷον.

¹ μελίνου Ald. H.; ἐλύμου Vo. Vin. cf. 8. 3. 2 and reff.

 ² cf. 8. 3. 3 n.
 3 Plin. 18. 53.
 4 ἀπόχυσις conj. Sch. from G; ἀπόφυσις P₂Ald. cf. 4. 4. 10, ἀποχεῖται; 8. 10. 4; C.P. 3. 21. 5.

ENQUIRY INTO PLANTS, VIII. III. 3-5

such flowers. Those of millet and Italian millet ¹ are also 'downy,' ² those of sesame and *erysimon* 'leafy.' Another difference is that in some cases the flowers are round the fruit; thus those of corn and millet are round the ear; while in leguminous plants the fruit comes as it were from the flower itself, or at least from the same starting-point. Another difference is that some produce all their flowers at once, others in succession. And there are other differences akin to these.

In like manner there are differences in the fruits; some have an 'ear,' leguminous plants a pod, and millet-like plants a 'plume' 3—which is the name given to an inflorescence 4 such as reeds have. Again, generally speaking, 5 some have their seeds in a vessel, 6 some in pods, 7 some naked; and further some bear their fruit at the top, some at the sides; and there are other differences which bear on this enquiry.

In general the leguminous plants produce more fruit and are more prolific, and the summer crops millet and sesame are even more so than these, while among the leguminous plants themselves lentil is the most prolific. ⁸ Generally speaking, those that have small seeds are more prolific, as cummin among pot-herbs, which are all prolific of seed. The seeds of cereals are more robust as to standing winter and conditions of climate generally, while those of leguminous plants are stronger as to providing food. ⁹ However it may be that in this respect

 $^{^6}$ μèν ἐναγγειόσπερμα conj. Sch.; μèν ἐγγειόσπερμα $P_2Ald.$ cf. C.P. 4. 7. 5.

⁷ cf. 1. 11. 2. ⁸ cf. C.P. 4. 15. 2.

[•] i.e. what has just been said perhaps applies only to human food. Sense fixed by 8. 9. 3 ad fin.: cf. Plin. 18. 50.

χεδροπά. τάχα δὲ τοῦτό γε ἡμῖν τοῖς ἄλλοις ἀνάπαλιν.

IV. Τὰ μὲν οὖν ὅλα γένη τοιαύτας ἔχει διαφοράς τὰ δὲ ὁμογενη δηλον ὅτι κατὰ τὴν τῶν μερῶν ἀνωμαλίαν, οἶον τῶν σιτωδῶν πυρὸς κριθης στενοφυλλότερον καὶ λειοκαυλότερον καὶ πυκνότερον καὶ γλισχρότερον ἔχει τὸν καυλὸν καὶ δυσθλαστότερον ἄμα δὲ καὶ ὁ μὲν ἐν χιτῶσι πολλοῖς ἡ δὲ γυμνόν μάλιστα γὰρ δὴ γυμνοσπέρματον ἡ κριθή. πολύλοπον δὲ καὶ ἡ τίφη καὶ ἡ ὀλύρα καὶ πάντα τὰ τοιαῦτα καὶ μάλιστα πάντων ὡς εἰπεῖν ὁ βρόμος. ἔστι δὲ καὶ ὑψηλότερος ὁ κάλαμος τοῦ πυροῦ ἡ τῆς κριθης, καὶ τὸν στάχυν ἀπηρτημένον ἔχει τοῦ φύλλου μᾶλλον ὁ πυρός.

Τδιον δὲ καὶ τὸ ἄχυρον τοῦ κριθίνου τὸ πύρινον ἐγχυλότερον γὰρ καὶ μαλακώτερον. διαφέρει δὲ
 ἡ κριθἡ καὶ τούτῳ τῶν πυρῶν ἡ μὲν γὰρ στοιχειώδης, ὁ δὲ πυρὸς ἄστοιχος καὶ πανταχόθεν

δμαλής τις.

Τῷ μὲν οὖν ὅλῳ γένει πρὸς γένος τοιαῦταί τινές εἰσι διαφοραί. καθ' ἐκάτερον δὲ τούτων πάλιν, οἶον πυρῶν καὶ κριθῶν, πολλὰ γένη καὶ τοῖς καρποῖς αὐτοῖς διαφέροντα καὶ τοῖς στάχυσι καὶ ταῖς ἄλλαις μορφαῖς καὶ ἔτι ταῖς δυνάμεσι

3 δμογενη conj. Sch.; δμοιογενη UMAld. cf. 8. 3. 1.

⁵ Plin. 18. 61. πολύλοπον conj. Salm.; πολύλοβον Ald.

¹ cf. 7. 4. 9.

² Åfter διαφοράς UM add τὰ δμοιογενη, Ald. τὰ μη δμοιογενη; om. Sch. and W. after G.

⁴ δυσθλαστότερον conj. Scal. from G, ruptu difficiliorem; δυσαλθατώτερον UMAld.

ENQUIRY INTO PLANTS, VIII. III. 5-IV. 2

the other animals are affected in the opposite 1 way to men.

Of the differences between cereals.

IV. There are then these differences ² between the various classes; and as between plants of the same class ³ there are plainly differences due to the unlikeness in the various parts. Thus among cereals wheat as compared with barley has a narrower leaf, and a smoother stem of closer texture tougher and less brittle.⁴ Again the seed of wheat has several coats,⁵ that of barley is naked, that plant having its seeds specially naked. Also one-seeded wheat rice-wheat and all such plants have their seed in several coats, and above all, it may be said, is this true of oats.⁶ Also the 'reed' of wheat is taller than ⁷ that of barley, and wheat has its ear less distant from the 'leaf.'

Further the husk of wheat is distinct ⁸ from that of barley, being less dry and softer. Barley also differs from wheat in this respect; it has grains in a regular row, ⁹ whereas those of wheat are not in a row, but the ear is as it were quite simple in form. ¹⁰

Such then are the differences as between one whole kind and another. But in each of these kinds again, for instance in barley and wheat, there are many sub-divisions differing both in the actual fruits, in the ear, and in the other characteristic

' η conj. Sen. from Flin. l.c. and V; και Ald. H.

8 τδιον Ald.; ηδιον Vin. H. from G: so Sch. and W. cf.
Col. 6. 3. 3.

165

βρόμος conj. Scal. from Plin. l.c. and G; κρόμος PM;
 κρόκος Ald.; βρῶμος Vin.
 † conj. Sch. from Plin. l.c. and G; καὶ Ald. H.

⁹ στοιχειώδης. ? στοιχώδης: v. LS. ¹⁰ όμαλης conj. Sch.; όμαλη UMAld.

καὶ τοῖς πάθεσι. τῶν μὲν κριθῶν αἰ μέν εἰσι δίστοιγοι αι δὲ τρίστοιγοι αι δὲ τετράστοιγοι καὶ πεντάστοιγοι πλείστον δ' έξάστοιγον, καὶ γαρ τοιουτό τι γένος έστί. πυκνότεραι δὲ ἀεὶ κατὰ τὴν θέσιν ὡς ἐπὶ πᾶν αἱ πολυστοιχότεραι. διαφορά δὲ μεγάλη καὶ τὸ παραβλαστητικήν είναι, καθάπερ είπομεν την Ίνδικήν. καὶ οί στάχυες δὲ τῶν μὲν μεγάλοι καὶ μανότεροι ταῖς κριθαίς των δὲ ἐλάττους καὶ πυκνότεροι, καὶ ἀπέγοντες δὲ τοῦ Φύλλου τῶν μὲν πολύ τῶν δὲ μικρόν, ώσπερ των 'Αγιλλείων καλουμένων. καὶ αὐτῶν δὲ τῶν κριθῶν αἱ μὲν στρογγυλότεραι καὶ έλάττους αί δὲ προμηκέστεραι καὶ μείζους καὶ μανότεραι κατά τὸν στάχυν. ἔτι δὲ αί μὲν λευκαί, αί δε μέλαιναι καί επιπορφυρίζουσαι, αίπερ καὶ πολυάλφιτοι δοκοῦσιν είναι καὶ πρὸς τοὺς γειμώνας δὲ καὶ τὰ πνεύματα καὶ ὅλως τὸν ἀέρα τῶν λευκῶν ἰσχυρότεραι.

Πολλά δὲ γένη καὶ τῶν πυρῶν ἐστιν εὐθὺς ἀπὸ τῶν χωρῶν ἔχοντα τὰς ἐπωνυμίας, οἶον Λιβυκοὶ Ποντικοὶ Θρᾶκες ᾿Ασσύριοι Αἰγύπτιοι Σικελοί. διαφορὰς δὲ καὶ ταῖς χροιαῖς καὶ τοῖς μεγέθεσι καὶ τοῖς εἴδεσι καὶ ταῖς ἰδιότησιν ἔχουσι καὶ ἐν ταῖς δυνάμεσι ταῖς τε ἄλλαις καὶ μάλιστα ταῖς πρὸς τὴν σίτησιν. τινὲς καὶ ἀπ᾽

¹ Explained below, 8. 4. 4; cf. 8. 4. 3. ² πάθεσι: cf. 1. 1. 1 n. ³ Plin. 18. 78.

^{*} πλείστον δ' έξάστοιχον, και γάρ τοιούτο conj. W.; πλείστον έξάστοιχον τοιούτον UM; πλείσται και έξάστιχοι και γάρ τοιούτον Ald. H,

ENQUIRY INTO PLANTS, VIII. iv. 2-3

features: and again in capacities 1 and properties.2 3 Of barley different sorts have respectively two, three. four, and five rows of seeds; the largest number 4 known is six, for there is a kind which bears that number. And those which have more rows have generally the grains set closer together. Another great difference is that of having side-shoots, as we said of the Indian kind.5 Again in barley 6 the ears are in some kinds large and of looser make, in some smaller and set closer; in some kinds the ear is some way from the 'leaf,' in some it is nearer to it, as in the kind called 'Achillean.' 7 Again of the grains themselves some are rounder and smaller, some more oblong and larger and set at wider intervals on the ear. Moreover some are white, some black or reddish, and the latter are thought to produce much meal and to be more robust than the white as to bearing winter wind or conditions of climate generally.

There are 8 also many kinds of wheat which take their names simply from the places where they grow, 9 as Libyan Pontic 10 Thracian Assyrian Egyptian Sicilian. They show differences 11 in colour size form and individual character, and also 12 as regards their capacities 18 in general and especially their value as food. Some again get

⁵ Referred to 4. 4. 9, but without mention of this feature.

ται̂s κριθαι̂s conj. W.; τη̂s κριθη̂s Ald.
 cf. C.P. 3. 21. 3; 3. 22. 2.

⁸ έστιν εὐθὺς conj. W.; εὐθύς έστιν Ald.

⁹ cf. C.P. 3. 21. 2; Plin. 18. 2. ¹⁰ Ποντικοί conj. Sch.; πόντιοι Ald.

¹¹ διαφοράς conj. W.; διαφέροντες Ald. H.

¹² kal conj. W.; & Ald.

¹³ Explained below, 8. 4. 4: pace of growth.

ἄλλων τὰς προσηγορίας, οἶον καγχρυδίας στλεγγὺς ᾿Αλεξάνδρειος· ὧν ἀπάντων ἐν τοῖς εἰρημένοις
τὰς διαφορὰς ληπτέον. οὐχ ἤκιστα δ᾽ οἰκεῖαι
εἴ τις λαμβάνοι τὰς τοιαύτας· οἶόν εἰσιν οἱ μὲν
πρώϊοι οἱ δὲ ὄψιοι, καὶ εὐαυξεῖς καὶ πολύχοι οἱ
δὲ <μικροὶ> καὶ ὀλιγόχοι, καὶ μεγαλοστάχυες οἱ δὲ
μικροστάχυες. καὶ οἱ μὲν ἐν κάλυκι πολὺν χρόνον
οἱ δ᾽ ὀλίγον ἔνοντες, ὥσπερ ὁ Λιβυκός. καὶ κάλαμον οἱ μὲν λεπτὸν οἱ δὲ παχύν· καὶ τοῦτο ὁ Λιβυκὸς ἔχει, παχὺν δὲ καὶ ὁ καγχρυδίας. ἔτι δὲ
χιτώνας οἱ μὲν ὀλίγους οἱ δὲ πολλούς, ὥσπερ ὁ
Φράκιος. καὶ ὁ μὲν μονοκάλαμος ὁ δὲ πολυκάλαμος, καὶ μᾶλλον δὲ καὶ ἦττον.

Όμοίως δὲ καὶ εἴ τι παραπλήσιον τούτοις ἡ τοῖς πρότερον εἰρημένοις κατὰ τὰς δυνάμεις. αἱ γὰρ τοιαῦται φυσικώταται δόξαιεν ὰν εἶναι τῶν διαφορῶν. ἐν αἶς καὶ τὸ τῶν τριμήνων καὶ τὸ τῶν διμήνων καὶ εἴ τι γένος ἐν ἐλάττοσιν ἡμέραις τελειοῦται, καθάπερ φασὶν εἶναι περὶ τὴν Αἰνείαν, οῖ τετταράκοντα ἡμέραις ἀπὸ τῆς σπορᾶς ἀδρύνονται καὶ τέλος ἴσχουσιν· εἶναι δ' ἰσχυρὸν τοῦτον καὶ βαρὺν οὐχ ὥσπερ τὸν τρίμηνον κοῦφον, δι' ὁ καὶ τοῖς οἰκέταις παρέχειν, καὶ γὰρ οὐδὲ πίτυρον ἔχειν πολύ. σπανιώτατος μὲν οὖν καὶ τάχιστος εἰς τελείωσιν

² cf. Geop. 3. 3. 11. ³ i.e. colour, size, etc. ⁴ μικροὶ add. W. to correspond to εὐαυξεῖς (conj. Sch.; εὐαξεῖς Ald.).

168

¹ στλεγγύs. Sir W. Thiselton-Dyer conjectures σίλιγνις: cf. Plin. 18. 184, LS. σιλίγνιον.

ENQUIRY INTO PLANTS, VIII. IV. 3-4

their distinctive names for other reasons, as kankhrydias stlengys 1 'Alexandrian'2; all of which must be distinguished by the above-mentioned 8 characters. Again, if one takes such differences as the following. they are quite characteristic—thus some are early, some late, some are vigorous and prolific, some are small and produce little, some have a large, some a small ear. The ears of some remain 5 a long time in the sheath,6 of some it remains but a short time, as that of the Libvan kind, 7 Again some have a slender, some a stout haulm; the Libyan kind has this characteristic also, and that of kankhrudias is also stout. Again the grain of some has few coats,8 of some many, for instance the Thracian.9 Some kinds have a single 'reed,' some more than one, and in the latter class the number varies.

10 So too must we distinguish any differences like these or those mentioned above which are found in the several capacities; for these would seem to be the most essential differences. In this connexion we may distinguish kinds which mature in three or in two months, and those, if there be such, which take a less number of days; for instance, they say that in the region of Aineia there is a kind which ripens and attains perfection within forty days from the time of sowing; they say too that this grain is strong and heavy, not light like that which takes three months; wherefore they give it even to the servants, for it also does not contain much bran. Now this kind is the rarest and the quickest to

⁵ ξνοντες conj. W.; ξχοντες Ald. ⁶ cf. 8. 2. 4. ⁷ Plin. 18. 69. 8 i.e. glumes. ° cf. C.P. 4. 12. 5; Plin. l.c. 10 Plin. 18. 70.

ούτος, είσι δε και δίμηνοί τινες οίπερ και εκ Σικελίας εκομίσθησαν είς 'Αχαίαν' όλιγοχόοι δε καὶ όλιγογόνοι καὶ κοῦφοι κατὰ τὴν προσφορὰν καὶ ήδεις. καὶ ἄλλοι δέ τινες οι περὶ Ευβοιαν είσι και μάλιστα έν τη Καρυστία. τρίμηνοι δέ πολλοί και πανταγού κούφοι ούτοι και όλιγογόοι καὶ μονοκάλαμοι κατὰ τὴν ἔκφυσιν καὶ τὸ ὅλον 5 ἀσθενείς. κουφότατος μέν οὖν ὡς ἀπλῶς εἰπεῖν πυρὸς ὁ Ποντικός βαρύτερος δὲ τῶν εἰς τὴν Έλλάδα παραγινομένων ο Σικελός τούτου δ' έτι βαρύτερος ο Βοιωτός σημείον δε λέγουσιν ότι οί μεν άθληταὶ ἐν τῆ Βοιωτία τρί ἡμιχοίνικα μόλις ἀναλίσκουσιν, ᾿Αθήναζε δὲ ὅταν ἔλθωσι πένθ' ήμιγοίνικα ραδίως. κοῦφος δὲ καὶ ὁ ἐν τη Λακωνική. τούτων μέν οὖν ἔν τε ταίς χώραις καὶ τῷ ἀέρι τὸ αἴτιον ἐπεὶ καὶ περὶ τὴν Ασίαν ου πόρρω Βάκτρων έν μέν τινι τόπω ούτως άδρον είναι φασι τον σίτον ώστε πυρήνος έλαίας μέγεθος λαμβάνειν, έν δε τοις Πισσάτοις καλουμένοις ούτως ισχυρον ώστ' εί τις πλείον προσενέγκοιτο διαρρήγνυσθαι, καὶ τῶν Μακεδόνων καὶ πολλούς τοῦτο παθεῖν. ἄτοπον δὲ καὶ ἀνομολογούμενον πρὸς τὴν τῶν τριμήνων κουφότητα τὸ περὶ τοὺς Ποντικοὺς συμβαίνον. είσι γάρ οί μεν σκληροί ήρινοι οί δε μαλακοί χειμερινοί πολύ γάρ διαφέρει τη κουφότητι ό 6 μαλακός. [όμοίως δε καὶ δύο ἀρότους ώς ἔοικε

 $^{^1}$ τρί' ἡμιχοίνικα conj. Sch.; τριημισχοίνικα ; τριημιχοίνικα $P_2Ald.\,H.$

¹⁷⁰

ENQUIRY INTO PLANTS, VIII. IV. 4-6

mature. But there is also a kind which takes two months: this was brought to Achaia from Sicily: it is not however prolific nor fertile, though as food it is light and sweet. There is another such kind which grows in Euboea and especially in the region of Karystos. There are several kinds that take three months, and these, wherever they are found, are light and not prolific; their growth consists of a single 'reed,' and in general they are not robust. Lightest of all we may say is the Pontic wheat: the Sicilian is heavier than most of those imported into Hellas, but heavier still than this is the Boeotian; in proof of which it is said that the athletes in Boeotia consume scarcely three pints,1 while, when they come to Athens, they easily manage five.2 The Laconian kind is also light. The reason for these differences is to be found in the respective soils and in the climate; 3 for in Asia not far from Bactra they say that in a certain place the corn is so vigorous that the grains grow as large as an olive-stone, while in the country called that of the Pissatoi it is so strong that, if a man eats too much of it, he bursts, which was actually the fate of many of the Macedonians.4 There is one curious thing about the corn of Pontus, which is an exception 5 to the rule as to the lightness of crops raised in three months; for there the hard crops are those of the spring, the soft ones those of the winter; for soft kinds are exceedingly light. Two sowings, as it appears, are made of all corn

i.e. in Alexander's army.

 $^{^2}$ πενθ' ήμιχοίνικα conj. Sch.; πενθημισχοίνικα M; πενθημιχοίνικα $P_2Ald.H.$ 3 Plin. 18. 70.

⁵ ἀνομολογούμενον: cf. C.P. 4. 8. 2; Plat. Gorg. 495 A.

παυτὸς τοῦ σίτου ποιοῦνται, τὸν μὲν χειμερινὸν τὸν δὲ ἦρινόν, ἐν ὧ καὶ τὰ ὅσπρια καταβάλ-

λουσιν].

Εἰσὶ δὲ καὶ οἱ μὲν καθαροὶ αἰρῶν, ὅσπερ ὁ Ποντικὸς καὶ ὁ Αἰγύπτιος· καθαρὸς δὲ ἐπιει-κῶς καὶ ὁ Σικελὸς καὶ μάλιστα ὁ ᾿Ακραγαντίνος

οὐκ αἰρώδης.

Ο δε Σικελος ίδιον έχει το μελάμπυρον καλούμενον, ο έστιν άβλαβες και ούχ ώσπερ ή αίρα βαρύ και κεφαλαλγές. άλλα τα μεν τοιαῦτα, καθάπερ ελέχθη, ταις χώραις άναθετέον και οσον

έπιβάλλει τοις γένεσιν.

V. Έν δὲ τοῖς ὀσπρίοις οὐχ ὁμοίως ἐστὶ λαβεῖν τὰς τοιαύτας διαφοράς, εἴτ' οὖν διὰ τὸ μὴ ἐξετάζειν ὁμοίως εἴτε καὶ διὰ τὸ μονοειδέστερα τυγχάνειν· ἔξω γὰρ ἐρεβίνθου καὶ φακοῦ καὶ ἐπ' ὀλίγου κυάμου καὶ ὀρόβου, καθ' ὅσον ἡ τῶν χρωμάτων καὶ τῶν χυλῶν διαφορά, τῶν γ' ἄλλων οὐ ποιοῦσιν ἰδέας. οἱ δὲ ἐρέβινθοι καὶ τοῖς μεγέθεσι καὶ τοῖς χυλοῖς καὶ τοῖς χρώμασι καὶ ταῖς μορφαῖς διαφέρουσιν, οἶον κριοὶ ὀροβιαῖοι οἱ ἀνὰ μέσον. ἐπὶ πᾶσι δὲ τὰ λευκὰ γλυκύτερα· καὶ γὰρ ὁ ὄροβος καὶ φακὸς καὶ ἐρέβινθος καὶ κύαμος καὶ σήσαμον ἔστι γὰρ καὶ σήσαμον λευκόν.

2 'Αλλά μάλλον ἐν τοις τοιοισδε ποιειν ἐστι τὰς διαφορὰς οἰον, ἐπεὶ πάντα ταῦτ' ἔλλοβα, τὰ μὲν

¹ δμοίως . . . καταβάλλουσιν bracketed by Sch. as a gloss.

² But cf. 8. 8. 3. ³ cf. Diosc. 4. 116.

i.e. when it gets into the bread. fr. Plin. 18. 156; Diosc. 2. 100.

⁶ δσον επιβάλλει: cf. Arist. Pol. 1. 13. 8.

ENQUIRY INTO PLANTS, VIII. IV. 6-V. 2

alike, one in winter and one in spring, at which time

they also plant the seed of the pulses.1

Some kinds are free from darnel, as the Pontic and the Egyptian; the Sicilian is also fairly free from it, and that of Akragas is especially immune from darnel.

² Peculiar however to the Sicilian is the plant called *melampyron*, ³ which is harmless ⁴ and not, like the darnel, injurious and productive of headache. ⁵ However such peculiarities, as was said, must be ascribed to the soil, and to a certain extent ⁶ to the different characters of different kinds.

Of the differences between pulses.

V. In pulses we cannot find such differences to the same extent, whether for the want of equally careful enquiry or because there is actually less diversity in these plants. ⁷ For, apart from chickpea lentil and to a certain extent bean and vetch (in so far as in these we find differences of colour and taste), among the rest ⁸ no distinct forms are recognised. Chick-peas however differ in size colour taste and shape; thus there are the varieties called 'rams,' 'vetch-like' chick-peas and the intermediate forms.⁹ In all pulses the white are the sweeter, and this applies to vetch lentil chick-pea bean and sesame, of which also there is a white form.

. ¹⁰ However it is more possible to recognise the differences in such points as these:—all these plants have pods, ¹¹ but whereas the pods in some kinds have

Plin. 18. 124. ⁸ γ' conj. Sch.; τ' Ald. H.

10 Plin 18, 125.

of after δριβιαΐοι add. Dalec. For and μέσον cf. 3. 18, 2.

¹¹ έλλοβα conj. Scal. from G ; έλλέβυρα Ald. Η.; έλλόβορα U ; έπει πάντα ταῦτ' conj. W.; ἐπὶ πάντα τὰ Ald. Η.

αὐτῶν ἀδιάφρακτα καὶ ὥσπερ συμψαύοντα τυγχάνει, καθάπερ ὅροβος πισὸς καὶ σχεδὸν τὰ πλεῖστα, τὰ δὲ διαπεφραγμένα, καθάπερ θέρμος, ἔτι δὲ μᾶλλον καὶ ἰδίως τὸ σήσαμον. καὶ τὰ μὲν μακρόλοβα τὰ δὲ καὶ στρογγυλόλοβα, καθάπερ ὁ ἐρέβινθος. ἀνὰ λόγον δὶ ἀκολουθεῖ καὶ τὰ πλήθη τῶν σπερμάτων· ἐλάττω γὰρ ἐν τοῖς μικροῖς, ὥσπερ ἔν τε τῷ τῶν ἐρεβίνθων καὶ ἐν τῷ τῶν φακῶν.

Καὶ παραπλήσιαι δὲ ἴσως αι τοιαῦται καὶ ᾶς ἐπὶ τῶν σιτηρῶν ἐλέγομεν περὶ τῶν σταχύων καὶ αὐτῶν τῶν καρπῶν· ἐπεὶ καὶ οὶ καλούμενοι λοβοὶ σχεδὸν ἀκόλουθοι τοῖς σπέρμασίν εἰσιν, οἱ μὲν ἐπιπλατεῖς, ὥσπερ οἱ τοῦ φακοῦ καὶ τῆς ἀφάκης, οἱ δὲ κυλινδρώδεις μᾶλλον, ὡς οἱ τοῦ ὀρόβου καὶ τοῦ πισοῦ· τὰ γὰρ σπέρματα ἐκατέρων τοιαῦτα τοῖς σχήμασι· ἀλλὰ τὰς μὲν τοιαύτας διαφορὰς πολλὰς ἄν τις εὕροι καθ' ἔκαστον, ὧν αἱ μὲν

κοιναί πασιν αί δὲ ίδιαι κατά γένος.

"Ότι δὲ πάντα προσπέφυκε τοῖς λοβοῖς καὶ ἔχει καθάπερ ἀρχήν τινα, τὰ μὲν προέχουσαν, ὅσπερ ὁ κύαμος καὶ ὁ ἐρέβινθος, τὰ δὲ καὶ ἔγκοιλον, ὅσπερ θέρμος καὶ ἄλλ' ἄττα, τὰ δὲ οὕτω μὲν οὐ φανερὰν ἐλάττω δὲ καὶ ὅσπερ ἀποσημαίνουσαν μόνον, δῆλον μὲν ἀπὸ τῆς ὅψεως ἐξ ἡς καὶ ὅταν σπαρῆ βλαστάνει καὶ ῥιζοῦται, καθάπερ ἐλέχθη, κατ' ἀρχὰς δὲ καὶ αὐτὰ τρέφεται προσηρτημένα τῷ λοβῷ, μέχρι οῦ ἄν

³ διαπεφραγμένα conj. Sch.; λεία πεφραγμένα Ald. Η.

4 cf. 3. 18. 13.

 ¹ адафракта conj. Scal. from G, non intersepta; дафракта Ald. Ĥ.
 ² cf. 1. 11. 5.

ENQUIRY INTO PLANTS, VIII. v. 2-4

no divisions, but the seeds as it were touch one another, as in vetch pea and most kinds, in some there are divisions, as in lupin and still more in sesame, in which the divisions are of a peculiar kind. Again some have long, some round pods, as chick-pea. And the number of seeds follows in proportion, since they are fewer in the small pods, as in those of chick-pea and lentil.

Possibly these differences correspond to those which we mentioned in the case of cereals as to the ears and the actual fruits; for what are called 'pods' also 5 fairly correspond to the shape of the seeds, some being flat, as those of lentil and tare, some more or less cylindrical, as those of vetch and pea 6: for in the case of either pair of plants the seeds correspond in shape. However one might discover and distinguish many such differences, of which

some are common to a whole kind,7 others special to

particular varieties.

In all cases the seeds are attached to the pods and have a sort of starting-point, which in some cases projects, as in bean and chick-pea, in some is hollow, as in lupin and some others, and in some is not thus conspicuous but smaller and, as it were, only indicated; this is plain from observation; it is from this point that the seeds germinate and take root when they are sown, as was said 8: but to start with they are themselves nourished by being so attached to the pod until they are matured. This

5 i.s. as does the form of the ear in cereals.

8 8, 2, 1,

⁶ καὶ τοῦ πισοῦ· τὰ γὰρ conj. Scal. from Plin. l.c. and G; τοῦ πισοῦ γὰρ τὰ UMAld.

⁷ i.e. which either differentiate (e.g.) pea from lentil, or one variety of pea from another. cf. 8, 4, 2 n.

τελειωθ $\hat{\eta}$ · φανερον δέ έστι καλ έκ τῶν νῦν καλ έκ τῶν προειρημένων. περλ μὲν οὖν τῶν κατὰ τὰς

διαφοράς άλις.

VI. Σπείρειν δὲ ξυμφέρει πάντα μάλιστα μὲν ἐν τοῖς ὡραίοις ἀρότοις· οὐ μὴν ἀλλὰ καὶ ἐν ξηρᾳ τινες καταβάλλουσι καὶ οὐχ ἤκιστα πυροὺς καὶ κριθὰς ὡς μάλιστα αὐταρκεῖν δυνάμενα, ὅπου μὴ ὅρνισιν ἡ ἄλλοις θηρίοις ἐπισινὴς ἡ χώρα. δοκεῖ γὰρ ὡς ἐπὶ πῶν ὁ πρῶτος ἄροτος ἀμείνων εἶναι, χείριστος δὲ σπόρος ἐν ταῖς ἡμιβρόχοις· ἀπόλλυται γὰρ καὶ ἐκγαλακτοῦται τὰ σπέρματα, καὶ ἄμα ξυμβαίνει πόαν ἀναφύεσθαι πολλήν. μετὰ δὲ τὴν σπορὰν ὕδωρ ἐπιγίνεσθαι πᾶσι ξυμφέρει, πλὴν ὅσα δυσβλαστῆ γίνεται μᾶλλον, ὡσπερ ὅ τε κύαμος δοκεῖ καὶ τῶν θερινῶν σήσαμον καὶ κύμινον καὶ ἐρύσιμον.

Πυκνοσπορείν δὲ καὶ μανοσπορείν καὶ πρὸς τὰς χώρας βλέποντα χρή· πλείον γὰρ ἡ πίειρα καὶ ἀγαθὴ δύναται φέρειν τῆς ὑφάμμου τε καὶ λεπτῆς. καίτοι λέγεταί τις λόγος ὡς ὁτὲ μὲν πλέον ὁτὲ δὲ ἔλαττον ἡ αὐτὴ δέχεται χώρα· καὶ οἰωνίζονται τὸ πλέον ὡς οὐκ ἀγαθόν, πεινῆν γὰρ εὐθύς φασι τὴν γῆν· οὖτος μὲν οὖν ἴσως εὐηθέστερος λόγος. εἰ δὲ τις πρὸς τὰ σπέρματα θεωροίη καὶ μάλιστά γε πρὸς αὐτοὺς τοὺς τόπους ἄμα τῷ ἐδάφει καὶ τὴν θέσιν ἀναθεωρῶν τήν γε πρὸς τὰ πνεύ-

i.e. after the rains.

¹ αὐταρκεῖν Ald.; ἀνταρκεῖν U.

² ἐπισινὴs conj. Dalec.; ἐπινὴs UMAld.; obnoxia G.

ENQUIRY INTO PLANTS, VIII. v. 4-vi. 2

is clear both from what is said now and from what was said before. Enough then about the points of difference.

Of sowing, manuring, and watering.

VI. It is expedient to sow all these, if possible, at the early seed-time; however some plant the seed even in dry ground, and especially wheat and barley, on the theory that they are most likely to hold their own 1 at a time when the ground is not infested 2 with birds or other creatures. For it appears that in general the first sowing is better, and worst that which is made in half-soaked ground; 3 for then the seeds perish and become 'milky'; 4 moreover many weeds come up at that time. After the sowing however it is beneficial for all that rain should fall on them, except in the case of some which appear to germinate then with more difficulty, as seems 5 to be the case with beans, and among summer crops with sesame cummin and erusimon.

⁶ As to sowing thickly or scantily one should have regard to the soil as well as to other considerations; for a fat good soil can bear more than one which is sandy and light. However there is a saying that the same soil can take at one time more, at another less seed; and in general the former condition is taken as an unfavourable omen, for then they say at once that the soil is hungry; however this is perhaps a rather foolish saying. If a man should have regard to the kind of the seed and especially to the actual situation, considering the aspect in respect of winds

177

VOL. II. N

cf. γαλάκτωσις, C.P. 4. 4. 7 and 8.
 δοκεί conj. W.; ἐδόκει Ald.
 Plin. 18. 196.

ματα καὶ τὸν ήλιον, οἰκειότερον αν λαμβάνοι τὰς διαφοράς.

- 3 'Ανὰ λόγον δὲ ἔχει καὶ ἡ κόπρισις τοῖς σπόροις πρὸς τὰς χώρας νειὸς δ' ἀμείνων ἡ χειμέριος τῆς ἐαρινῆς. ἐνιαχοῦ δὲ οὐ ξυμφέρειν βαθεῖαν ἀροτριᾶν, ὥσπερ καὶ ἐν Συρία, δι' ὁ μικροῖς ἀρότροις χρῶνται. παρ' ἄλλοις δὲ τὸ λίαν ἐξεργάζεσθαι βλάπτει, καθάπερ ἐν Σικελία, δι' ὁ καὶ τῶν ξένων ὡς ἔοικε πολλοὶ διαμαρτάνουσι. πάντα μὲν οὖν πρὸς τὰς χώρας.
- Διαιροῦσι δὲ καὶ τὰ σπέρματα ποία ποίον πρόσφορον ἐν γὰρ ταῖς χειμεριναῖς πυρὸν μᾶλλον ἢ κριθήν, καὶ ὅλως σῖτον ἢ χεδροπὰ κελεύουσιν ἐν ταῖς χέρσοις καὶ διὰ χρόνου κινουμέναις· καὶ γὰρ αὖται πυρὸν φέρουσι μᾶλλον ἢ κριθήν. δέχεται δὲ καὶ ἐπομβρίαν μᾶλλον πυρὸς τῆς κριθῆς, καὶ ἐν τοῖς ἀκόπροις φέρει μᾶλλον. ὡσαύτως δὲ καὶ αὐτῶν τῶν πυρῶν ποῖος τῆ ποία πρόσφορος, οἶον ἀγαθῆ καὶ πιείρα καὶ ψαφαρᾶ καὶ λεπτῆ <καὶ> ταῖς ἄλλαις ὁμοίως.
- 5 "Υδωρ δὲ ὅταν μὲν χλοηφορήση καὶ κυήση πλεῖον ἄπασι ξυμφέρει· ἀνθοῦσι δὲ πυροῖς μὲν καὶ κριθαῖς καὶ τοῖς σιτώδεσι βλαβερόν· ἀπόλ-

¹ αν λαμβάνοι conj Sch.; αναλαμβάνοι Ald. Η.

² κόπρισις conj. Sch.; κόπρησις Ald.

 ³ cf. C.P. 3. 20. 7.
 4 cf. C.P. 3. 20. 5.
 5 πάντα μὲν οδν M; ταῦτα μὲν Ald. H.; ταῦτα μὲν οδν conj.
 Sch. followed by W.

⁶ κελεύουσιν conj. W.; και δλως Ald. H.

 $^{^7}$ κινουμέναις conj. Sch. (cf. C.P. 3. 21. 4, ή διὰ χρόνου γεωργουμένη γ ή); κενουμέναις UAld; καινουμέναις Vin.

ENQUIRY INTO PLANTS, VIII. vi. 2-5

and sun, as well as the soil itself, he would more

properly gauge1 the differences.

Similarly manuring ² for the sown crops should be done with regard to the soil; and it is better to turn up fallow ⁸ land in winter than in spring. And there are some ⁴ places in which deep ploughing is not expedient, as in Syria; wherefore they use small plough-shares. In other parts to work the ground too much is injurious, as in Sicily: wherefore many settlers in the country, it appears, make a mistake. From every point of view ⁵ therefore the soil must be considered.

The seeds are also classified according as each suits a particular soil; in wintry lands wheat is sown rather than barley, and in general they say 6 that corn rather than leguminous plants should be sown in barren soils which are only disturbed 7 at long intervals; and such soils bear wheat better than barley. 8 Moreover wheat welcomes abundant rain 9 more than barley, and bears better on land which is not manured. In like manner they distinguish among wheats themselves which suits which kind of soil, namely which grows best in good 11 fat soil and which in crumbling light soil, and 12 so on with other kinds of soil.

¹³ More abundant rain is beneficial to all crops when they have come into leaf and formed the flower; however it is harmful to wheats and barleys and other cereals when they are actually in flower; for

 $^{^8}$ τη̂s κριθη̂s conj.W.; καὶ κριθη̂s UM; ή κριθὴ Ald.; η κριθὴ H. 9 cf. C.P. l.c.

¹⁰ Explained C.P. l.c.

¹¹ ἀγαθη conj. Casaub: so Vin.; ἀγαθη Ald. (and so with the other datives).

12 καὶ add. St.

¹⁸ Plin. 18, 151 and 152.

λυσι γάρ· ὀσπρίοις δ' ἀβλαβὲς πλὴν ἐρεβίνθων· οὐτοι γὰρ ἀποκλυσθείσης τῆς ἄλμης ἀπόλλυνται σφακελίζοντες καὶ ὑπὸ καμπῶν κατεσθιόμενοι· ἰσχυρότερος δὲ ὁ μέλας ἐρέβινθος καὶ ὁ πυρρὸς τοῦ λευκοῦ· συμφέρει δέ, φασίν, ἐν τοῖς ἐφύδροις τόποις ὀψὲ σπείρειν αὐτόν. κύαμος δὲ ἀνθῶν μάλιστα φιλεῖ βρέχεσθαι, δι' ὁ καὶ οὐκ ἐθέλουσιν ὀψισπορεῖν, ὅσπερ εἶπομεν, ὅτι πολὺν ἀνθεῖ· μετὰ δὲ τὴν ἀπάνθησιν ὀλίγου πάμπαν ὕδατος δεῖται· σύνεγγυς γὰρ ἡ τελείωσις. ἀλλ' ὅταν ἀδρυνθῆ καὶ βλάπτειν δοκεῖ τὰ σιτώδη καὶ

κριθην δὲ πυροῦ μᾶλλον.

και Κυν Αίγύπτω δὲ καὶ Βαβυλῶνι καὶ Βάκτροις, ὅπου μὴ ὕεται ἡ χώρα <ἡ> σπανίως, αὶ δρόσοι τὸ ὅλον ἐκτρέφουσιν. ἔτι καὶ οἱ περὶ Κυρήνην καὶ Εὐεσπερίδας τόποι. καιριώτατα δὲ πᾶσιν ὡς ἀπλῶς εἰπεῖν τὰ ἠρινά· δι ὁ καὶ ἡ Σικελία πολύσιτος· πολλὰ γὰρ τοῦ ἠρος καὶ μαλακὰ γίνεται, τοῦ δὲ χειμῶνος ὀλίγα. ζητεῖ δὲ ἡ μὲν λεπτόγεως πολλὰ κατὰ μικρόν· ἡ δὲ πίειρα καὶ πλῆθος μὲν ἐνεγκεῖν δύναται καὶ ἀϋδρίαν—πρὸς δὲ τὴν χώρας ἀϋδρίαν πόντια πνεύματα καὶ αὐραι δοκοῦσι ξυμφέρειν, ἄλλα δὲ παρ' ἄλλοις τοιαῦτα, καθάπερ καὶ πρότερον εἴρηται,—ὡς ἐπὶ τὸ πᾶν δὲ μᾶλλον αὐχμὸς ἡ ἐπομβρία ξυμφέρει τῷ σίτῳ· το γὰρ ὅμβροι καὶ ἄλλως ἐναντίοι καὶ πολλάκις αὐτὰ τὰ σπέρματα διαφθείρουσιν, εἰ δὲ μὴ πλῆ-

1 σφακελίζοντες: cf. 4. 14. 4.

3 cf. C.P. 3. 22. 3.

² δ πυρρός τοῦ λευκοῦ conj. Scal. from G and Plin. 18. 124; δ λευκός τοῦ πυροῦ UAld; δ λ. τ. πυρροῦ Η.; δ λ. τ. πυρος Μ.

⁴ δείται conj. Sch.; δείσθαι Ald. H.

ENQUIRY INTO PLANTS, VIII. vi. 5-7

it destroys the flower. But to pulses it is harmless, except to chick-peas; for these, if the salt is washed off them, perish from rot 1 or from being eaten by caterpillars. However the black and the red 2 chick-pea are stronger than the white, and it is beneficial, they say, to sow this crop late in moist soil. The bean 3 likes especially to receive rain when it is in flower; wherefore men are unwilling, as we said, to sow it late, because it flowers for a long time; but after it has shed its flowers, it needs 4 very little water, since its time of maturity is now near. But, when cereals have matured, it appears that water actually injures them, and barley more than wheat.

In Egypt Babylon and Bactra, where the country receives no rain, or 5 but little, the dews are sufficient nourishment; and so is it also 6 in the regions about Cyrene and the Euesperides. However to all, generally speaking, it is the spring rains which are the most seasonable; and that is why Sicily is rich in corn; for there is abundance of soft rain in spring and little of it in winter. A light soil requires plenty of rain, but little at a time; while that which is fat can indeed bear both an abundance of rain and a drought; (for a droughty country seawinds 7 and breezes seem to be helpful, and various breezes of this kind prevail in various countries, as has been said already). Yet in general drought suits corn better than excessive rain; for heavy showers, apart from the harm which they do in other ways, often actually destroy the seed, or at

⁵ h add. Scal. from G: so Vin.

⁶ έτι conj. St. from G (?); ἐπεὶ Ald.

⁷ πόντια conj. Sch.; πάντα Ald. cf. 8. 7. 6.

θός γε ποιοῦσι βοτάνης, ὥστε καταπνίγεσθαι καὶ ἀτροφεῖν.

VII. Τῶν μὲν οὖν ἄλλων σπερμάτων οὐδὲν εἰς ἄλλο πέφυκε μεταβάλλειν φθειρόμενον, πυρὸν δὲ καὶ κριθὴν εἰς αἰράν φασι καὶ μᾶλλον τὸν πυρόν, γίνεσθαι δὲ τοῦτ' ἐν ταῖς ἐπομβρίαις καὶ μάλιστα ἐν τοῖς εὐύδροις καὶ ὀμβρώδεσι χωρίοις. ὅτι δ' οὐκ ἔστιν ἠρινὸν ἡ αἰρα καθάπερ ἡ ἄλλη πόα, πειρῶνται γάρ τινες τοῦτο λέγειν, ἐκεῖθεν δῆλον· εὐθὺς γὰρ τοῦ χειμῶνος φανερὰ γίνεται πεφυκυῖα· καὶ διαφέρει πολλοῖς· ἔχει γὰρ τὸ φύλλον στενὸν καὶ δασὺ καὶ λιπαρόν, καὶ τοῦτων ἰδιώτατον τὸ λιπαρόν· ἡ γὰρ δασύτης καὶ τοῖς τοῦ αἰγίλωπος ὑπάρχει, ἀλλ' ἐκφανὴς γίνεται ἐπὶ τοῖς τοῦ αἰγίλωπος τοῦ ἦρος. τοῦτο μὲν οὖν ἴδιον τούτων, καὶ ἔτι τοῦ λίνου· καὶ γὰρ ἐκ τούτου φασὶ γίνεσθαι τὴν αἰραν.

2 Τοῦ δὲ ἐρεβίνθου πρὸς τὰ ἄλλα χεδροπὰ τό τε περὶ τὴν ἄνθησιν λεχθὲν καὶ τὸ τάχιστα τελειοκαρπεῖν ἰσχυρότατον ον καὶ ξυλωδέστατον, καὶ

² πόα: ? grasses; cf. 8. 6. 1.

4 εὐθὺς γὰρ τοῦ conj. Sch.; εὐθὺς τὸ τοῦ Ald.

182

¹ Plin 18. 149 and 150; cf. C.P. 4. 5. 2.

³ πειρώνται γάρ τινες Η.; ἀπειρώνται αίτιώνται γάρ τινες U; ὰ πειρώνται αίτιώνται γάρ τινες PM: so also Ald. Bas. Can. with mark of corruption.

ENQUIRY INTO PLANTS, VIII. vi. 7-vii. 2

least cause a luxuriant growth of leafage, so that the grain is choked and becomes abortive.

Of the degeneration of cereals into darnel.

VII. 1 Now, while it is not the nature of any other of these seeds to degenerate and change into something else, they say that wheat and barley change into darnel, and especially wheat; and that this occurs with heavy rains and especially in well-watered and rainy districts. But that darnel is not a plant of the spring, like other weeds 2 (for some endeavour 3 to make this out) is clear from the following consideration: it springs up and becomes noticeable directly 4 winter comes; and it is distinguished in many ways; the foliage 5 is narrow abundant and glossy, and this gloss is the most marked of these differences; (the 6 leaves of aigilops 7 are indeed also abundant, 8 but this character does not shew itself in them till spring). This then is peculiar to the seeds of wheat and barley, and also to those of flax; for that too, they say, turns into darnel.

Of the peculiar character of chick-pea.

A peculiarity of chick-pea as compared with other leguminous plants is that which has been mentioned as to its flowering; and also the fact that it is the quickest to mature its fruit, being very strong and woody; and again there is the fact that in

⁷ Plin. 18. 155.

Digitized by Google

⁵ cf. C.P. 4. 4. 11. ⁶ τοι conj. Sch.; τη Ald.

ἀλλ'... alγίλωπος: text a makeshift. Wanting in Ald.
 and all MSS. except U; ἀλλ' ἐκφανεῖς γίνονται καὶ τοῖς τοῦ αἰγ.
 U; ἐκφανὴς γίνεται conj. Sch.; ἐπὶ for καὶ conj. W.

τὸ ὅλον μὴ ποιεῖν νειὸν <ώς> καρπιζόμενον τὴν δὲ πόαν ἐξαπόλλυσι καὶ μάλιστα δὲ καὶ τάγιστα τὸν τρίβολον. ὅλως δὲ οὐδὲ ἡ τυγοῦσα δύναται φέρειν αὐτόν, άλλα μελάγγειόν τινα δεῖ καὶ πίειραν είναι. των δε άλλων ή αρίστη νειος από των κυάμων καίπερ πυκνοσπορουμένων και πολύν καρπον φερόντων.

- Τὰ δὲ ἐν τοῖς θερινοῖς ἀρότοις ὀλίγου δεῖ πάντα. φασί δε και τα ναματιαία συμφέρειν μάλλον αὐτοῖς τῶν ἐκ διός, μέλινοι δὲ καὶ κέγγροι έλάττους ὕδατος έὰν γὰρ ἔχωσι πλείον φυλλοβολούσιν. ἰσχυρότερον δὲ ὁ κέγχρος οἱ δὲ μέλινοι γλυκύτεροι καὶ ἀσθενέστεροι. σήσαμον δὲ οὐδὲν <ζῶον> ἐσθίει χλωρὸν οὐδὲ θέρμον. εἰ δὲ μηδ' έρύσιμον μηδέ δρμινον σκεπτέον καὶ ταῦτα πικρά. ἔστι δὲ τὸ μὲν ἐρύσιμον ὅμοιον σησάμφ καὶ λίπος ἔχει· τὸ δὲ ὅρμινον κυμινῶδες μέλαν· σπείρεται δὲ ἄμα καὶ τὸ σήσαμον. περὶ μὲν οὖν τούτων σκεπτέον.
- Έν δὲ ταῖς ἀγαθαῖς χώραις πρὸς τὸ μὴ φυλλομανείν επινέμουσι καὶ επικείρουσι τὸν σίτον,

¹ Lit. 'does not make fallow land.' cf. C.P. 4. 8. 3.

³ ώς καρπιζόμενον Ι conj. after W. (καρπιζόμενον την γην); καρπός U; καρπός M; καρπόν Ald. cf. C.P. l.c. and 4. 8. 1; 4. 8. 3: μη καρπίζεσθαι την γην άλλα νειον ποιείν (? < νέοις >) καρποῖς, 'for fresh crops.'

⁴ cf. C.P. 4. 8. 3. 3 δè conj. W.; γε Ald.

ο το τους. τι, γε Αια.

δ η αρίστη νειος conj. W. (cf. 8. 9. 1; C.P. 4. 8. 1); χειρίστη νήπιος U; χειρίστην νήπιος MP; καλλίστη νειος Ald. cf. also C.P. 3, 20, 7.

ENQUIRY INTO PLANTS, VIII. VII. 2-4

general it does not reinvigorate the ground,1 since it exhausts 2 it; but it destroys weeds,8 and above all and soonest caltrop. And in general 4 it is not every kind of soil which suits it: the soil should be black and fat. Of the other leguminous plants the bean best 5 reinvigorates the ground, even if it is sown thick and produces much fruit.

Of special features of 'summer crops.'

All those crops 6 sown at the summer seed-time need little water, and they say also that spring water is better for them than rain water; and Italian millet 8 and millet need less water, for, if they have too much, they shed their leaves. Millet is the robuster plant, Italian millet is sweeter and less robust. Sesame and lupin are not eaten green by any animal 9; whether the same is true of erysimon and horminon is matter for enquiry; for these too are bitter. Erysimon is like sesame and is oily; horminon is like cummin and black, and is sown at the same time as sesame. These matters then require investigation.

Of treatment of cereals peculiar to special localities.

10 In good soils to prevent the crop running wildly to leaf they graze and cut down the young corn,

⁶ Plin. 18, 96 and 101.

⁷ δλίγου, sc. δδατος, but the omission is strange; perhaps due to misunderstanding of onlyou bei by a scribe. Sch. joins the words τὰ δὲ . . . πάντα to the last sentence, and supplies καρπίζεται την γην (δλίγου δε $\hat{i} = \text{almost}$).

 ⁸ μέλινοι Ald. Η.; έλυμοι Vin. cf. 8. 1. 1. n.
 ⁹ ζφον add. Sch. from G and Plin. 18. 96. cf. C. P. 6. 12. 12.

¹⁰ Plin. 18. 157-162,

ώσπερ καὶ ἐν Θετταλία. συμβαίνει δ' αν μὲν έπινέμωσιν όποσακισούν μηδέν άλλοιούσθαι τὸν καρπόν, αν δε επικείρωσιν απαξ μόνον εξίστασθαι τὸν πυρὸν καὶ γίνεσθαι μακρὸν καὶ οὐχ άδρόν, δν καλούσι καμακίαν, καὶ οὐκ ἀποκαθίστασθαι πάλιν σπειρόμενον τοῦτο μεν οὖν ώς παύροις συμβαίνου Θετταλοί λέγουσιν. ἐν Βαβυλώνι δὲ άεὶ καὶ ὥσπερ τεταγμένως ἐπικείρουσι μὲν δίς, τὸ δὲ τρίτον τὰ πρόβατα ἐπαφιᾶσιν οὕτω γὰρ φύει τον καυλόν, εί δὲ μὴ φυλλομανεί γίνεται δὲ μὴ καλώς έργασαμένοις πεντηκονταχόα, τοις δε έπιμελως έκατονταχόα. ή δὲ ἐργασία τὸ ὡς πλεῖστον χρόνον εμμένειν το ύδωρ, δπως ίλυν ποιήση πολλήν πίειραν γάρ ούσαν καὶ πυκνήν την γήν δεί ποιήσαι μανήν. ύλην δε ου φέρει και πόαν ώσπερ έν Αίγύπτω. τὰ μὲν οὖν τοιαῦτα γώρας ἀρετής.

Φύεται δὲ καὶ ἀπὸ ριζῶν πυρὸς καὶ κριθὴ πολλαχοῦ τῷ ὑστέρῳ ἔτει αὐτοετὴς δὲ καὶ ἀπὸ τῶν εἰς κράστιν κειρομένων ἐτέρου καλάμου παραβλαστάνοντος. ὡσαύτως δὲ κᾶν ὑπὸ χειμῶνος ἐκπαγῆ· παραβλαστάνει γὰρ ὑδάτων ἐπιγινομένων ὁ δὲ στάχυς ἀτελὴς καὶ μικρὸς ἀπὸ τῶν τοιούτων. βλαστάνουσι δὲ τῷ ὑστέρῳ ἔτει καὶ ἀπὸ τῶν καταπονουμένων καὶ συμπατουμένων, ὥστε μηδὲν εἶναι δῆλον ὡς εἰπεῖν, οἶον ὅταν

² Text perhaps defective: cf. Plin. l.c.

¹ λλυ conj. Sch. from Plin. 18. 162; ὅλην Ald. Η.

ENQUIRY INTO PLANTS, VIII. vii. 4-5

for instance in Thessalv. And the result is that, however often they graze it, the crop is not impaired; while if they cut it down not more than once, the wheat changes in character and becomes tall and weak-what they call 'long-shafted' corn, and, if seed of this is sown, it does not recover its character. This the Thessalians tell of as having occurred in a few cases. At Babylon however they cut it down twice always and as it were systematically, and after that they let the sheep on to it: for in that case it makes its straw, but otherwise it runs wildly to leaf; and, if the ground is ill cultivated, it produces fifty fold, if it is carefully cultivated, a hundred fold. And the 'cultivation' consists in letting the water lie on it as long as possible, so that it may make much silt1; for the soil being fat and close must be made open. And at Babylon 2 the ground does not produce weeds and grasses, as it does in Egypt. Such are the things which depend on the quality of the soil.

Of cereals which grow a second time from the same stock.

⁸ Wheat and barley also in many places grow from the root in the next year, or in the same year from crops cut down for fodder, since a second haulm shoots up. The like happens also if the plant has been nipped by winter; for it shoots again when rain comes; but such plants produce an ear which is imperfect and under-sized. There is also new growth the next year from plants which are roughly treated or trodden down ⁴ so that hardly anything remains visible, as happens when an army has marched over

³ cf. C.P. 4. 8. 5. ⁴ cf. C.P. l.c.

διέλθη στρατόπεδον, καὶ οἱ στάχυες μικροὶ καὶ τούτων, οθς καλοῦσιν ἄρνας· τῶν δὲ χεδροπῶν οὐδὲν δύναται τοιοῦτον ποιεῖν ἡ οὐχ ὁμοίως. καὶ αἱ βλαστήσεις τοσαυταχῶς.

Πρὸς αὔξησιν δὲ καὶ τροφὴν μέγιστα μὲν ἡ τοῦ ἀέρος κρᾶσις συμβάλλεται, καὶ ὅλως ἡ τοῦ ἔτους κατάστασις· εὐκαίρων γὰρ ὑδάτων καὶ εὐδιῶν καὶ χειμώνων γινομένων ἄπαντα εὕφορα καὶ πολύκαρπα, κὰν ἐν ἀλμώδεσι καὶ λεπτογείοις ἢ·δι' δ καὶ παροιμιαζόμενοι λέγουσιν οὐ κακῶς ὅτι ἔτος φέρει οὐχὶ ἄρουρα."

Μέγα δὲ καὶ αὶ χῶραι διαφέρουσιν οὐ μόνον τῷ πίειραι καὶ λεπταὶ καὶ ἔπομβροι καὶ αὐχμώδεις <εἶναι> ἀλλὰ καὶ τῷ ἀέρι τῷ περιέχοντι καὶ τοῖς πνεύμασιν. ἔνιαι γὰρ οὖσαι λεπταὶ καὶ φαῦλαι τελεσφοροῦσι διὰ τὸ πρὸς τὰ πνεύματα τὰ πόντια κεῖσθαι καλῶς. ἄλλα δὲ ἄλλαις τοιαῦτα, καθάπερ πολλάκις εἴρηται. ταῖς μὲν γὰρ τὰ ζεφυρικὰ ταῖς δὲ τὰ βόρεια ταῖς δὲ τὰ νότια.

Συμβάλλεται δὲ καὶ οὐ μικρὰ ἡ ἐργασία καὶ μάλισθ' ἡ <πρὸ> τοῦ σπόρου· κατεργασθεῖσα γὰρ ράδίως ἐκφέρει. καὶ ἡ κόπρος δὲ μεγάλα βοηθεῖ τῷ διαθερμαίνειν καὶ συμπέττειν· προτρέχει γὰρ τὰ κοπριζόμενα τῶν ἀκόπρων καὶ εἴκοσιν ἡμέραις·

T88

¹ τούτων conj. Sch.; τούτους Ald. H.

² cf. Lewis and Short s.v. agna.
³ C.P. l.c. gives the reason.

⁴ τοσαυταχώς conj. Scal.; τοσαυταχείς Ald. H.

ENQUIRY INTO PLANTS, VIII. vii. 5-7

the field; the ears in such cases 1 too are undersized and are called 'lambs.' 2 But no kind of leguminous plant 3 can do anything of the kind, or at least not to the same extent. In these various ways 4 may new growth occur.

Of the effects of climate, soil, and manuring.

For growth and nourishment the climate is the most important factor, and in general the character of the season as a whole; for when rain, fair weather and storms occur opportunely, all crops bear well and are fruitful, even if they be in soil which is impregnated with salt or poor. Wherefore there is an apt proverbial saying 5 that "it is the year which bears and not the field."

But the soil also makes much difference, according as it is ⁶ fat or light, well watered or parched, and it also makes quite as much difference what sort of air and of winds prevails in that region; for some soils,⁷ though light and poor, produce a good crop because the land has a fair aspect in regard to sea breezes. But, as has been repeatedly said already, the same breeze has not this effect in all places; some places are suited by a west, some by a north, some by a south wind.

Again the working of the soil and above all that which is done before 8 the sowing has an important effect; for when the soil is well worked it bears easily. Also dung is helpful by warming and ripening the soil, for manured land gets the start by as much as twenty days of that which has not been

8 πρδ add. W. cf. C.P. 3. 20. 6.

⁵ Quoted also C.P. 3. 23. 4.

⁶ elvai add. Sch. 7 cf. C.P. 3. 23. 5.

άπασι δε οὐ ξυμφέρει καὶ χρήσιμος οὐ μόνον τοῖς περὶ τὸν σῖτον ἀλλὰ καὶ τοῖς ἄλλοις πλὴν πτερίδος, ταύτην δε φθείρειν φασὶν ἐπιβαλλομένην. ἀπόλλυται δε ἡ πτερὶς καὶ ἐπικοιμωμένων τῶν προβάτων, ὡς δε τινες λέγουσι καὶ ἡ Μηδικὴ διὰ τὴν κόπρον καὶ τὸ οὖρον.

VIII. Των δε σπερμάτων εκαστα και προς την της χώρας φύσιν άρμόττει, καὶ όλως γένη πρὸς γένος καὶ ἐν αὐτοῖς τοῖς ὁμογενέσιν, ἃ δὴ πειρώνται διαιρείν. μεταβάλλει δὲ τὰ ξενικὰ τών σπερμάτων μάλιστα μεν έν τρισίν έτεσιν είς τὰ έπιχώρια. συμφέρει δὲ ἐκ τῶν ἀλεεινῶν εἰς τὰ μικοον ήττον άλεεινα και έκ των ψυχεινων ανα λόγον ποιείσθαι την μεταβολήν. τὰ δ' ἐκ τῶν δυσχειμερινών έν τοις πρωίοις όψε ἀποχείται, ωστ' ἀπ' αὐχμοῦ φθείρεται, ἐὰν μὴ ὄψιον ὕδωρ σώση. διὰ τοῦτο καὶ εὐλαβητέον φασὶ τὸ μίσγειν τὰ ξενικά τοῖς ἐπιχωρίοις ἐὰν μὴ ἐξ ὁμοίας, ὅτι ασύμφωνα τη χώρα κατά τὸν σπόρον καὶ κατά την γένεσιν, ώστε καὶ έργασίας έτέρας δείται τάς τε τής γής διαφοράς καὶ τὰς τῶν σπερμάτων δυνάμεις καὶ ἔτι τὰς ἐκάστων ὥρας.

¹ cf. Col. 2. 2. 13. The reference is perhaps to fern grown for litter, or possibly for medicinal use. cf. 9. 20. 5.

² χώραs conj. Sch.; ωραs Ald. ³ à conj. Dalec.; àν Ald.

⁴ ψυχεινών conj. W.; ψυχικών UM; ψυχρών Ald.

ο ἀποχείται conj. Sch., cf. ἀπόχυσις 8. 3. 4; ἀποκείται Ald. cf. 4. 4. 10.

ENQUIRY INTO PLANTS, VIII. vii. 7-viii. 2

manured. However manure is not good for all crops; and further it is beneficial not only to corn and the like but to most other things, except fern, which they say it destroys if it is put on. (Fern is also destroyed if sheep lie on it, and, as some say, lucerne is destroyed by their dung and urine.)

Of different qualities of seed.

VIII. There is a particular kind of soil 2 which best suits each kind of seed, whether we compare one class with another or those of the same class; and attempts are made to distinguish these.3 Foreign seeds change into the native sorts in about three It is well that they should be imported from a warm climate to one that is rather less warm, or from a cold one 4 to one that is rather less cold. Those imported from a wintry climate, if they be those of early crops, are late in coming into ear,5 so that they get destroyed by drought unless rain late in the season saves them. Wherefore they say that one should take good heed not to mix foreign with native seeds, unless they come from a similar place, since 6 they do not agree with the soil 7 as to the time of being sown and of germinating, and accordingly need different cultivation; and so that one should take good heed to the differences of soil, the properties of the seed, and further the seasons appropriate to each.

When however there is a good season, the grain also is fuller.⁸ For instance at Athens the barley pro-

δτι conj. Sch.; ἔτι UMAld.
 χώρα conj. Sch.; ἄρα UMAld.

^{*} πολυνοστότερα: cf. νόστιμος, C.P. 4. 13. 2, Geop. 2. 16. 1, and other reff. in Sch.'s exhaustive note.

τὰ πλείστα ποιούσιν ἄλφιτα κριθοφόρος γὰρ άρίστη τοῦτο δ' οὐν ὅταν πλεῖσται γένωνται άλλ' όταν λάβη τινὰ κρᾶσιν. ἐν δὲ τῆ Φωκίδι περί 'Ελάτειαν οί πυροί ποιούσιν ήμιόλια τὰ άλευρα, καὶ ἐν Σόλοις τῆς Κιλικίας καὶ οἱ πυροὶ καὶ αί κριθαί· καὶ παρ' ἄλλοις ἄλλα πρὸς ἄπερ εὐφυὴς ἐκάστη. βελτίω μὲν οὖν καὶ γείρω τὰ σπέρματα καὶ διὰ τὴν ἐργασίαν καὶ διὰ τὴν γῆν γίνεται και γάρ άπαγριοθται και ήμεροθται, καθάπερ τὰ δένδρα: καὶ ὅλως μεταβάλλει <κατὰ> την χώραν, ώσπερ τινα των δένδρων εύθυς έστηκε πρὸς τὸ χεῖρον.

3 Γένος δ' όλον εξαλλάττειν είς ετερον οὐδεν άλλο πέφυκε πλην τίφη καλ ζειά, καθάπερ εἴπομεν έν τοις πρώτοις λόγοις, καὶ ή αίρα δ' έκ τῶν πυρών καὶ κριθών διαφθειρομένων ή εἰ μή τοῦτο άλλὰ φιλεί γε μάλιστα ἐν τοῖς πυροῖς γίνεσθαι, καθάπερ καὶ ὁ μελάμπυρος ὁ Ποντικὸς καὶ τὸ των βολβων σπέρμα, καὶ ἄλλα δὲ ἐν ἄλλοις των σπερμάτων έπει και ο αιγίλωψ δοκει μαλλον έν ταις κριθαις, έν δὲ τοις φακοις ἄρακος τὸ τραχύ καὶ σκληρόν, ἐν δὲ ταῖς ἀφάκαις ὁ πελεκῖνος ομοιον τη όψει τφ πελέκει σχεδον δε καθ' εκαστόν έστι τὸ συνεκτρεφόμενον καὶ συνανα-

3 (eid conj. Scal.; Vea Ald. H. 4 2. 4. 1.

¹ κατὰ add. W. cf. 2. 4. 1.
2 τινα conj. W.; τε P; τφ Ald.; τδ H. Vin. Vo. cf. 2. 2. 6.

ENQUIRY INTO PLANTS, VIII. viii. 2-3.

duces more meal than anywhere else, since it is an excellent land for that crop; and this is so, not merely when a very large crop is sown, but when the weather has been favourable for it. And in Phocis about Elateia the wheats produce half as much meal again as elsewhere; while at Soli in Cilicia this is true of both wheat and barley; and in other parts there are other crops for which the soil is severally well adapted. Wherefore grain turns out better or worse because of the soil as well as because of cultivation; for in some places it changes into the cultivated from the wild form, or the reverse, like trees; and in general it changes according 1 to the soil in which it is grown, just as some 2 trees, when transplanted, forthwith deteriorate.

Of degeneration of cereals, and of the weeds which infest particular crops.

But no kind can change altogether into another, except one-seeded wheat and rice-wheat,³ as we said ⁴ in our previous discussions, and darnel which comes from degenerate wheat and barley: at least, if this is not the true account, darnel loves chiefly to appear among wheat, as does the Pontic ⁵ melampyros and the seed of purse-tassels,⁶ even as other seeds appear in other crops; thus aigilops seems to grow for choice among barley, and among lentils the rough hard kind of arakos, while among tares occurs the axe-weed,⁷ which resembles an axe-head in appearance. Indeed in the case of nearly every crop there is a plant which grows up with it and

⁷ Plin. 18. 155; 27. 121; Diosc. 3. 130; Hesych. s.v. βέλλεκυς.

193

0

⁵ cf. 8. 4. 6, where μ ελάμπυρον was said to be peculiar to Sicily. 6 cf. C.P. 4. 6. 1.

- μιγνύμενον εἴτε διὰ τὰς χώρας, ὅπερ οὐκ ἄλογον, εἴτε δι' ἄλλην τινὰ αἰτίαν. ἔνια δὲ καὶ φανερῶς ἐστι κοινὰ πλειόνων, ἀλλὰ διὰ τὸ μάλιστα ἔν τισιν εὐθενεῖν ἴδια τούτων φαίνεται, καθάπερ ἡ ὀροβάγχη τῶν ὀρόβων καὶ ἡ ἀπαρίνη τῶν φακῶν ἀλλὰ ἡ μὲν μάλιστα ἐπικρατεῖ τῶν ὀρόβων διὰ τὴν ἀσθένειαν ἡ δὲ ἀπαρίνη μάλιστα ἐν τοῦς φακοῦς εὐτροφεῦ τρόπον δέ τινα καὶ παραπλήσιόν ἐστι τῆ ὀροβάγχη <τῷ> ἐπιβάλλειν καὶ κατέχειν ὅλον ὥσπερ πλεκτάναις ἀποπνίγει γὰροὕτως, ὅθεν καὶ τοῦνομα εἴληφε.
 - Τὸ δ' ὑποφυόμενον εὐθὺς ἐκ τῆς ῥίζης τῷ κυμίνῷ καὶ τῷ βουκέρῷ τὸ αἰμόδωρον καλούμενον μᾶλλον ἰδίᾳ. ἔστι δὲ τὸ αἰμόδωρον μονόκαυλον οὐκ ἀπεμφερές [τῷ καυλῷ], πλὴν βραχύτερόν τε πολύ, καὶ ἄνωθέν τι κεφαλῶδες ἔχει ῥίζαν δὲ ὑποστρόγγυλον· οὐθὲν δὲ ἔτερον ἀφαυαίνεται παρὰ τὸ βούκερας. γίνεται δὲ ταῦτα ἐν ταῖς λεπταῖς οὐκ ἐν ταῖς πιείραις, ὥσπερ καὶ τῆς Εὐβοίας ἐν τῷ Ληλάντῷ μὲν οὐ γίνεται περὶ δὲ τὸν Κάνηθον καὶ εἴ τις ἄλλος τοιοῦτος τόπος.

² τῶ add. Sch.

¹ άλλην τινα conj. Sch.; άλλης τινα U; άλλην Ald.

³ πλεκτάναις conj. W.; πλεκτᾶνες U; πλεκτάνης M; πλεκτάνην Ald.: veluti brachiis G.

⁴ Plin. 19. 176, who however calls this αἰμόδωρον. See Index App. (26).

ENQUIRY INTO PLANTS, VIII. viii. 3-5

mingles with it, whether this is due to the soil, which is a reasonable explanation, or to some other 1 cause. Some plants of this character evidently attach themselves to more than one kind of crop, but, because they are specially vigorous in some one particular crop, they are thought to be peculiar to that one, as 'vetch-strangler' (dodder) to vetches and bedstraw to lentils. But the former gains the mastery over the vetches especially because of the weakness of that plant; and bedstraw is specially luxuriant among lentils; to some extent it resembles dodder, in that 2 it overspreads the whole plant and holds it fast as it were in coils, 3 for it is thus that dodder strangles the plant, and this is the origin of its name ('vetch-strangler').

The plant which springs up straight from the roots of cummin and the plant called broom-rape which similarly attaches itself to 'ox-horn' ⁵ (fenugreek) are somewhat more peculiar in their habits. ⁶ Broom-rape has a single stem, ⁷ and is not unlike . . . , ⁸ but is much shorter and has on the top a sort of head, while its root is more or less round; and there is no other plant which it starves except fenugreek. These plants grow in light and not in fat soils; thus in Euboea they do not occur at Lelanton, ⁹ but only about Kanethos ¹⁰ and in districts of like character.

⁵ Plin. 24, 184.

⁶ idla MSS.; ? Toia.

⁷ cf. C.P. 5. 15. 5, where the same is said of λειμόδωρον (cf. Plin. 19. 176). But Ald. Bas. Cam. give αἰμόδωρον here; hemodorum G.

⁸ τῷ καυλῷ probably conceals the name of a plant.

⁹ cf. Strabo, 10. 1. 9. L. is the name of a Euboean river in Plin. 4. 64.

¹⁰ cf. Strabo, 10. 1. 8, Ap. Rhod. 1. 77.

ταῦτα μὲν οὖν κοινὰ πλειόνων ὄντα κατισχύει μᾶλλον ἐν τοῖς εἰρημένοις διὰ τὴν ἀσθένειαν.

Τὸ δὲ τέραμον καὶ ἀτέραμον λέγεται μὲν ἐπὶ τῶν ὀσπρίων μόνον, οὐκ ἄλογον δὲ καὶ ἐπὶ τῶν σιτωδών παραπλήσιον ή καὶ ταὐτό τι συμβαίνειν, άλλὰ διὰ τὸ μὴ τὴν αὐτὴν εἶναι γρείαν οὐγ ομοίως εμφανές επεί ουδ' επί τούτων απάντων όμοίως άλλα μάλιστα έπι των κυάμων λέγεται καὶ φακῶν, εἴτ' οὖν καὶ μάλιστα πασχόντων είτε και διά την γρείαν φαινομένων. γίνεται γοῦν πλεοναχώς πολλαχοῦ γάρ τόποι τινές είσιν οι αίει φέρουσι τεράμονα και ἄλλοι πάλιν άτεράμονα· τὸ δὲ ὡς ἐπὶ πᾶν οἱ λεπτόγεω μᾶλλον 7 τεράμονα καὶ ἀέρος κατάστασίς τις ποιεί τὴν τοιαύτην παραλλαγήν σημείον δε ότι ταὐτά χωρία και δμοίως έργασθέντα φέρει ποτέ μέν τεράμονα <ποτε δε ἀτεράμονα.> περί Φιλίππους δε δ κύαμος λικμώμενος, εαν ύπο πνεύματος ενγωρίου ληφθή, τεράμων ὢν ἀτεράμων γίνεται. ταθτα μεν οθν μηνύει διότι πολλαχώς των αθτών

² πλεοναχῶς· πολλαχοῦ I conj. ; πλέον· πολλαχῶς MSS.

8 ποτè δè ἀτεράμονα add. Η. from G.

¹ cf. 2. 4. 2; C.P. 4. 12; Plin. 18. 155, who makes ateramum, teramum plants.

⁴ cf. C.P. 4. 12. 8; Plut. Quaest. Conv. 7. 2. 3; Plin. l.c. 196

ENQUIRY INTO PLANTS, VIII. viii. 5-7

The reason then why these plants, which attach themselves to more than one kind, grow stronger when attached to the plants specified, is that the latter are not robust.

Of the conditions in the seeds of pulses known as 'cookable' and 'uncookable,' and their causes.

¹The terms 'cookable' and 'uncookable' are only applied to pulses, but it is not unreasonable to suppose that conditions like those indicated, if not identical with them, occur also in cereals, though they are not so obvious, since these plants are not put to the same use. Indeed it is said that these terms are not applied even to all pulses alike, but chiefly to beans and lentils, either because these are specially subject to these conditions, or because the use to which they are put makes them more conspicuous. At all events the conditions occur for a variety of reasons; for in many parts 2 there are places which regularly produce seeds that are 'cookable,' while others again produce seeds that are 'uncookable'; in general however it is light soils which tend to produce the former. Now it is a certain condition of the climate which causes this variation: a proof of which is the fact that the same piece of land, tilled in the same manner, produces sometimes seeds that are 'cookable,' sometimes seeds that are 'uncookable.' 3 In the district of Philippi, if the beans, while being winnowed,4 are caught by the prevailing wind of the country, they become 'uncookable,' having previously been 'cookable.' These facts prove that for various reasons. of districts 5 which are close together, have the same

⁸ αὐτῶν conj. W.; δὲ τῶν Ald. cf. a similar expression 8.2.10.



χωρίων ένια σύνορα καὶ όμοίως καθήμενα καὶ οὐδεμίαν έχοντα κατὰ τὴν γῆν διαφορὰν τὸ μὲν τεράμονα τὸ δ' ἀτεράμονα φέρει, καὶ ἐνίοτε μόνον αὔλακος διοριζούσης.

ΙΧ. Καρπίζεται τὴν γῆν μάλιστα πυρὸς εἶτα κριθή, δι' ὁ καὶ ὁ μὲν ἀγαθὴν ζητεῖ χώραν ἡ δὲ κριθὴ δύναται καὶ ἐν ταῖς ψαφαρωτέραις ἐκφέρειν τῶν δὲ χεδροπῶν μάλιστα ἐρέβινθος καίπερ ἐλάχιστον χρόνον ἐν τῆ γῆ μένων, ὁ δὲ κύαμος, ὥσπερ ἐλέχθη, καὶ ἄλλως οὐ βαρὺ καὶ ἔτι κοπρίζειν δοκεῖ τὴν γῆν διὰ μανότητα καὶ εὐσηψίαν δι' ὁ καὶ οἱ περὶ Μακεδονίαν καὶ Θετταλίαν ὅταν ἀνθῶσιν ἀνατρέπουσι τὰς ἀρούρας.

Τῶν δὲ ὁμοιοπύρων καὶ ὁμοιοκρίθων, οἶον ζειᾶς τίφης ὀλύρας βρόμου αἰγίλωπος, ἰσχυρότατον καὶ μάλιστα καρπιζόμενον ἡ ζειά· καὶ γὰρ πολύρριζον καὶ βαθύρριζον καὶ πολυκάλαμον· ὁ δὲ καρπὸς κουφότατος καὶ προσφιλὴς πᾶσι τοῖς ζώοις. τῶν δὲ ἄλλων ὁ βρόμος· πολύρριζος γὰρ καὶ οὖτος καὶ πολυκάλαμος. ἡ δὲ ὀλύρα μαλακώτερον καὶ ἀσθενέστερον τούτων. ἡ δὲ τίφη πάντων κουφότατον· καὶ γὰρ καὶ μονοκάλαμον <καὶ λεπτοκάλαμον,> δι' δ καὶ χώραν ζητεῦ

³ Plin. 18. 120; Varro 1, 23, 3; Col. 2, 10 ⁴ 8, 7, 2,

¹ cf. C.P. 4. 12. 1. ² cf. C.P. 4. 8. 3. ³ Plin. 18. 120; Varro 1. 23. 3; Col. 2. 10. 7.

⁵ i.e. dig in the bean-plants if the soil is poor, before the pools are formed enough to make it worth while to gather the beans. So Varro l.c. ⁶ Cited by Galen.

ENQUIRY INTO PLANTS, VIII. viii. 7-1x. 2

aspect and shew no difference of soil, some bear 'cookable' some 'uncookable' seeds, and that sometimes when there is only 1 the breadth of a furrow between them.

Of the grains and pulses which most exhaust the soil, or which improve it.

IX. Wheat exhausts the land more than any other crop, and next to it barley; wherefore the former requires good soil, while barley will bear even on somewhat crumbling soils; ² and of leguminous plants chick-pea is the most exhausting, although this crop is in the ground only a very short time. ³ Beans, as was said, ⁴ are in other ways not a burdensome crop to the ground, they even seem to manure it, because the plant is of loose growth and rots easily; wherefore the people of Macedonia and Thessaly turn over the ground when it is in flower. ⁵

Gof the plants which resemble wheat or barley—such as zeia (rice-wheat) one-seeded wheat olyra (rice-wheat) oats aigilops—zeia is the strongest and most exhausts the ground; for it has many roots which run deep and many stems; but its fruit is the lightest and is welcome to all animals. Of the rest oats is the most exhausting; for this too has many roots and many stems. Olyra is a more delicate plant and not so robust as these. But one-seeded wheat is the crop which is of all the least burdensome to the soil; for it has but a single slender stem 10; wherefore also it requires a light soil and not, like

10 και λεπτοκάλαμον add. Bod. from Galen.

⁷ See Index.

⁸ Ισχυρότατον conj. W. from Galen; Ισχυρότερον Ald.

⁹ βρόμος· πολύρριζος γάρ conj. Sch.; β. πολ.· καὶ γάρ Ald.

λεπτήν, οὐχ ὥσπερ ή ζειὰ πίειραν καὶ ἀγαθήν. έστι δὲ δύο ταῦτα καὶ ὁμοιότατα τοῖς πυροῖς ή τε <ζειὰ καὶ ἡ τίφη,> ὁ δ' αἰγίλωψ καὶ ὁ Βρόμος ωσπερ άγρι' άττα καὶ ἀνήμερα.

Έπικαρπίζεται δὲ σφόδρα καὶ ὁ αἰγίλωψ τὴν γην, καί έστι πολύρριζον και πολυκάλαμον ή δε αίρα παντελώς άπηγριωμένον. των δε έν τοίς θερινοίς αρότοις τὸ σήσαμον δοκεί γαλεπώτατον είναι τη γη καὶ μάλιστα καρπίζεσθαι καίτοι πολυκαλαμώτερον καὶ παχυκαλαμώτερον καὶ πολυρριζότερον κέγχρος. διαφέρει δὲ τά τε πρὸς την γην κουφα καὶ τὰ πρὸς την ημετέραν τροφήν. ένια γαρ έναντίως, ώσπερ τα γεδροπα και οί κέγχροι καὶ τὰ πρὸς ἡμᾶς δέ, ὥσπερ ἐλέχθη, καὶ τἆλλα ζῶα. καὶ περὶ μὲν τούτων ἄλις.

Χ. Νοσήματα δὲ τῶν σπερμάτων τὰ μὲν κοινὰ πάντων έστίν, οίον ή έρυσίβη, τὰ δ' ἴδιά τινων. οίον ό σφακελισμός του έρεβίνθου, και το ύπο καμπών κατεσθίεσθαι καὶ ὑπὸ ψυλλών, τινὰ δὲ καὶ ὑπ' ἄλλων θηριδίων. ἔνια δὲ καὶ ψωριᾶ καὶ άλμα, καθάπερ καὶ τὸ κύμινον. τὰ δ' ἐπιγινόμενα ζωα μη έξ αὐτων άλλ' έκ των έξωθεν οὐχ ομοίως βλάπτει. ἐπιγίνεται γὰρ ἡ μὲν κανθαρίς

¹ ή τε ζειά καὶ ή τίφη add. W. from Galen.

² δδ' conj. Scal.; δτ' Ald. H.; ητ' UMP. 8 καίτοι conj. W.; καὶ Ald. cf. C.P. 4. 15. 1.
4 τὰ add. St.
5 8. 3. 5 ad fin.

⁶ καὶ τὰ Ald.: καὶ οἱ τὰ UMP: ? καὶ αὖ τὰ W. .

ENQUIRY INTO PLANTS, VIII. 1x. 2-x. 1

zeia, one that is fat and good. These last two,¹ zeia and one-seeded wheat, are also those which are likest to wheat, while ² aigilops and oats are as it

were wild and uncultivated things.

Aigilops also greatly exhausts the land, having many roots and many stems; while darnel is a plant which has become altogether wild. Of the crops sown at the summer seed-time sesame seems to be most severe on the land and to exhaust it most; yet 3 millet has more numerous and stouter stems and more roots. Moreover there is a difference between crops which 4 are called 'light' in relation to the soil and those called 'light' in regard to human use. For some, such as leguminous plants and millet, are light in one sense but not in the other; and, as was said, 5 what 6 is light for men is not necessarily so for the other animals. Now enough of these matters.

Of the diseases of cereals and pulses, and of hurtful winds.

X. ⁷ As to diseases of seeds—some are common to all, as rust, some are peculiar to certain kinds; thus chick-pea is alone subject to rot⁸ and to being eaten by caterpillars and by spiders⁹; and some seeds are eaten ¹⁰ by other small creatures. Some again are liable to canker and mildew, ¹¹ as cummin. But creatures which do not come from the plant itself but from without do not do so much harm; thus the kantharis ¹² is a visitor among wheat, the

Digitized by Google

Plin. 18. 152 and 154.
 ψυλλῶν: described by Arist. H.A. 9. 39. 1.

¹⁰ δè add. Sch.; ? κατεσθίεσθαι· κατεσθίεται δè καὶ ὑπὸ ψ. W.

ψωριᾶ καὶ άλμᾶ conj. W.; ψώραις καὶ ἄλμαις Ald. cf. 7.5.4 n.
 Plin. 18. 156.

έν τοῖς πυροῖς, τὸ δὲ φαλάγγιον ἐν ὀρόβοις, ἄλλα δ' ἐν ἄλλοις.

- 'Ερυσιβά δ' ώς άπλως είπειν τὰ σιτώδη μάλλον τῶν ὀσπρίων αὐτῶν δὲ τούτων κριθὴ μᾶλλον η πυρός και των κριθών έτεραι ετέρων, μάλιστα δ' ώς είπειν ή 'Αχιλληίς. διαφέρει δε και ή των χωρίων θέσις καὶ ή φύσις οὐ μικρόν τὰ γὰρ προσήνεμα καὶ μετέωρα οὐκ ἐρυσιβᾶ ἡ ἡττον, άλλα τα έγκοιλα και άπνοα γίνεται δε ή έρυσίβη 8 πανσελήνοις μάλιστα. ἀπόλλυται δὲ καὶ ὑπὸ των πνευμάτων καὶ πυρὸς καὶ κριθή, ὅταν ἡ άνθοῦντα ληφθή ή άρτι ἀπηνθηκότα καὶ ἀσθενή. μαλλον δε κριθή, πολλάκις δ' ήδη εν τω άδρύνεσθαι οὖσα, ἐὰν μεγάλα καὶ πλείω χρόνον ἐπιγένηται Εηραίνει γαρ και άφαυαίνει, δ καλουσί τινες έξανεμοῦσθαι. διαπόλλυσι δὲ καὶ ήλιος ό έκνέφελος ἄμφω καὶ μᾶλλον πυρὸν ἡ κριθήν, ὥστε μηδ' ἐπίδηλον είναι τὸν στάχυν τῆ ὄψει ὄντα κενόν.
 - Τὸν δὲ πυρὸν ἀπολλύουσι καὶ οἱ σκώληκες οἱ μὲν εὐθὺς κατεσθίοντες φυόμενοι τὰς ῥίζας, οἱ δὲ ὅταν αὐχμῶντες ἀποχυθῆναι μὴ δύνωνται· τότε γὰρ ἐγγινόμενος ὁ σκώληξ ἐσθίει τὸν ἀποπηνιζόμενον κάλαμον· ἐσθίει δὲ ἄχρι τοῦ στάχυος, εἶτ'

¹ Plin. 18. 154.

ξρυσιβά conj. W.; ἐρυσίβαι Ald.; els add. Sch.
 τὰ add. Sch.
 τζ. C. P. 3. 22. 2.

⁵ έρυσιβα conj. Sch.; έρυσίβαι Ald. 6 cf. C.P. 4. 13. 4; Plin. 18. 151.

⁷ μέγαλα conj. Sch.; μεγάλη UMAld.

ENQUIRY INTO PLANTS, VIII. x. 1-4

phalangion in vetches, and other pests in other

crops.

¹Generally speaking, cereals are more liable to rust 2 than pulses, and among these barley is more liable to it than wheat; while of barleys some kinds are more liable than others, and most of all, it may be said, the kind called 'Achillean.' Moreover the position and character of the land make no small difference in this respect; for lands which 8 are exposed to the wind and elevated are not liable to rust,5 or less so, while those that lie low and are not exposed to wind are more so. And rust occurs chiefly at the full moon. wheat and barley are destroyed by winds, if they are caught by them either when in flower, or when the flower has just fallen and they are weak; and this applies specially to barley, indeed it occurs when the grain is already ripening, if the winds are violent and last a long time; for they dry up and parch the grain, which some call being 'wind-bitten.' Also a hot sun after cloudy weather destroys both, and wheat more than barley, so that the ear is not even conspicuous, since it is emptv.

Wheat is also destroyed by grubs; sometimes they eat the roots, as soon as they appear,⁸ sometimes they do their work when by reason of drought the ear cannot be formed ⁹; for at such times the grub is engendered, and eats the haulm as it is becoming unrolled ¹⁰; it eats right up to the ear and then,

10 ἀποπηνιζόμενον: lit. 'unwinding itself.' All edd. mark the word as corrupt.

 ⁸ φυόμενοι conj. Sch.; φυόμενον Ald. cf. C.P. 3. 22. 4.
 ⁹ ἀποχυθῆναι conj. Sch. after Vin. Vo.G; ἀπολυθῆναι UM Ald. cf. C.P. 3. 22. 4; 4. 14. 1.

έξαναλώσας ἀπόλλυται· καὶ ἐὰν μὲν ὅλον ἐκφάγη τελέως αὐτὸς ὁ πυρός, ἐὰν δὲ ἐπὶ θάτερον τοῦ καλάμου καὶ ἐκβιάσηται τὴν ἀπόχυσιν, τοῦτο μὲν αὖον τοῦ στάχυος θάτερον δὲ ὑγιές. γίνεται δὲ οὐ πανταχοῦ τὸ περὶ τοὺς πυρούς, οἶον ἐν Θετταλίᾳ, ἀλλὰ κατὰ χώρας τινάς, ὥσπερ ἐν τῆ Λιβύη καὶ τῆς Εὐβοίας ἐν τῷ Ληλάντφ.

Σκώληκες δὲ γίνονται καὶ ἐν τοῖς ἄχροις καὶ τοῖς λαθύροις καὶ τοῖς πισοῖς, ὅταν ὑγρανθῶσι καὶ θερμημερίαι γένωνται, καθάπερ καὶ ἐν τοῖς ἐρεβίνθοις αἰ κάμπαι. πάντα δὲ ἐξαναλώσαντα τὰς τροφὰς ἀπόλλυται καὶ ἐν τοῖς χλωροῖς καὶ ἐν τοῖς ξηροῖς καρποῖς, οἶον οἴ τε ἶπες καὶ οἱ ἐν τοῖς κυάμοις ἐγγινόμενοι καὶ ἐν τοῖς ἄλλοις, ὥσπερ καὶ ἐν τοῖς δένδρεσι καὶ ἐν τοῖς ξύλοις ἐλέχθη, πλὴν τῶν κεραστῶν καλουμένων. πρὸς ἄπαντα δὴ ταῦτα μεγάλα διαφέρουσιν αἱ χῶραι οὐκ ἀλόγως ὁ γὰρ ἀὴρ εὐθὺς διάφορος τῷ θερμὸς ἡ ψυχρὸς εἶναι ἡ ὑγρὸς ἡ ξηρός οῦτος δ' ἦν ὁ γονεύων δι' δ καὶ ἐν οῖς εἰώθασι γίνεσθαι οὐκ ἀεὶ γίνονται.

ΧΙ. Τῶν δὲ σπερμάτων οὐχ ἡ αὐτὴ δύναμίς ἐστιν εἴς τε τὴν βλάστησιν καὶ εἰς θησαυρισμόν.

³ cf. C P. 3. 22. 3. ωχροις conj. St.; σχροις Ald. H.

4 καθάπερ και conj. Sch.; και καθάπερ Ald.

⁵ πάντα conj. W.; τὰ Ald.

¹ αύτδς: sc. the grain. ἀναίνεται conj. W.; ? αὐαίνεται αὐτδς. 2 θάτερον conj. Sch.; θατέρου Ald.

ENQUIRY INTO PLANTS, VIII. x. 4-xi. 1

having consumed it, perishes. And, if it has entirely eaten it, the wheat itself perishes; if however it has only eaten one side of the haulm and the plant has succeeded in forming the ear, half the ear withers away, but the other half remains sound. However it is not everywhere that the wheat is so affected; for instance this does not occur in Thessaly, but only in certain regions, as in Libya and at Lelanton in Euboea.

Grubs occur also in okhros³ lathuros and peas. whenever these crops get too much rain and then hot weather supervenes; and caterpillars occur in chick-peas under the same conditions.4 All 5 these pests perish, when they have exhausted their food, whether the fruit in which they occur be green or dry, just as wood-worms do and the grubs found in beans and other plants, as was said of the pests found in growing trees and in felled timber. But the creature called 'horned worm' 6 is an exception. Now in regard to all these pests the position makes a great difference, as might be expected. climate, it need hardly be said, makes a difference according as it is hot or cold, moist or dry; and it was the climate which gave rise to these pests 7; wherefore they are not always found even in places in which they ordinarily occur.8

Of seeds which keep or do not keep well.

XI. The seeds have not all the same capacity for germination and for keeping well. Some germinate

⁶ cf. 4. 14. 5; C.P. 5. 10. 5.

⁷ δ' ήν δ γονεύων Ι conj.; δ' ήν δ νεύων UAld.; δ' ήνονεύων M; δ' έστιν δ γονεύων conj. Sch.; δ' δ γονεύων conj. W.

⁸ i.e. because the atmospheric conditions are not always favourable to the pest.

ένια μεν γὰρ βλαστάνει καὶ τελειοῦται τάχιστα καὶ θησαυρίζεται κράτιστα, καθάπερ ἔλυμος καὶ κέγχρος ἔνια δὲ βλαστάνει μὲν εὖ ταχέως δὲ σήπεται, καθάπερ ὁ κύαμος καὶ μᾶλλον ὁ τεράμον ταχὺ δ' ἡ ἀφάκη καὶ ὁ δόλιχος κριθὴ δὲ πυροῦ <θᾶττον> θᾶττον δὲ καὶ ὁ κονιορτώδης σῖτος καὶ ὁ ἐν οἰκήμασι κονιατοῖς ἡ ἀκονιάτοις.

Γίνεται δὴ φθειρομένοις σπέρμασι ἴδια ζῶα, καθάπερ ἐλέχθη, πλὴν ἐρεβίνθου· μόνος γὰρ οὖτος οὐ ζωογονεῖ. καὶ σηπομένοις μὲν πᾶσι σκώληξ κοπτομένοις δὲ καθ' ἔκαστον ἴδιον. πάντων δὲ μάλιστα διαμένουσιν ἐρέβινθος καὶ ὅροβος, τούτων δ' ἔτι μᾶλλον ὁ θέρμος· ἀλλ' ἔοικέ γ' οὖτος

ὥσπερ ἀγρίω.

Διαφέρει δὲ ὡς ἔοικε χώρα χώρας καὶ ἀὴρ ἀέρος εἰς τὸ κόπτεσθαι καὶ μὴ τὰ σπέρματα ἐν ᾿Απολλωνία γοῦν τῆ περὶ τὸν Ἰόνιον οὐκ ἐσθίεσθαί φασιν ὅλως κύαμον, δι ὁ καὶ εἰς θησαυρισμὸν ἀποτίθεσθαι διαμένει δὲ καὶ περὶ Κυζικὸν ἐπὶ πλείω. μέγα δὲ πρὸς διαμονὴν καὶ τὸ ξηρὰ θερίζειν ἐλάττων γὰρ ἡ ὑγρότης θερίζουσι δ' ἐγχυλότερα τὰ μὲν χεδροπὰ πρὸς τὸ μᾶλλον καὶ ρῷον συλλέξαι, ταχὸ γὰρ καταρρεῖ καὶ αὐανθέντα

⁵ καὶ ὁ κον.... ἀκονιάτοις conj. W., cf. C.P. 4. 16. 1; ὁ κονιορτώδης καὶ ὁ κονίορτος· καὶ ὁ ἐν τοῖς κονιορτοῖς ἐν ἄπασιν, οῖον κονιάτοις ἡ ἀκονιάτοις Ald.; so also UM, but omitting τοῖς; Ü gives κονιατοῖς; ἡ ὁ ἀκονίορθος for καὶ ὁ κονίορτος mBas. cf. Plin. 18. 301, Varro 1. 57. 1, where the use of a cement of pounded marble is recommended.

6 δη φθειρομένοις conj. Sch.; διαφθειρομένοις UMAld.

¹ εδ conj. W.; οὐ Ald. 2 σήπεται add. W. 3 cf. 8. 8. 6; C. P. 5. 18. 2. 4 θᾶττον add. W.

ENQUIRY INTO PLANTS, VIII. xi. 1-3

and mature very quickly, and keep excellently, as Italian millet and millet. Some germinate well, but soon rot, as beans, and especially those that are cookable; so do tare and calavance; also barley perishes sooner than wheat; and dusty grain and that which is kept in plastered store-rooms perishes sooner than that which is kept in unplastered rooms.

Again, as seeds decay, they engender special creatures, except chick-pea, which alone engenders none. As they rot, all produce a grub; but, as they get worm-eaten, each produces a special creature. Chick-pea and vetch keep best of all, and better still than these lupin; but this, as it were, is like a wild kind.

⁹ It appears that soil and climate make a difference as to whether the seed gets worm-eaten or not; at least they say that at Apollonia on the Ionian Sea beans do not get eaten in this way at all, and therefore they are put away and stored; and about Cyzicus they keep an even longer time. It also makes a great difference to keeping that the seed should be gathered dry, for then there is less moisture in it.¹⁰ However the seeds of leguminous plants are gathered with a certain amount of moisture in them,¹¹ because then they can be collected in greater quantity and more easily; for otherwise they are soon shed and get shrivelled up and split ¹²;

8 i.e. and so the seed is hard and not liable to these attacks.

cf. 8. 11. 8; C.P. 4. 16. 2.

of. C.P. 4. 16. 2. 10 i.e. liability to rot.

 $^{^{7}}$ i.e. rot is produced in all cases by the same creature (σκώληξ), but the condition called being 'worm-eaten' is due in each plant to a different pest.

¹¹ έγχυλότερα conj. Sch.; εὐχυλότερα Ald. H. Cam.; εὐχηλότερα Bas. cf. C.P. 4. 13. 3.

θρύπτεται, τοὺς δὲ πυροὺς καὶ γένος τι κριθῶν διά τὸ βελτίους εἰς τὰ ἄλφιτα γίνεσθαι μὴ ἀπεξηραμμένας.

- Δι' δ καὶ εἰς θώμους συντιθέασι καὶ πυρούς καὶ κοιθάς, καὶ δοκοῦσιν άδρύνεσθαι ἐν θώμω μᾶλλον ή λιποσαρκείν, οὐκ ἐσθίεται δὲ σῖτος, ὅταν ὑσθεὶς θερισθή άθέριστος δὲ μάλιστα διαμένει ὁ πυρός, έτι δὲ μᾶλλον ὁ θέρμος οὐδὲ γὰρ θερίζουσι τοῦτον πρότερον η ύδωρ γενέσθαι, δια το έκπηδαν θεριζόμενον καὶ ἀπόλλυσθαι τὸ σπέρμα.
- Πρὸς ἔκφυσιν δὲ καὶ τὴν ὅλην σπορὰν ἄριστα δοκεί τὰ ένάενα· τὰ δὲ δίενα γείρω καὶ τὰ τρίενα. τὰ δ' ὑπερτείνοντα σχεδὸν ἄγονα, πρὸς δὲ τὴν σίτησιν άρκοθντα. βίος γάρ έστιν έκάστοις ώρισμένος είς γονήν. καίτοι καὶ ταύταις παραλλάττει ταις δυνάμεσι δια τούς τόπους έν οίς αν θησαυρίζωνται, της γοῦν Καππαδοκίας ἐν χωρίω τινὶ τῷ καλουμένω Πέτρα καὶ τετταράκοντα ἔτη διαμένειν φασί γόνιμα καί χρήσιμα πρὸς σπόρον, είς δὲ τὴν σίτησιν ἐξήκοντα ἡ ἐβδομήκοντα· τὸ γὰρ όλον οὐ κόπτεσθαι· τὰ δὲ ἱμάτια καὶ τὴν ἄλλην ε γάζαν κόπτεσθαι. τὸ γὰρ χώριον ἄλλως τε ύψηλον είναι καὶ εὔπνουν καὶ ἔναυρον αἰεὶ καὶ άπ' ἀνατολης ἔχουσι καὶ δύσεως καὶ μεσημβρίας.

θερισθή conj. W.; περιφυή MSS.

¹ ή λιποσαρκείν conj. Η.; ήλίκα σωρών U; ήλίκα σαρκών Μ. W. brackets as due to a gloss. cf. C.P. 4. 13. 6.
² δταν δσθεις conj. Scal.: so Vo.; δ τανυσθείς Bas.Cam.

ENQUIRY INTO PLANTS, VIII. x1. 3-6

and wheat and one kind of barley are gathered before they are dry, because then they are better for meal.

Wherefore the grain of wheat and barley is put into heaps, and it seems to ripen in a heap rather than to lose substance.¹ (However corn does not get worm-eaten when it is reaped after exposure to rain.)² Also corn lasts better than other things if it is left standing, and so does lupin to an even greater extent; indeed this crop is not even gathered till rain has fallen,³ because, if it is gathered, the seed springs out and is lost.

Of the age at which seeds should be sown.

⁴For propagation and sowing generally seeds one year old seem to be the best; ⁵ those two or three years old are inferior, while those kept a still longer time are infertile, though they are still available as food. For each kind has a definite period of life in regard to reproduction. However these seeds too differ in their capacity according to the place in which they are stored. For instance, in Cappadocia at a place called Petra they say that seed remains even for forty years fertile and fit for sowing, while as food it is available for sixty or seventy years; for that it does not get wormeaten at all like clothes and other stored-up articles: for that the region is, apart from this, elevated and always exposed to fair winds and breezes which prevail alike from ⁶ the east, the west, and the

209

Vol., II.

þ

³ πρότερον ή conj. W.; τον τρόπον UAld. cf. C.P. 4. 13. 3; Plin. 18. 133. ⁴ Plin. 18. 195.

cf. 7. 5. 5; Geop. 2. 16.
 ἀπ' conj. Sch.; ἐπὶ P₂Ald.

φασί δὲ καὶ ἐν Μηδεία καὶ ταῖς ἄλλαις ταῖς ὑψηλαῖς χώραις διαμένειν θησαυριζόμενα πολὺν χρόνον. ἐρέβινθον δὲ δὴ καὶ θέρμον καὶ ὄροβον καὶ κέγχρον καὶ τὰ τοιαῦτα δῆλον ὅτι πολλῷ πλείω τούτων, ὥσπερ καὶ ἐν τοῖς περὶ τὴν Ἑλλάδα τόποις. ἀλλὰ ταῦτα μέν, ὥσπερ εἴρηται, τῶν τόπων ἴδια.

Δοκεί δὲ καὶ γῆ τις εἶναι παρά τισιν ἡ διαπαττομένη συντηρεί τὸν πυρόν, ὥσπερ ἡ τε ἐν 'Ολύνθω καὶ ἐν Κηρίνθω τῆς Εὐβοίας· ποιεί δὲ χείρω μὲν εἰς τὴν σίτησιν ἀδρότερον δὲ τῆ προσόψει· παραπάττουσι δὲ χοίνικα εἰς τὸν μέδιμνον.

Πυρωθέντα πάντα τὰ σπέρματα ἀπόλλυται καὶ ἀβλαστῆ γίνεται· καίτοι περί γε Βαβυλῶνά φασι τὰς κριθὰς καὶ τοὺς πυροὺς ἐπὶ τῆς ἄλω πηδᾶν, ὥσπερ τὰ φρυγόμενα· ἀλλὰ δῆλον ὅτι διαφορά τίς ἐστι τῆς θερμότητος, ἡ ἀπλῶς πως θερμασία γίνεται [καὶ] ἡ πήδησις. καὶ τὰ μὲν τοιαῦτα σχεδὸν ὡσπερεὶ κοινὰ δόξειεν ἄν εἶναι πάντων ἡ τῶν πλείστων.

² διαπαττομένη conj. Η.; διαπλαττομένη UMAld.; διακοπτομένη P₂. cf. Plin. 18. 305.

³ παραπάττουσι conj. Sch., cf. Geop. 2. 21. 3; (ἐμπάσσειν); παραπάττουσι UMAld. cf. Varro 1. 57. 1.

 $^{^{1}}$ cf. C.P. 5. 18. 3; for millet-seed see J.H.S. vol. xxxv. part i. p. 22.

ENQUIRY INTO PLANTS, VIII. xi. 6-7

south. They say that in Media 1 also and other elevated countries the seed when stored keeps for a long time. And it is plain that chick-pea lupin vetch millet and the like will keep a far longer time than these seeds, as they do even in districts of Hellas. However these peculiarities, as has been said, are due to the particular region.

Of artificial means of preserving seed.

There appears to be a kind of earth in some places, which when sprinkled ² over the seed helps to make wheat keep, for instance, the earth found at Olynthos and at Kerinthos in Euboea; this makes the grain inferior for food, but fuller in appearance; the earth is sprinkled ³ in the proportion of one pint to twenty-four of grain.

Of the effect of heat on seeds.

All seeds if exposed to fire perish and become infertile. Yet they say that at Babylon 4 the grains of barley and wheat jump on the threshing-floor like corn which is being parched. However it is plain that it is some particular kind of warmth 5 which produces this effect: or else the jumping is simply another effect of heat. 6 Such behaviour would appear to be common to most, 7 if not to all kinds.

5 i.e. the sun's heat is different in kind, and therefore in effect, to that of a fire.

6 θερμασία conj. Sch.; θερμασία Ald. H.

2 I I

⁴ i.e. the grain is there exposed to great sun-heat. cf. de igne 44.

ο ωσπερεί κοινά conj. Sch. from G; ωσπερ εἰκόνα UM; ωσπερ εἰκόνε Ald. Η.

- Ένια δὲ ἔγει τινὰ ἰδιότητα καὶ τῶν δοκούντων ώσπερ αγρίων είναι καὶ κατὰ τὴν γένεσιν καὶ τὴν έκφυσιν, ώσπερ ο θέρμος καὶ ο αἰγίλων ο μεν γαρ θέρμος καίπερ ισχυρότατος ων όμως, εαν μή εὐθὺς ἀπὸ τῆς ἄλω καταβληθῆ, κακοφυὴς γίνεται, καθάπερ έλέχθη, καὶ τὸ ὅλον δὲ οὐκ ἐθέλει κρύπτεσθαι τη γη, δι' δ και ούχ ύπαρουντες σπείρουσι πολλάκις δε καν είς ύλην ή βοτάνην τινα πέση, διωσάμενος ταύτην συνάπτει την δίζαν τη γη και βλαστάνει. χώραν δὲ ὕφαμμον ζητεῖ καὶ φαύλην μᾶλλον, τὸ δὲ ὅλον οὐκ ἐθέλει φύεσθαι έν διειργασμένη.
 - Ο δὲ αἰγίλωψ ἀνάπαλιν ἐν γὰρ τῆ γεωργουμένη κάλλιον καὶ ένιαχοῦ δὲ πρότερον άβλαστής ων έαν γεωργηθη βλαστάνει και γίνεται πολύς, καὶ όλως δὲ φιλεῖ χώραν ἀγαθήν. ἴδιον δὲ αὐτοῦ λέγεται πρὸς τὰ ἄλλα τὰ σιτώδη σπέρματα καὶ ή παρ' ενιαυτον βλάστησις εκατέρου των σπερμάτων. δι' δ καὶ οἱ βουλόμενοι τελέως φθείραι, δύσφθαρτον γὰρ δὴ φύσει τυγχάνει, τὰς ἀρούρας ανιασιν ασπόρους έπι δύο έτη, και όταν αναβλαστήση τὰ πρόβατα ἐπαφιᾶσι πολλάκις, ἔως αν εκνεμηθώσι, και αυτη γίνεται φθορά παντελής άμα δὲ τοῦτο μαρτυρεί καὶ τὴν παρὰ μέρος βλάστησιν.

¹ καl conj. Sch.; ħ Ald. H.

cf. 8. 1. 3. β cf. C.P. 4. 7. 3. δπαροῦντες conj. Η.; δπαποροῦντες UMAld. cf. C.P. 3. 20. 8.

ENQUIRY INTO PLANTS, VIII. xi. 8-9

Of certain peculiarities of the seed of lupin and aigilops.

Some even 1 of those kinds which seem to be more or less wild have peculiarities as to their germination and growth, for instance, lupin and aigilops. For lupin, although it is very robust, unless it is planted immediately after leaving the threshing-floor, 2 turns out of poor growth, as was said, and refuses altogether to be buried in the ground; 3 wherefore they sow it without first ploughing 4 the land. And often if the seed has fallen amid thick undergrowth or herbage, 5 it thrusts this aside, fastens on to the earth with its root and grows vigorously. It seeks sandy and poor soil for choice, and will not grow at all in 6 cultivated 7 soil.

Aigilops has the opposite character; it grows better in tilled soil; and in some places where at first it would not grow, if the ground is tilled, it grows and yields a large crop, and in general it likes good soil. A peculiarity mentioned in regard to it as compared with other cereal seeds is that one seed in two does not germinate for a year. Wherefore those who wish to destroy it entirely, (since it is naturally hard to destroy), leave the fields unsown for two years, and, when it springs up, send in the sheep several times till they have grazed it down, and this is a way of completely destroying it. At the same time this testifies to the fact that the seed does not all germinate at once.

⁹ ἐπαφιᾶσι conj. Sch., cf. 8. 7. 4; ἀφιῆσι M; ἀφίησι P; ἀφίασι Ald,

BOOK 1X

Ι. Ἡ ὑγρότης οἰκεία τῶν φυτῶν, ἢν δὴ καλοῦσί τινες ὀπὸν ὀνόματι κοινῷ προσαγορεύοντες. δύναμιν δὲ ἔχει δῆλον ὅτι τὴν καθ' αὐτὴν ἐκάστη. χυμὸς δὲ ταῖς μὲν μᾶλλον ταῖς δ' ἢττον ἀκολουθεῖ, ταῖς δ' ὅλως οἰκ ἀν δόξειεν, οὕτως ἀσθενὴς καὶ ὑδαρής τίς ἐστι. πλείστη μὲν οὖν ὑπάρχει πᾶσι κατὰ τὴν βλάστησιν, ἰσχυροτάτη δὲ καὶ μάλιστα ἐκφαίνουσα τὴν ἑαυτῆς φύσιν ὅταν ἤδη παύσηται καὶ βλαστάνοντα καὶ καρπογονοῦντα. συμβαίνει δὲ τισι τῶν φυτῶν καὶ χρόας ἰδίας ἔχειν. τοῖς μὲν λευκὰς οἶον τοῖς ὀπώδεσι, τοῖς δ' αἰματώδεις οἷον τῷ κενταυρίω καὶ τῆ ἀτρακτυλίδι καλουμένη ἀκάνθη, τοῖς δὲ χλωρόν, τοῖς δ' ἐν ἄλλη χρόα, ἔνδηλα δὲ μᾶλλον ταῦτα ἐν τοῖς ἐπετείοις καὶ διὶς ἐπετείοις καὶ διὶς ἐπετείοις ἐπετεί

'Η δ' ὑγρότης τῶν μὲν πάχος ἔχει μόνον, ὥσπερ τῶν ὀπωδῶν· τῶν δὲ καὶ δακρυώδης γίνεται, καθάπερ ἐλάτης πεύκης τερεβίνθου πίτυος ἀμυγδαλῆς κεράσου προύμνης ἀρκεύθου κέδρου τῆς ἀκάνθης τῆς Αἰγυπτίας πτελέας, καὶ γὰρ αὕτη φέρει κόμμι

¹ cf. C.P. 6. 11. 16.

² I have omitted ή and restored δὲ before ἔχει (om. Scal.; found in UMAld.).

³ τῷ κενταυρίφ conj. Scal. cf. Plin. 25. 32; κενταυρίδι conj. St.; κευτηρία P₂Ald.G, cf. 9. 8. 7.

BOOK IX

OF THE JUICES OF PLANTS, AND OF THE MEDICINAL PROPERTIES OF HERBS.

Of the various kinds of plant-juices and the methods of collecting them.

I. 1 Moisture belongs to plants as such and some call it the 'sap,' to give it a general name; and it plainly has 2 special qualities in each plant. This moisture is attended by a taste, in some cases more, in some less, while in some it would seem to have none, so weak and watery is it. Now all plants have most moisture at the time of making growth, but it is strongest and most shows its character when the plant has ceased to grow and to bear fruit. Again in some plants the juice has a special colour; in some it is white, as in those which have a milky juice; in some blood-red, as in centaury 3 and the spinous plant which is called distaff-thistle; in some green: and in some of other colours. And these qualities are more obvious in annual 4 plants and those with annual stems than in trees.

Again in some plants the juice is merely thick, as in those in which it is of milky character; but in some it is of gummy character, as in silver-fir fir terebinth Aleppo pine almond kerasos (bird-cherry) bullace Phoenician cedar prickly cedar acacia elm.⁵ For

ἐν inserted here by W. instead of before τοις ἐπετείοις.
 πτελέας after κέδρου P₂Ald.; transposed by Sch. after Tobias Aldinus. cf. Plin. 13, 67.

πλην οὐκ ἐκ τοῦ φλοιοῦ ἀλλ' ἐν τῷ κωρύκῳ, ἔτι δὲ ἀφ' ὧν ὁ λίβανος καὶ ἡ σμύρνα, δάκρυα γὰρ καὶ ταῦτα, καὶ τὸ βάλσαμον καὶ <ή> χαλβάνη καὶ εἴ τι τοιοῦτον ἔτερον, οἶόν φασι τὴν ἄκανθαν τὴν Ἰνδικήν, ἀφ' ἢς γίνεταί τι ὅμοιον τῆ σμύρνης συνίσταται δὲ καὶ ἐπὶ τῆς σχίνου καὶ ἐπὶ τῆς ἀκάνθης τῆς ἰξίνης καλουμένης, ἐξ ὧν ἡ μαστίχη.

καὶ τητα δὲ ταῦτα εὕοσμα καὶ σχεδὸν ὅσα πιότητά τινα ἔχει καὶ λῖπος ὅσα δ' ἀλιπῆ ταῦτα δ' ἄοσμα, καθάπερ τὸ κόμμι καὶ τὸ τῆς ἀμυγδαλῆς. ἔχει δὲ δάκρυον καὶ ἡ ἰξία ἡ ἐν Κρήτη καὶ ἡ τραγάκανθα καλουμένη ταύτην δὲ πρότερον ῷοντο μόνον ἐν Κρήτη φύεσθαι, νῦν δὲ φανερὰ καὶ ἐν ᾿Αχαίδι τῆς Πελοποννήσου καὶ ἄλλοθι καὶ τῆς ᾿Ασίας περὶ τὴν Μήδειαν. καὶ ἄλλοθι καὶ τῆς ᾿Ασίας περὶ τὴν Μήδειαν.

τούτων μèν πάντων ἔν τε τοῖς καυλοῖς καὶ τοῖς στελέχεσι καὶ τοῖς ἀκρεμόσι τὸ δάκρυον ἐνίων δ' ἐν ταῖς ρίζαις, ὥσπερ τοῦ ἱπποσελίνου καὶ τῆς σκαμμωνίας καὶ ἄλλων πολλῶν φαρμακωδῶν. τῶν δὲ καὶ ἐν τῷ καυλῷ καὶ ἐν τῷ ρίζη καὶ γὰρ τὸν καυλὸν ὀπίζουσιν ἐνίων καὶ τὰς ρίζας, ὥσπερ καὶ τοῦ σιλφίου.

Τὸ μὲν οὖν τοῦ ἱπποσελίνου παρόμοιον τῆ σμύρνη καί τινες ἀκούσαντες ὡς ἐντεῦθεν ἡ σμύρνα ἡγοῦνται βλαστάνειν ἐξ αὐτῆς ἱπποσέ-

¹ κωρύκφ conj. Sch.; ἀγγείφ Η.; ἀγείφ P₂Ald. probably a gloss on κωρύκφ, for which of. 2. 8. 3 and reff. in note. Plin. l.c. has preserved the right word through an absurd blunder—in Coryco monte Ciliciae.

ENQUIRY INTO PLANTS, IX. 1, 2-4

this last also produces a gum, though it does not exude from the bark, but is found in the 'bag'1 of the leaves: there are also the juices from which come frankincense and myrrh; for these too are gums; so too are balsam of Mecca khalbane 2 and any others of the kind that there may be, such as, they say, the Indian akantha, from which comes something 3 resembling myrrh; and a similar substance forms on mastich and the spinous plant called ixine

(pine-thistle), whence mastic-gum is made.

All these have a fragrant odour, as in general have those which contain a viscous substance and are fatty: while those that are not fatty have no scent. as gum and the juice which exudes from the almond. The pine-thistle 4 of Crete has also a gum, and so has the plant called tragacanth; 5 this was formerly supposed to grow only in Crete, but now it is well known to grow also in Achaia in the Peloponnese and elsewhere in Hellas and in Asia in the Median country. In all these plants the gum occurs in the stems the trunks and the branches, but some plants it is found in the roots, as in alexanders scammony and many other medicinal plants. In some it is found in the stem and also in the root:6 for of some 7 plants they tap the stem and the roots as well, as is done with silphium.

Now the juice of alexanders is like myrrh, and some, having heard that myrrh comes from it, have supposed that, if myrrh is sown, alexanders comes up

^{**} galbanum. cf. Plin. 12. 121; 24. 21. Verg. G. 3. 415; 4. 264. See 9. 7. 2; 9. 9. 2 n. ** τ_1 I conj.; τ_2 MSS. ** $\xi(\alpha = i\xi(\nu_1)$. See Index. ** Plin. 13. 115. ** cf. C.P. 6. 11. 15.

⁷ ένίων και conj. Sch.; και ένίων Ald.

λινον· φυτεύεται γάρ, ὥσπερ ἐλέχθη, καὶ ἀπὸ δακρύου τὸ ἱπποσέλινον, καθάπερ ἡ κρινωνία καὶ ἄλλα. τὸ δὲ τοῦ σιλφίου δριμύ, καθάπερ αὐτὸ τὸ σίλφιον· ὁ γὰρ ὀπὸς καλούμενος τοῦ σιλφίου δάκρυόν ἐστιν. ἡ δὲ σκαμμωνία καὶ εἴ τι ἄλλο τοιοῦτον, ὥσπερ ἐλέχθη, φαρμακώδεις ἔχουσι τὰς δυνάμεις.

Πάντων δὲ τῶν εἰρημένων τὰ μὲν αὐτομάτως συνίσταται, τὰ δ' ἀπ' ἐντομῆς, τὰ δ' ἀμφοτέρωθεν τέμνουσι δε δήλον ὅτι τὰ χρήσιμα καὶ τὰ μάλλον επιζητούμενα, τοῦ δ' ἀπὸ τῆς ἀμυγδαλῆς οὐδεμία χρεία δακρύου, δι' δ κούκ άφελκοῦσι. πλην έκεινό γε φανερον ότι ων αυτόματος ή πηξις 6 τούτων πλείων ή επιρροή της ύγρότητος. οὐ τὴν αὐτὴν δ' ὥραν ἀπάντων αί ἐντομαὶ καὶ ἡ πῆξις. άλλὰ τὸ μὲν τῆς ἀμπέλου μάλιστα συνίστασθαί φασιν έὰν μικρὸν πρὸ τῆς βλαστήσεως τμηθῆ, τοῦ δὲ μετοπώρου καὶ ἀρχομένου τοῦ χειμῶνος ήττον καίτοι πρός γε καρποτοκίαν αι ωραιόταται ταίς γε πλείσταις αύται. της δε τερμίνθου καὶ της πεύκης και εί έκ τινων άλλων ρητίνη γίνεται μετά την βλάστησιν το δ' όλον ουκ επέτειος ή τούτων, άλλ' εἰς πλείω χρόνον ἡ ἐντομή. τὸν δὲ λιβανωτὸν καὶ τὴν σμύρναν ὑπὸ Κύνα φασὶ καὶ ταις θερμοτάταις ήμέραις εντέμνειν ωσαύτως δὲ καὶ τὸ ἐν Συρία βάλσαμον.

' Ακριβεστέρα δὲ καὶ ἐλάττων ἡ καὶ τούτων

^{1 &}amp; abτηs conj. Scal.: cf. Plin. 19. 162, where smyrnium is given as a synonym; ἐν αὐτοῖs Ald.

ENQUIRY INTO PLANTS, IX. 1. 4-7

from it: 1 for, as was said, 2 this plant can be grown from an exudation, like the krinonia (lily) and other plants. The juice of silphium is pungent like the plant itself; for what is called the 'juice' of silphium is a gum. Scammony and similar plants, as was

said.8 have medicinal properties.

In all the plants mentioned the juice either forms naturally, or when incisions are made, or in both ways,4 but it is obvious that men only make incisions in plants whose juice is of use and is specially sought after.5 Now there is no use in the gum which exudes from the almond, wherefore men do not tap it.6 However it is plain that in plants whose gum forms naturally the flow of juice is greater. The incisions and the clotting of the juice do not take place at the same season in all cases; 7 but the juice of the vine clots best they say if the incision is made a little before budding begins, less well in the autumn or at the beginning of winter; (although in regard to production of fruit these 8 seasons are the best in the case of most 9 vines). However with terebinth fir or any other tree which produces resin the best time is after the period of budding; yet in general these trees are not cut every year, but at longer intervals. The frankincense and myrrh trees they say should be cut at the rising of the Dogstar and on the hottest days, and so also the 'Syrian balsam' (balsam of Mecca).

The cutting of these is also a more delicate matter

6 κουκ αφελκουσιν conj. Scal., cf. 9. 2. 1; καν αφέλκουσιν U;

κὰν ἀφέλκωσι MAld. of. Plin. 24. 105.

7 Plin. 24. 106.

8 αl conj. W.; καl UPAld.

9 γε conj. Sch.; δὲ Ald. of. U.P. 3. 13. 2.

⁵ μαλλον ἐπιζητούμενα· τοῦ δ' ἀπὸ τῆς ά. conj. W. supported by G; μᾶλλον ἐπὶ γοῦν τὸ ἀπὸ τῆς à. UMAld.

έντομή· καὶ γὰρ ἡ συρροὴ τῆς ὑγρότητος ἐλάττων· ὧν δὲ καὶ ὁ καυλὸς ἐντέμνεται καὶ ἡ ῥίζα, τούτων ὁ καυλὸς πρότερον, ὥσπερ καὶ τοῦ σιλφίου, καὶ καλοῦσι δὲ τῶν ὀπῶν τούτων τὸν μὲν καυλίαν τὸν δὲ ῥιζίαν· καὶ ἐστι βελτίων ὁ ῥιζίας· καθαρὸς γὰρ καὶ διαφανὴς καὶ ξηρότερος. ὁ δὲ καυλίας ὑγρότερος· καὶ διὰ τοῦτο ἄλευρον αὐτῷ περιπάττουσι πρὸς τὴν πῆξιν. τὴν ὥραν τῆς ἐντομῆς ἴσασιν οἱ Λίβυες· οὖτοι γὰρ οἱ σίλφιον λέγοντες. ὡσαύτως δὲ καὶ οἱ ῥιζοτόμοι καὶ οἱ τοὺς φαρμακώδεις ὀποὺς συλλέγοντες· καὶ γὰρ οὖτοι τοὺς καυλοὺς ὀπίζουσι πρότερον. ἀπλῶς δὲ πάντες καὶ οἱ τὰς ῥίζας καὶ οἱ τοὺς ὀποὺς συλλέγοντες τὴν οἰκείαν ὥραν ἑκάστων τηροῦσι. καὶ τοῦτο μὲν δὴ κοινόν.

II. Ἡ δὲ ρητίνη γίνεται τόνδε τὸν τρόπου ἐν μὲν τῆ πεύκη ὅταν ἀφελκωθείσης ἡ δᾶς ἐξαιρεθῆ, συρρεῖ γὰρ εἰς τὸ ἔλκωμα τοῦτο πλείων ἡ ὑγρότης, ἐν δὲ τῆ ἐλάτη καὶ τῆ πίτυι ὅταν γευσάμενοι τῶν ξύλων ἀφελκώσωσιν οὐ γὰρ πᾶς ἀφορισμὸς ὁμοίως ἀφελκοῦσι γὰρ καὶ τὰς τερμίνθους ἐν ἀμφοῦν καὶ ἐν τῷ στελέχει καὶ ἐν τοῖς ἀκρεμόσιν αἰεὶ δὲ πλείων καὶ βελτίων ἡ εἰς τὸ στέλεχος

συρρέουσα της είς τους άκρεμόνας.

Διαφέρουσι δὲ καὶ κατὰ τὰ δένδρα. βελτίστη μὲν γὰρ ἡ τερμινθίνη καὶ γὰρ συνεστηκυῖα καὶ εὐωδεστάτη καὶ κουφοτάτη τῆ ὀσμῆ ἀλλ' ὀλίγη. δευτέρα δὲ ἡ ἐλατίνη καὶ πιτυίνη, κουφότεραι γὰρ τῆς πευκίνης. πλείστη δὲ ἡ πευκίνη καὶ

¹ cf. 6. 3. 2; C.P. 6. 11. 16.

² σίλφιον conj. St.; σιλφίου UM; σιλφιολέγοντες PAld.

ENQUIRY INTO PLANTS, IX. 1. 7-11. 2

and is done on a smaller scale; for the flow of juice is less. In those plants whose stem and root are both cut the stem is cut first, as also with silphium; and the juices so obtained are called respectively stalk-juice and root-juice, of which the latter is the better, for it is clear transparent and less liquid. The stalk-juice is more liquid, and for this reason they sprinkle meal 1 over it to make it clot. The Libyans know the season for cutting, for it is they that gather the silphium. So also do the root-diggers and those that collect medicinal juices, for these too tap the stems earlier. And in general all those who collect whether roots or juices observe the season which is appropriate in each case. And this remark applies generally.

Of resinous trees and the methods of collecting resin and pitch.

II. 8 Resin is made in the following manner:—in fir it is done by removing the resinous wood after the tree has been tapped; for then the juice flows into the hole so made in greater abundance; in silver-fir and Aleppo pine it is done by tapping the wood, after tasting it. For there is no fixed rule for all alike; thus with terebinth they tap both the stem and the branches; but the juice which runs into the stem is always more abundant and better than that which flows into the branches.

There are also differences in 4 the resin obtained from different trees. The best is that of terebinth; for it sets firm, is the most fragrant, and has the most delicate smell; but the yield is not abundant. Next comes that of silver-fir and Aleppo pine, for these are more delicate than that of the fir. But that of the fir

⁸ Plin, 16. 57. ⁴ κατὰ conj. W.; ταῦτα Ald. H.

βαρυτάτη καὶ πιττωδεστάτη διὰ τὸ μάλιστα ένδαδον είναι την πεύκην. ἄγεται δε εν άσκοις ύγρά, κάπειτα οΰτω συνίσταται. καίτοι φασί καί την τέρμινθον πιττοκαυτείσθαι περί Συρίαν έστι γαρ όρος, καθάπερ έν τοῖς ἔμπροσθεν εἴπομεν, μέγα

τερμίνθων μεστον απαν μεγάλων.

Ενιοι δέ φασι καὶ τὴν πίτυν καὶ τὴν κέδρον δὲ την Φοινικικήν άλλα ταθτα μεν ώς ενδεχόμενα ληπτέον διὰ τὸ σπάνιον ἐπεὶ οί γε περὶ Μακεδονίαν οὐδὲ τὴν πεύκην πιττοκαυτοῦσιν ἀλλ' ἢ την άρρενα καλούσι γαρ άρρενα την μη καρποφόρου. της δε θηλείας εάν τινα των ριζών λάβωσιν άπασα γὰρ ἔνδαδος πεύκη ταις ρίζαις. καλλίστη δὲ πίττα γίνεται καὶ καθαρωτάτη ἡ ἐκ τῶν σφόδρα προσείλων καὶ προσβόρρων, ἐκ δὲ τῶν παλισκίων βλοσυρωτέρα καὶ βορβορώδης ἐν γάρ τοις σφόδρα παλισκίοις οὐδε φύεται πεύκη τὸ παράπαν.

Έστι δὲ καὶ ἀφορία τις καὶ εὐφορία καὶ πλήθους καὶ καλλονής. ὅταν μὲν γὰρ χειμών μέτριος γίνηται, πολλή γίνεται καὶ καλή καὶ τῷ χρώματι λευκοτέρα, ὅταν δὲ ἰσχυρός, ὀλίγη καὶ μοχθηροτέρα. καὶ ταῦτά γέ ἐστι τὰ ὁρίζοντα πλήθος καὶ καλλουήν πίττης, ούχ ή πολυκαρπία τῶν πευκῶν.

⁵ Plin. 16, 59.

¹ πιττοκαυτεῖσθαι conj. Sch., cf. 9. 3. 4 ; πιττοκαυθίσαι U ; ττωθεῖσαι Ald. ² 3. 2. 6. πιττωθείσαι Ald.

^{*} δè conj. W.; καὶ Ald. H. cf. 3. 12. 3.
* μὴ conj. W.; γε Ald. H.; γε μὴ Cod. Casaub. Vin.; γε μὴν
Vo. (τὴν ἄκαρπον mBas.). cf. 3. 9. 2.

ENQUIRY INTO PLANTS, IX. II. 2-4

is the most abundant, the grossest and the most pitchlike, because this tree has the greatest amount of resinous wood. It is carried about in baskets in a liquid state, and so acquires the more solid form which we know. However they say that in Syria pitch is extracted even from the terebinth by burning 1; for there is in that land a mountain which, as we said

before,2 is all covered with great terebinths.

Some ³ say the same of Aleppo pine and also of Phoenician cedar; but this must be taken as only indicating what can be done, the practice not being common; for the people of Macedonia do not extract pitch by burning even from fir, except from the 'male' kind (they call the kind which bears no fruit ⁴ the 'male'); the 'female' kind they only treat in this way when they have found roots containing pitch; for all firs have resinous wood extending to the roots. ⁵The finest and purest pitch is that obtained from trees growing in a sunny position and facing north ⁶; that obtained from trees growing in shade is coarser ⁷ and muddy; (in exceedingly shady places the fir does not even grow at all).

Again the yield may be either good or bad as to amount and as to quality; thus, when there is a moderate winter, it is abundant and good and whiter in colour, but, when there is a severe winter, it is scanty and of inferior quality. And it is these conditions, and not the tree's capacity for bearing fruit, which determine the amount and quality of pitch.

225

VOL. II.

⁶ Apparently because this is the dry quarter in the Balkan peninsula.

⁷ βλοσωρωτέρα conj. Sch.; βροσηροτέρα M; βλοσηροτέρα Ald. cf. C.P. 6. 12. 5.

- 5 Οἱ δὲ περὶ τὴν Ἰδην φασί, διαιροῦντες τὰς πεύκας καὶ τὴν μὲν καλοῦντες Ἰδαίαν τὴν δὲ παραλίαν, τὴν ἐκ τῆς Ἰδαίας πλείω καὶ μελαντέραν γίνεσθαι καὶ γλυκυτέραν καὶ τὸ ὅλον εὐωδεστέραν ὤμήν, ἐψηθεῖσαν δὲ ἐλάττω ἐκβαίνειν· πλείω γὰρ ἔχειν τὸν ὀρρόν, δι' ὁ καὶ λεπτοτέραν εἶναι. τὴν δὲ τῆς παραλίας ξανθοτέραν καὶ παχυτέραν ὡμήν, ὥστε καὶ τὴν ἄφεψιν ἐλάττω γίνεσθαι, δαδωδεστέραν δὲ τὴν Ἰδαίαν. ὡς δὲ ἀπλῶς εἰπεῖν ἐκ τῆς ἴσης δαδὸς πλείω καὶ ὑδαρεστέραν ἐν ταῖς ἐπομβρίαις γίνεσθαι ἡ ἐν τοῖς αὐχμοῖς, καὶ ἐκ τῶν χειμερινῶν καὶ παλισκίων τόπων ἡ ἐκ τῶν εὐείλων καὶ εὐδιεινῶν. ταῦτα μὲν οὖν οὕτως ἐκάτεροι λέγουσιν.
 - 'Αναπληροῦσθαι δὲ συμβαίνει τὰ κοιλώματα πρὸς τὸ πάλιν ἐξαιρεῖν τῶν μὲν ἀγαθῶν πευκῶν ἐνιαυτῷ, τῶν δὲ μετριωτέρων ἐν δυσὶν ἔτεσι, τῶν δὲ μοχθηρῶν ἐν τρισίν. ἡ δὲ ἀναπλήρωσις οὐ τοῦ ξύλου καὶ τῆς συμφύσεως ἀλλὰ τῆς πίττης ἐστίν· ἐπεὶ τὸ ξύλον ἀδύνατον συμφῦναι καὶ ἔν γενέσθαι πάλιν, ἀλλ' ἡ ἐργασία διὰ τοσούτου χρόνου γίνεται τῆς πίττης· ἀναγκαῖον δὲ δῆλον ὅτι καὶ τῷ ξύλφ γίνεσθαί τινα πρόσφυσιν, εἴπερ

³ Plin. *l.c.*

¹ Plin. 16. 60.

² εὐωδεστέραν Η.; εὐκρινωδεστέραν UMAld.; ? εὐκρινεστέραν καὶ εὐωδεστέραν W. cf 3. 9. 2.

ENQUIRY INTO PLANTS, IX. 11. 5-6

The people of Mount Ida distinguish different kinds of fir, calling one 'that of Mount Ida' (Corsican pine), another the 'fir of the seashore,' (Aleppo pine); and they say that the pitch obtained from the former is more abundant blacker sweeter and generally more fragrant 2 in the raw state, but that boiling down reduces the amount; for that it contains a larger proportion of watery matter, wherefore it is less substantial; but that derived from the 'fir of the seashore' is browner and thicker in the raw state, so that the amount is less reduced by boiling down; that the 'fir of Mount Ida' however contains more resinous wood. And, speaking generally, they say that from an equal amount of resinous wood more pitch is obtained and in a more liquid state in wet weather than during a drought, and from a wintry and shady position than from one that is sunny and enjoys fair weather. the account given by the peoples of Mount Ida and of Macedonia respectively.

³ The holes for the pitch fill up, so that the pitch can be again removed, in good firs in a year, in those of more moderate quality in two years, in poor trees in three. The filling-up is composed of the pitch; it is not caused by closing up of the wood; for the wood cannot close up and become one again, but the effect which takes the time mentioned is due to the formation of the pitch. However it is clearly inevitable that there should be some new growth of the wood too, seeing that the resinous wood is

4 ¿ξαιρείν conj. Sch.; ¿ξαίρειν Ald.H.

⁵ So W. explains ἀλλά... πιττῆs. Or perhaps (as Sch.), 'however this is the interval which must elapse before the pitch can be worked again.'

έξαιρουμένης της δαδός καὶ καιομένης της πίττης ή έκροή. τοῦτο μὲν οὖν οὕτω ληπτέον.

- Οἱ δὲ περὶ τὴν Ἰδην φασίν, ὅταν λεπίσωσι τὸ στέλεχος,—λεπίζουσι δὲ τὸ πρὸς ἥλιον μέρος ἐπὶ δύο ἡ τρεῖς πήχεις ἀπὸ τῆς γῆς—ἐνταῦθα τῆς ἐπιρροῆς γινομένης ἐνδαδοῦσθαι ἐνιαυτῷ μάλιστα, τοῦτο δ' ὅταν ἐκπελεκήσωσιν ἐν ἑτέρω πάλιν ἐνδαδοῦσθαι καὶ τὸ τρίτον ὡσαύτως, μετὰ δὲ ταῦτα διὰ τὴν ὑποτομὴν ἐκπίπτειν τὸ δένδρον ὑπὸ τῶν πνευμάτων σαπέν τότε δ' ἐξαιρεῖν αὐτοῦ τὴν καρδίαν, τοῦτο γὰρ μάλιστα δαδῶδες, ἐξαιρεῖν δὲ ἐκ τῶν ῥιζῶν καὶ γὰρ ταύτας, ὥσπερ εἴπομεν, ἐνδάδους πασῶν.
- Εἰκὸς δὲ δῆλον ὅτι τὰς μὲν ἀγαθάς, ὅσπερ ἐλέχθη, συνεχῶς τοῦτο δρᾶν τὰς δὲ χείρονας διὰ πλείονος χρόνου· καὶ ταμιευομένων μὲν πλείω χρόνον ἀντέχειν, ἐὰν δὲ πᾶσαν ἐξαιρῶσιν ἐλάττω· δύναται δ' ὡς ἔοικε τρεῖς μάλιστα τοιαύτας ἐξαιρέσεις ὑπομένειν. οὐχ ἄμα δὲ καρποφοροῦσιν αὶ πεῦκαι καὶ δαδοφοροῦσι· καρποφοροῦσι μὲν γὰρ εὐθὺς νέαι, δαδοφοροῦσι δὲ ὕστερον πολλῷ πρεσβύτεραι γινόμεναι.

ΙΙΙ. Τὴν δὲ πίτταν καίουσι τόνδε τὸν τρόπον ὅταν κατασκευάσωσιν ὁμαλῆ τόπον ὥσπερ ἄλω

¹ i.e. and so this kind of wood at least is replaced by fresh growth.

² Plin. 16. 57.

³ τῆς ἐπιρροῆς γινομένης ἐνδαδοῦσθαι conj. W.; τὴν ἐπιρροὴν γινομένην ἔνδαδον Ald.; τῆς ἐπιρροῆς γινομένης ἔνδαδον γίνεσθαι conj. Sch.

ENQUIRY INTO PLANTS, IX. II. 6-III. 1

removed 1 and burnt when the discharge of pitch takes

place. So much for this account.

² The people of Mount Ida however say that, when they bark the stem,—and they bark the side towards the sun to a height of two or three cubits from the ground,—the flow of pitch takes place in that part,³ and in about a year the wood becomes full of pitch; and that, when they have hewn this part out, pitch forms again in the next year, and in the third year in like manner; after which ⁴ that the tree, because it has been cut away underneath, is rotted by the winds and falls; and that then ⁵ they take out its heart, for that is especially full of pitch, and that they also extract pitch from ⁶ the roots; for that these too, as we said,⁷ are full of pitch in all firs.

Now it is plainly to be expected that they should, as was said, repeatedly thus treat a good tree, but an inferior one at longer intervals, and that, if the tree is husbanded, the supply should hold out longer, while, if they remove all the pitch, it will not hold out so long; it appears as a matter of fact that the tree will stand about three such removals of its substance. ⁸ However firs do not produce both fruit and pitch at once; they begin to bear fruit when they are quite young, but they only produce pitch

much later, when they are older.

Of the making of pitch in Macedonia and in Syria.

III. This is the manner in which they make pitch by fire:—having prepared a level piece of ground,

 $^{^4}$ μετὰ δὲ ταῦτα conj. Sch.; τὰ δὲ ταῦτα UM; τὰ δὲ τοιαῦτα Ald.

⁵ τότε conj. Sch. from G; τοὺs Ald.

⁶ ἐκ Ald.; καὶ conj. W.

⁷ 9. 2. 3. ⁸ cf. C.P. 5. 16. 2.

ποιήσαντες έχουσαν είς τὸ μέσον συρροήν καὶ ταύτην έδαφίσωσι, κατασχίσαντες τοὺς κορμοὺς συντιθέασι παραπλησίαν σύνθεσιν της των άνθρακευόντων, πλην οὐκ ἔμβοθρον άλλὰ τὰς σχίζας ὀρθὰς πρὸς ἀλλήλας, ὥστε λαμβάνειν ὕψος αἰεὶ κατὰ πλήθος· γίνεσθαι δέ φασιν, ὅταν ἡ σύνθεσις ἢ κύκλφ μὲν ὀγδοήκοντα καὶ ἑκατὸν πηχέων είς ύψος δε εξήκοντα πλείστον ή πεντήκοντα ἡ ἐκατὸν ἀμφοτέροις, ἐάνπερ ἡ δậς τυγχάνη 2 πίειρα. συνθέντες οθν αθτήν οθτως και κατασκεπάσαντες ύλη γην επιβαλόντες κατακρύπτουσιν δπως μηδαμώς διαλάμψη τὸ πῦρ, ἀπόλλυται γαρ ή πίττα τούτου συμβάντος. υφάπτουσι δέ κατά την υπολειπομένην δίοδον είτα δε καί ταθτα επιφράξαντες τη ύλη και επιχώσαντες τηροῦσιν ἀναβαίνοντες κατὰ κλίμακος, ή αν δρωσι τον καπνον ωθούμενον, και επιβάλλουσιν αίει της γης όπως μηδ' ἀναλάμψη, κατεσκεύασται δὲ ὀχετὸς τῆ πίττη διὰ τής συνθέσεως τής ἀπορροής εἰς βόθυνον ὅσον ἀπέχοντα πεντεκαίδεκα πήχεις ή δ απορρέουσα της πίττης 8 ψυχρὰ γίνεται κατὰ τὴν άφήν. καίεται δὲ μάλιστα δύο ήμέρας και νύκτας τη γαρ ύστεραία προ ήλίου δύναντος εκκεκαυμένη γίνεται και ενδέδωκεν ή πυρά τοῦτο γὰρ συμβαίνει μηκέτι ρεούσης. τοῦτον δὲ τὸν χρόνον ἄπαντα τηροῦσιν

² cf. 5 9, where however the 'pit' is not described.

¹ ἐδαφίσωσι: cf. 9. 4. 4.

^{*} γίνεσθαι δὲ. Something seems to have dropped out at the beginning of this clause. ? "and they say that the pile at largest is 180". . .: so Sch. supplying μεγίστη after ἢ. The omitted words might also throw light on the preceding sentence.

ENQUIRY INTO PLANTS, IX. III. 1-3

which they make like a threshing-floor with a slope for the pitch to run towards the middle, and having made it smooth,1 they cleave the logs and place them in an arrangement like that used by charcoal-burners,2 except that there is no pit; but the billets are set upright against one another, so that the pile goes on growing in height according to the number used. And they say that the erection is complete. 8 when the pile is 180 cubits in circumference, and fifty, or at most sixty, in height; or again when it is a hundred cubits in circumference and a hundred in height,4 if the wood happens to be rich in pitch. Having then thus arranged the pile and having covered it in with timber they throw on earth and completely cover it, so that the fire may not by any means show through; for, if this happens, the pitch is ruined. Then they kindle the pile where the passage is left, and then, having 5 filled that part 6 up too with the timber and piled on earth, they mount a ladder and watch wherever they see the smoke pushing its way out, and keep on piling on the earth, so that the fire may not even shew itself. And a conduit is prepared for the pitch right though the pile, so that it may flow into a hole about fifteen cubits off, and the pitch as it flows out is now cold to the touch. The pile burns for nearly two days and nights; for on the second day before sunset it has burnt itself out and the pile has fallen in; for this occurs if the pitch is no longer flowing. All this time 7 they keep watch and do not

⁶ ταῦτα Ald ; ταύτην W. after Sch.'s conj.

⁴ αμφοτέροις: αμφοτέρωσε conj. Sch.

⁵ ἐπιφράξαντες conj. Scal. from G, elipant; ἐπάξαντες UM Ald.; ἐπισάξαντες Η.

⁷ τοῦτον δέ τὸν χρόνον conj. Sch. from G, totum tempus ; τόνδε δὲ τὸν τρόπον UMAld. Η.

άγρυπνοῦντες, ὅπως μὴ διαλάμπη, καὶ θύουσι δὲ καὶ ἑορτάζουσιν εὐχόμενοι πολλήν τε καὶ καλὴν γίνεσθαι τὴν πίτταν οἱ μὲν δὴ περὶ

Μακεδονίαν καίουσι τὸν τρόπον τοῦτον.

Έν δὲ τῆ ᾿Ασία φασί περὶ Συρίαν οὐκ ἐκπελεκῶντας τὴν δάδα ἀλλ᾽ ἐπ᾽ αὐτῷ τῷ δένδρῷ
προσκαίειν φέροντας ὅργανόν τι περιπεποιημένον
καὶ τούτῷ περιάπτοντας, εἶθ᾽ ὅταν ἐκτήξωσι
ταύτην πάλιν ἐπ᾽ ἄλλο καὶ ἄλλο μεταφέρειν
ὅρος δέ ἐστιν αὐτοῖς τις καὶ σημεῖα τοῦ παύεσθαι
καὶ μάλιστα δῆλον ὅτι τὸ μηκέτι ῥεῖν. πιττοκαυτοῦσι δέ, ὥσπερ καὶ πρότερον ἐλέχθη,
τὰς τερμίνθους· πεύκην γὰρ οὐ φέρουσιν οἱ
τόποι. τὰ μὲν οὖν περὶ τὴν ῥητίνην καὶ τὴν
πίτταν οὕτως ἔχει.

IV. Περὶ δέ λιβανωτοῦ καὶ σμύρνης καὶ βαλσάμου καὶ εἴ τι τοιοῦτον ἔτερον ὅτι μὲν καὶ ἀπ' ἐντομῆς γίνεται καὶ αὐτομάτως εἴρηται. ποία δε τις ἡ τῶν δένδρων φύσις καὶ εἴ τι περὶ τὴν γένεσιν ἡ τὴν συλλογὴν ἡ τῶν ἄλλων ἴδιον αὐτοῖς ὑπάρχει πειρατέον εἰπεῖν, ὡσαύτως δὲ καὶ περὶ τῶν λοιπῶν εὐόσμων σχεδὸν γὰρ τά γε πλεῖστα ἀπὸ τῶν τόπων ἐστὶ τῶν τε πρὸς μεσημβρίαν καὶ

ἀνατολήν.

Σ Γίνεται μὲν οὖν ὁ λίβανος καὶ ἡ σμύρνα καὶ ἡ κασία καὶ ἔτι τὸ κινάμωμον ἐν τἢ τῶν ᾿Αράβων Χερρονήσφ περί τε Σαβὰ καὶ ʿΑδραμύτα καὶ

² The sense given to περιπεποιημένον is unexampled, and the word may be corrupt.

 $^{^1}$ ἐκπελεκῶντας conj. W.; ἐκπλέκοντες U; ἐκπλέοντες M; ἐκπλήπτοντες Ald.

ENQUIRY INTO PLANTS, IX. III. 3-IV. 2

go to rest, in case the fire should come through; and they offer sacrifice and keep holiday, praying that the pitch may be abundant and good. Such is the manner in which the people of Macedonia make

pitch by fire.

They say that in Asia in the Syrian region they do not extract the pitch by cutting 1 out of the tree the wood containing it, but use fire to the tree itself, applying an instrument fashioned on purpose,2 with which they set fire to it.3 And then, when they have melted out the pitch at once place, they shift the instrument to another. But they have a limit and indications when to stop, chiefly of course the fact that the pitch ceases to flow. They also, as was said before,4 use fire to get pitch out of the terebinth; for the places where this tree grows do not produce the fir. Such are the facts about resin and pitch.

Of frankincense and myrrh: various accounts.

IV. As to frankincense myrrh balsam of Mecca and similar plants it has been said that the gum is produced both by incision and naturally. Now we must endeavour to say what b is the natural character of these trees and to mention any peculiarities as to the origin of the gum or its collection or anything belse. So too concerning the other fragrant plants; most of these come from places in the south and east.

Now frankincense myrrh cassia and also cinnamon are found in the Arabian peninsula 7 about

^{*} τούτφ περιάπτοντας seems to have been G's reading (Scal.); τοῦτο περιαλείφοντας MSS.

4 9. 2. 2.

ποία conj. W.; πολλή Ald.
 τι before τῶν add. Sch.
 Χερρονήσω conj. Salm.; χώρα νήσω Ald. cf. Plin. 6. 28.

Κιτίβαινα καὶ Μαμάλι. φύεται δὲ τὰ τοῦ λιβανωτοῦ καὶ τῆς σμύρνης δένδρα τὰ μὲν ἐν τῷ ὅρει τὰ δ' ἐν ταῖς ἰδίαις γεωργίαις ὑπὸ τὴν ὑπώρειαν, δι' ὁ καὶ τὰ μὲν θεραπεύεται τὰ δ' οὔ τὸ δ' ὅρος εἶναί φασιν ὑψηλὸν καὶ δασὰ καὶ νιφόμενον, ρεῖν δ' ἐξ αὐτοῦ καὶ ποταμοὺς εἰς τὸ πέδιον. εἶναι δὲ τὸ μὲν τοῦ λιβανωτοῦ δένδρον οὐ μέγα, πεντάπηχυ δέ τι καὶ πολύκλαδον, φύλλον δ' ἔχειν ἐμφερὲς τῆ ἀπίφ, πλὴν ἔλαττον πολὰ καὶ τῷ χρώματι ποῶδες σφόδρα, καθάπερ τὸ πήγανον

λειόφλοιον δὲ πᾶν, ὥσπερ τὴν δάφνην.

Τἡν δὲ σμύρναν ἔλαττον ἔτι τῷ μεγέθει καὶ θαμνωδέστερον δέ, τὸ δὲ στέλεχος ἔχειν σκληρὸν καὶ συνεστραμμένον ἐπὶ τῆς γῆς, παχύτερον δὲ ἡ κνημοπαχές· φλοιὸν δὲ ἔχειν λεῖον ὅμοιον τῆ ἀνδράχλη. ἔτεροι δὲ οἱ φάσκοντες τεθεωρηκέναι περὶ μὲν τοῦ μεγέθους σχεδὸν συμφωνοῦσιν· οὐδέτερον γὰρ εἶναι μέγα τῶν δένδρων, ἔλαττον δὲ τὸ τῆς σμύρνης καὶ ταπεινότερον· φύλλον δὲ ἔχειν τὸ τοῦ λιβανωτοῦ δαφνοειδὲς καὶ λειόφλοιον δ' εἶναι· τὸ δὲ τῆς σμύρνης ἀκανθῶδες καὶ οὐ λεῖον, φύλλον δὲ προσεμφερὲς ἔχειν τῆ πτελέα, πλὴν οῦλον ἐξ ἄκρου δὲ ἐπακανθίζον, ὥσπερ τὸ τῆς πρίνου.

"Εφασαν δε οὖτοι κατὰ τὸν παράπλουν δν εξ Ἡρώων ἐποιοῦντο κόλπου ζητεῖν ἐκβάντες ὕδωρ ἐν τῷ ὄρει καὶ οὕτω θεωρῆσαι τὰ δένδρα καὶ τὴν συλλογήν. εἶναι δ' ἀμφοτέρων ἐντετμημένα καὶ

* έχειν conj. Sch.; έχει P₂Ald.

¹ Plin. 12. 55 and 56. ² Plin. 12. 67.

⁴ λίβανωτοῦ δαφνοειδές καὶ λειόφλοιον δ' εἶναι conj. Sch.; λιβά-νου τοῦ δαφνοειδὲς καὶ λειόφυλλον δ' εἶναι UM; λιβάνου δαφνοειδὲς

ENQUIRY INTO PLANTS, IX. IV. 2-4

Saba Hadramyta Kitibaina and Mamali. The trees of frankincense and myrrh grow partly in the mountains, partly on private estates at the foot of the mountains; wherefore some are under cultivation, others not; the mountains, they say, are lofty, forest-covered and subject to snow, and rivers from them flow down to the plain. The frankincense-tree, it is said, is not tall, about five cubits high, and it is much branched; it has a leaf like that of the pear, but much smaller and very grassy in colour, like rue; the bark is

altogether smooth like that of bay.

The myrrh-tree ² is said to be still smaller in stature and more bushy; it is said to have ³ a tough stem, which is contorted near the ground, and is stouter than a man's leg; and to have a smooth bark like that of andrachne. Others who say that they have seen it agree pretty closely about the size; neither of these trees, they say, is large, but that which bears myrrh is the smaller and of lower growth; however they say that, while the frankincense-tree ⁴ has a leaf like that of bay and smooth bark, that which bears myrrh is spinous and not smooth, and has a leaf like that of the elm, except that it is curly and spinous ⁵ at the tip like that of kermes-oak.

⁶ These said that on the coasting voyage which they made from the bay of the Heroes they landed to look for water on the mountains and so saw these trees and the manner of collecting their gums. ⁷ They reported that with both trees incisions had

καὶ λειόφλοιον δ' εἶναι P_2 ; λιβάνου \cdot άλλα τοῦ μὲν δαφνοειδὲς καὶ λειόφυλλον εἶναι Ald.H. of. Plin.~12.~57.

⁵ cf. 3. 10. 1; 3. 11. 3.

⁷ Plin. 12, 58-62.

⁶ cf. Arr. Anab. 3. 5. 4; 7. 20. 1 and 2.

τὰ στελέγη καὶ τοὺς κλάδους, ἀλλὰ τὰ μὲν ὥσπερ άξίνη δοκείν τετμήσθαι τούς δε λεπτοτέρας έχειν τὰς ἐντομάς τὸ δὲ δάκρυον τὸ μὲν καταπίπτειν τὸ δὲ καὶ πρὸς τῶ δένδρω προσέγεσθαι. μεν ύποβεβλησθαι ψιάθους εκ φοινίκων πεπλεγμένας, ένιαχοῦ δὲ τὸ ἔδαφος μόνον ήδαφίσθαι καὶ καθαρον είναι και τον μεν επι των ψιάθων λιβανωτὸν είναι καὶ καθαρὸν καὶ διαφανή, τὸν δ' έπὶ της γης ήττον τον δ' έπὶ τοις δένδροις προσεχόμενον αποξύειν σιδήροις, δι' δ καλ φλοιον ενίοις 5 προσείναι. τὸ δὲ ὄρος ἄπαν μεμερίσθαι τοῖς Σαβαίοις, τούτους γάρ είναι κυρίους, δικαίους δέ τὰ πρὸς ἀλλήλους, δι' δ καὶ οὐδένα τηρείν ὅθεν καὶ αὐτοὶ δαψιλώς εἰς τὰ πλοῖα λαβόντες ἐνθέσθαι τοῦ λιβανωτοῦ καὶ τῆς σμύρνης ἐρημίας ούσης καὶ ἀποπλεῖν. ἔλεγον δ' ούτοι καὶ τόδε καὶ ἔφασαν ἀκούειν, ὅτι συνάγεται πανταχόθεν ἡ σμύρνα καὶ ὁ λιβανωτὸς εἰς τὸ ἱερὸν τὸ τοῦ ἡλίου. τοῦτο δ' είναι μεν των Σαβαίων άγιώτατον δε πολύ τῶν περὶ τὸν τόπον, τηρεῖν δέ τινας "Αρα-6 βας ενόπλους όταν δε κομίσωσιν, εκαστον σωρεύσαντα τὸν αύτοῦ καὶ τὴν σμύρναν ὁμοίως καταλιπείν τοίς έπὶ της φυλακής, τιθέναι δὲ έπὶ τοῦ σωροῦ πινάκιον γραφην έχον τοῦ τε πλήθους τῶν μέτρων και της τιμης ης δεί πραθηναι το μέτρον έκαστον όταν δε οί έμποροι παραγένωνται, σκοπείν τὰς γραφάς, ὅστις δ' αν αὐτοῖς ἀρέσκη

 $^{^1}$ πραθήναι conj. Sch. from G ; πρασθήναι U ; προσθείναι P_2 Ald.

²³⁶

ENQUIRY INTO PLANTS, IX. iv. 4-6

been made both in the stems and in the branches. but that, while the stems looked as if they had been cut with an axe, in the branches the incisions were slighter; also that in some cases the gum was dropping, but that in others it remained sticking to the tree; and that in some places mats woven of palm-leaves were put underneath, while in some the ground underneath was merely made level and clean; and that the frankincense on the mats was clear and transparent, that collected on the ground less so; and that that which remained sticking to the trees they scraped off with iron tools, wherefore sometimes pieces of bark remained in it. whole range, they said, belongs to the portion of the Sabaeans; for it is under their sway, and they are honest in their dealings with one another. Wherefore no one keeps watch; so that these sailors greedily took, they said, and put on board their ships some of the frankincense and myrrh, since there was no one about, and sailed away. They also reported another thing which they said they had been told, that the myrrh and frankincense are collected from all parts into the temple of the sun; and that this temple is the most sacred thing which the Sabaeans of that region possess, and it is guarded by certain Arabians in arms. And that when they have brought it, each man piles up his own contribution of frankincense and the myrrh in like manner, and leaves it with those on guard; and on the pile he puts a tablet on which is stated the number of measures which it contains, and the price for which each measure should be sold 1; and that, when the merchants come, they look at the tablets, and whichsoever pile pleases them, they

μετρησαμένους τιθέναι τὴν τιμὴν εἰς τοῦτο τὸ χωρίον ἔνθεν ἂν ἔλωνται, καὶ τὸν ἱερέα παραγενόμενον τὸ τρίτον μέρος λαβόντα τῆς τιμῆς τῷ θεῷ τὸ λοιπὸν αὐτοῦ καταλιπεῖν καὶ τοῦτο σῶν εἶναι τοῖς κυρίοις, ἔως ἂν ἔλωνται παραγενόμενοι.

"Αλλοι δέ τινες το μεν τοῦ λιβανωτοῦ δένδρον ὅμοιον εἶναί φασι σχίνφ καὶ τον καρπον ταῖς σχινίσι φύλλον δε ὑπέρυθρον εἶναι δε τον μεν ἐκ τῶν νέων λιβανωτον λευκότερον καὶ ἀοδμότερον, τον δ' ἐκ τῶν παρηκμακότων ξανθότερον καὶ εὐοσμότερον τὸ δε τῆς σμύρνης ὅμοιον τῆ τερμίνθφ, τραχύτερον δε καὶ ἀκανθωδέστερον, φύλλον δε μικρῷ στρογγυλότερον, τῆ δε γεύσει διαμασωμένοις ὅμοιον τῷ τῆς τερμίνθου εἶναι δε καὶ τούτων τὰ παρηκμακότα εὐοσμότερα.

Γίνεσθαι δὲ ἀμφότερα ἐν τῷ αὐτῷ τόπῳ· τὴν δὲ γῆν ὑπάργιλον καὶ πλακώδη, καὶ ὕδατα πηγαῖα σπάνια. ταῦτα μὲν οὖν ὑπεναντία τῷ νίφεσθαι καὶ ὕεσθαι καὶ ποταμοὺς ἐξιέναι· τὸ δὲ παρόμοιον εἶναι τὸ δένδρον τῆ τερμίνθῳ καὶ ἄλλοι τινὲς λέγουσιν, οἱ δὲ καὶ ὅλως τέρμινθον εἶναι· κομισθηναι γὰρ ξύλα πρὸς ᾿Αντίγονον ὑπὸ τῶν ᾿Αράβων τῶν τὸν λιβανωτὸν καταγόντων, ἃ οὐδὲν διέφερε τῶν τῆς τερμίνθου· πλὴν οὖτοί γε μεῖζον

¹ Plin. 12. 66 and 67. ² Plin. 12. 53.

³ πλακάδη: lit. 'with a crust'; so W., but the word does not seem to occur elsewhere in this sense.

⁴ cf. 9. 4. 2.

²³⁸

ENQUIRY INTO PLANTS, IX. IV. 6-8

measure, and put down the price on the spot whence they have taken the wares, and then the priest comes and, having taken the third part of the price for the god, leaves the rest of it where it was, and this remains safe for the owners until they come and claim it.

Others report that the tree which produces the frankincense is like mastich, and its fruit is like the fruit of that tree, but the leaf is reddish: also that the frankincense derived from young trees is whiter and less fragrant, while that derived from those which have passed their prime is yellower and more fragrant; also that the tree which produces myrrh is like the terebinth, but rougher and more thorny; that the leaf is somewhat rounder, and that, if one chews it, it resembles that of the terebinth in taste; also that of myrrh-trees too those that are past their prime give more fragrant myrrh.

¹ Both trees, it is said, grow in the same region; the soil is clayey ² and caked, ³ and spring waters are scarce. Now these reports are contradictory to ⁴ that which says that the country is subject to snow and rain and sends forth rivers. However others make the statement ⁵ that the tree is like the terebinth; in fact some say that it is the same tree; for that logs of it were brought to Antigonus by the Arabs who brought the frankincense down to the sea, and that these did not differ at all from logs of terebinth. ⁶ However these informants were guilty

⁶ τῶν conj. Sch.; τῷ UAld.

⁵ i.e. the statement quoted of the myrrh-tree, § 7. The 'tree' is here the λιβανωτός, but these authorities did not distinguish it from the myrrh-tree. See below.

ἔτερον ἀγνόημα προσηγνόουν· φοντο γὰρ ἐκ τοῦ αὐτοῦ δένδρου τόν τε λιβανωτὸν γίνεσθαι καὶ τὴν σμύρναν· διόπερ ἐκεῖνος ὁ λόγος πιθανώτερος ὁ παρὰ τῶν ἀναπλευσάντων ἐξ Ἡρώων πόλεως· ἐπεὶ καὶ τὸ ὑπὲρ Σάρδεων πεφυκὸς τοῦ λιβανωτοῦ δένδρον ἐν ἱερῷ τινι δαφνοειδὲς ἔχει τὸ φύλλον, εἴ τι δεῖ σταθμᾶσθαι τοῦτο· ὁ λιβανωτὸς δ' ἔχει, καὶ ὁ ἐκ τοῦ στελέχους καὶ ὁ ἐκ τῶν ἀκρεμόνων, ὁμοίως καὶ τῆ ὄψει καὶ τῆ ὀσμῆ θυμιώμενος τῷ ἄλλφ λιβανωτῷ. πέφυκε δὲ τοῦτο μόνον τὸ δένδρον οὐδεμιᾶς τυγχάνειν <θεραπείας>.

"Ενιοι δε λέγουσιν ώς πλείων μεν ό λιβανωτός έν τη 'Αραβία γίνεται, καλλίων δε έν ταις επικειμέναις νήσοις ων επάρχουσιν ένταιθα γαρ και σχηματοποιειν επί των δενδρων οι αν θελωσι και τάχα τοιτό γε οὐκ ἀπίθανον ενδέχεται <γαρ>
όποίαν αν βούλωνται ποιειν την εντομήν. εἰσι δε τινες και μεγάλοι σφόδρα των χόνδρων, ωστ εἰναι τῷ μεν ὅγκω χειροπληθιαίους σταθμῷ δε πλέον ἡ τρίτον μέρος μνας. ἀργὸς δε κομίζεται πας ὁ λιβανωτός, ὅμοιος δε τη προσόψει φλοιῷ. της σμύρνης δε ἡ μεν στακτη ἡ δε πλαστή. δοκιμάζεται δ' ἡ μεν ἀμείνων τῆ γεύσει, και ἀπὸ ταύτης την ὁμόχρων λαμβάνουσι. περι μεν οῦν λιβανωτοῦ και σμύρνης σχεδὸν τοσαῦτα ἀκηκόαμεν ἄχρι γε τοῦ νῦν.

* δ' έχει conj. W.; διέται P2Ald; om. H.

4 πλείων conj. W.; ἡδίων P2Ald.

¹ Plin. 15. 57; cf. 16. 135.

^{*} θεραπείαs add. Sch. from G; οὐδεμιᾶs τυγχάνειν UMAld.; οὐ μιᾶs τυγχάνειν P₂. But there is no sign of a lacuna in the MSS., and οὐδεμίαs is probably corrupt, as W. suggests. οὐδὲ θερ. is inconsistent with 9. 4. 2.

ENQUIRY INTO PLANTS, IX. IV. 8-10

of a further more important piece of ignorance; for they believed that the frankincense and the myrrh were produced by the same tree. Wherefore the account derived from those who sailed from the city of Heroes is more to be believed; ¹ in fact the frankincense-tree which grows above Sardes in a certain sacred precinct has ² a leaf like that of bay, if we may judge at all by this; and the frankincense derived both from its stem and its branches is like in appearance and in smell, when it is burnt as incense, to other frankincense. This is the only tree which can never be cultivated.³

Some say that the frankincense-tree is more abundant in Arabia, but finer in the adjacent islands 5 over which the Arabians bear rule; 6 for there it is said that they mould the gum on the trees to any shape that they please. And perhaps this is not incredible, since it is possible to make any kind of incision that they like. Some of the lumps 7 of gum are very large, so that one is large enough in bulk to fill the hand and in weight is more than a third of a pound. All frankincense is gathered in the rough and is like bark in appearance. Myrrh 8 is either 'fluid' (myrrh-oil) or 'solid' (agglutinated). That of better quality is tested by its taste, and of this they select that which is of uniform colour. 10 of frankincense and myrrh these are about all the facts that have come to our notice at present.

24 I

⁵ Plin. 12, 60.

⁶ επάρχουσιν conj. Coraës; ενυπάρχουσι P.Ald.

⁷ τῶν χόνδρων conj. Sch.; τῷ χόνδρῳ P₂Ald.

⁸ i.e. here the commodity so called.

⁹ cf. Odor. 29.

¹⁰ δμόχρων conj. Sch.; δμόχρω UM; δμόχρουν Ald.; την έτέραν καταλαμβάνουσι P₂.

V. Περί δὲ κιναμώμου καὶ κασίας τάδε λέγουσι· θάμνους μὲν ἀμφότερα ταῦτ' εἶναι οὐ μεγάλους ἀλλ' ἡλίκους ἄγνου, πολυκλάδους δὲ καὶ ξυλώδεις. ὅταν δὲ ἐκκόψωσιν, ὅλον τὸ κινάμωμον διαιρεῖν εἰς πέντε μέρη· τούτων δὲ τὸ πρῶτον πρὸς τοῖς βλαστοῖς βέλτιστον εἶναι, ὁ τεμνεται σπιθαμιαῖον ἡ μικρῷ μεῖζον· ἐπόμενον δὲ τὸ δεύτερον, ὁ καὶ τῆ τομῆ ἔλαττον· εἶτα τὸ τρίτον καὶ τέταρτον· ἔσχατον δὲ τὸ χείριστον τὸ πρὸς τῆ ρίζη· φλοιὸν γὰρ ἐλάχιστον ἔχειν· χρήσιμος δὲ οὖτος, οὐ τὸ ξύλον· δί ὁ καὶ τὸ ἀκροφυὲς κράτιστον, πλεῖστον γὰρ ἔχειν καὶ τὸν φλοιόν. οἱ μὲν οὕτω λέγουσιν.

"Αλλοι δὲ θαμνῶδες μὲν καὶ ἔτι μᾶλλον φρυγανῶδες εἶναί φασι δύο δ' αὐτοῦ γένη, τὸ μὲν
μέλαν τὸ δὲ λευκόν. λέγεται δέ τις καὶ μῦθος
ὑπὲρ αὐτοῦ· φύεσθαι μὲν γάρ φασιν ἐν φάραγξιν,
ἐν ταύταις δ' ὄφεις εἶναι πολλοὺς δῆγμα θανάσιμον ἔχοντας, πρὸς οῦς φραξάμενοι τὰς χεῖρας καὶ
τοὺς πόδας καταβαίνουσι καὶ συλλέγουσιν, εἶθ'
ὅταν ἐξενέγκωσι διελόντες τρία μέρη διακληροῦνται πρὸς τὸν ἥλιον, καὶ ἡν ἀν λάχη ὁ ἥλιος
καταλείπουσιν· ἀπιόντες δ' εὐθὺς ὁρῶν φασι καιομένην ταύτην· οὖτος μὲν οὖν τῷ ὄντι μῦθος.

Την δὲ κασίαν φασὶ τὰς μὲν ράβδους παχυτέρας ἔχειν, ἰνώδεις δὲ σφόδρα καὶ οὐκ εἶναι περιφλεῦσαι χρήσιμον δὲ καὶ ταύτης τὸν φλοιόν.

¹ Plin. 12. 85-94.

² A similar tale is told of frankincense by Herodotus (3. 107), who has an equally surprising tale about cinnamon (3. 111).

ENQUIRY INTO PLANTS, IX. v. 1-3

Of cinnamon and cassia: various accounts.

V. 1 Of cinnamon and cassia the following account is given: both are shrubs, it is said, and not of large size, but of the same size as bushes of chaste-tree. with many branches and woody. When they cut down the whole cinnamon-tree, they divide it into five parts; of these the first is that which grows next the branches and this is the best: this is cut in lengths a span long or a little longer; next comes the second kind, which is cut in shorter lengths; then come the third and the fourth, and last the least valuable wood, which grows next the root; for this has least bark, and it is the bark and not the wood which is serviceable; wherefore the part which grows high up the tree is the best, since it has the Such is the account given by some. most hark.

Others say that cinnamon is shrubby or rather like an under-shrub; and that there are two kinds, one black, the other white. ² And there is also a tale told about it; they say that it grows in deep glens, and that in these there are numerous snakes which have a deadly bite; against these they protect their hands and feet before they go down into the glens, and then, when they have brought up the cinnamon, they divide it in three parts and draw lots for it with the sun; and whatever portion falls to the lot of the sun they leave behind; and they say that, as soon as they leave the spot, they see this take fire. Now this is sheer fable.

³ Cassia, they say, has stouter branches, which are very fibrous and difficult to strip of the bark ⁴; and it is the bark of this tree also which is serviceable.

⁸ Plin. 12. 95-97.

⁴ περιφλοϊσαι Η.; περιφλεύσαι UMP2Ald. cf. de igne 72.

ὅταν οὖν τέμνωσι τὰς ῥάβδους, κατακόπτειν ὡς διδάκτυλα τὸ μῆκος ἡ μικρῷ μείζω, ταῦτα δ' εἰς νεόδορον βύρσαν καταρράπτειν εἶτ' ἐκ ταύτης καὶ τῶν ξύλων σηπομένων σκωλήκια γίνεσθαι, ἃ τὸ μὲν ξύλον κατεσθίει, τοῦ φλοιοῦ δ' οὐχ ἄπτεται διὰ τὴν πικρότητα καὶ δριμύτητα τῆς ὀσμῆς. καὶ περὶ μὲν κασίας καὶ κιναμώμου τοσαῦτα λέγεται.

VI. Το δε βάλσαμον γίνεται μεν εν τῷ αὐλῶνι τῷ περὶ Συρίαν. παραδείσους δ' εἶναί φασι δύο μόνους, τὸν μεν ὅσον εἴκοσι πλέθρων τὸν δ' ἔτερον πολλῷ ἐλάττονα. τὸ δε δένδρον μέγεθος μεν ἡλίκον ῥόα μεγάλη πολύκλαδον δε σφόδρα φύλλον δε ἔχειν ὅμοιον πηγάνῳ, πλὴν ἔκλευκον, ἀείφυλλον δὲ εἶναι· καρπὸν δὲ παρόμοιον τῆ τερμίνθῳ καὶ μεγέθει καὶ σχήματι καὶ χρώματι εὐῶδες σφόδρα καὶ τοῦτο καὶ μᾶλλον τοῦ δακρύου.

Τὸ δὲ δάκρυον ἀπὸ ἐντομῆς συλλέγειν, ἐντέμνειν δὲ ὄνυξι σιδηροῖς ὑπὸ τὸ ἄστρον, ὅταν μάλιστα πνίγη ὧσι, καὶ τὰ στελέχη καὶ τὰ ἄνω. τὴν δὲ συλλογὴν ὅλον τὸ θέρος ποιεῖσθαι οὐκ εἶναι δὲ πολὺ τὸ ῥέον, ἀλλ' ἐν ἡμέρα τὸν ἄνδρα συλλέγειν ὅσον κόγχην τὴν δ' ὀσμὴν διαφέρουσαν καὶ πολλήν, ὥστε ἀπὸ μικροῦ πολὺν ἐφικνεῖσθαι τόπον. ἀλλ' οὐ φοιτᾶν ἐνταῦθα ἄκρατον ἀλλὰ τὸ συνηγμένον κεκραμένον πολλὴν γὰρ δέχεσθαι

 $^{^1}$ νεόδορον conj. Sch.; νεόδερον P_2Ald . * Plin. 12 111–123. * εὐώδες . . . τοῦτο P_2Ald .; εὐώδη . . . τοῦτον W. after Sch.'s conj. But the clause begins without a conjunction, and some

ENQUIRY INTO PLANTS, IX. v. 3-vi. 2

When then they cut off the branches, they chop them up into lengths of about two fingers' breadth or rather more, and these they sew up in raw 1 hide; and then from the leather and the decaying wood little worms are engendered, which devour the wood but do not touch the bark, because it is bitter and has a pungent odour. This is all the information forthcoming about cinnamon and cassia.

Of balsam of Mecca.

VI. ² Balsam of Mecca grows in the valley of Syria. They say that there are only two parks in which it grows, one of about four acres, the other much smaller. The tree is as tall as a good-sized pomegranate and is much branched; it has a leaf like that of rue, but it is pale; and it is evergreen; the fruit is like that of the terebinth in size shape and colour, and this too is very fragrant, ³ indeed

more so than the gum.

⁴ The gum, they say, is collected by making incisions, which is done with bent pieces of iron at the time of the Dog star, when there is scorching heat; and the incisions are made both in the trunks and in the upper parts of the tree. The collecting goes on throughout the summer; but the quantity which flows is not large; in a day a single man can collect a shell-full ⁵; the fragrance is exceeding great and rich, so that that which comes from a small amount is perceived for a wide distance. However it does not reach us in a pure state; what is collected is mixed with other things; for it mixes freely with

words about the flower may have dropped out, to which this clause refers; cf. however Odor. 32.

⁴ Diose, 1, 19, ⁵ Plin, 12, 117,

κράσιν· καὶ τὸ ἐν τῆ Ἑλλάδι πολλάκις εἶναι κεκραμένον· εἴοσμα δὲ σφόδρα καὶ τὰ ῥαβδία·
δε καθαίρειν γὰρ καὶ τῶνδε ἔνεκα καί του διαφόρου·
πωλεῖσθαι <γὰρ> τίμια. καὶ τὴν ἐργασίαν τὴν
περὶ τὰ δένδρα σχεδὸν ἐν ταὐτῆ αἰτία εἶναι καὶ
τὴν βροχήν· βρέχεσθαι γὰρ συνεχῶς. συναιτίαν
δὲ δοκεῖν εἶναι τοῦ μὴ μεγάλα γίνεσθαι τὰ δένδρα
καὶ τὴν τῶν ῥαβδίων τομήν. διὰ γὰρ τὸ πολλά-
κις ἐπικείρεσθαι ῥάβδους ἀφιέναι καὶ οὐκ εἰς ἐν
ἐκτείνειν τὴν ὁρμήν.

"Αγριον δὲ οὐδὲν εἶναι βάλσαμον οὐδαμοῦ· γίνεσθαι δὲ ἐκ μὲν τοῦ μείζονος παραδείσου ἀγγείδια δώδεκα ὅσον ἡμιχοαῖα, ἐκ δὲ τοῦ ἑτέρου δύο μόνον πωλεῖσθαι δὲ τὸ μὲν ἄκρατον δὶς πρὸς ἀργύριον τὸ δ' ἄλλο κατὰ λόγον τῆς μίξεως καὶ τοῦτο μὲν διαφέρον τι φαίνεται κατὰ τὴν εὐοσ-

μίαν.

VII. 'Ο δὲ κάλαμος γίνεται καὶ ὁ σχοῖνος ὑπερβάλλοντι τὸν Λίβανον μεταξὺ τοῦ τε Λιβάνου καὶ ἄλλου τινὸς ὅρους μικροῦ ἐν τῷ αὐλωνίσκῷ τούτῷ, καὶ οὐχ ὡς τινές φασι τοῦ ᾿Αντιλιβάνου· ὁ γὰρ ᾿Αντιλίβανος μακρὰν ἀπέχει τοῦ Λιβάνου καὶ μεταξὺ τούτων ἐστὶν δν αὐλῶνα καλοῦσι πεδίον πολὺ καὶ καλόν. ὅπου δὲ ὁ κάλαμος καὶ ὁ σχοῖνος φύεται λίμνη μεγάλη τυγχάνει, πρὸς ταύτην δὲ ἐν τῷ ἔλει τῷ ἀνεξηραμμένῷ πεφύκασι·

 $^{^1}$ τov I conj.; $\tau o\hat{v}$ MSS. G's rendering shews that the explanation of the 'different reason' (i.e. to strengthen the tree) has dropped out of our texts. 2 $\gamma a \rho$ add. Sch.

ENQUIRY INTO PLANTS, IX. vi. 2-vii. 1

other things; and what is known in Hellas is generally mixed with something else. The boughs are also very fragrant. In fact it is on account of these boughs, they say, that the tree is pruned (as well as for a different reason 1), since 2 the boughs cut off can be sold for a good price. In fact the culture of the trees has the same motive 3 as the irrigation (for they are constantly irrigated). And the cutting of the boughs seems likewise to be partly the reason why the trees do not grow tall; for, since they are often cut about, they send out branches instead of putting out all their energy in one direction.

Balsam is said not to grow wild anywhere. From the larger park are obtained twelve vessels containing each about three pints, from the other only two such vessels; the pure gum sells for twice its weight in silver, the mixed sort at a price proportionate to its purity. Balsam then appears to be of exceptional fragrance.

Of other aromatic plants-all oriental, except the iris.

VII. ⁴ Sweet-flag and ginger-grass grow beyond the Libanus between that range and another small range, in the depression thus formed; and not, as some say, between Libanus and Anti-Libanus. For Anti-Libanus is a long way from Libanus, and between them is a wide fair plain called 'The Valley.' But, where the sweet-flag and ginger-grass grow, there is a large lake,⁵ and they grow near it in the dried up marshes, covering an extent

The 'motive' is the production of boughs. ἐν ταὐτῆν alτίᾳ I conj.; ταὐτην alτίαν UMP, Ald.; ἐν ταὐτην alτίᾳ conj. W.
 Plin. 12. 104 and 105,
 cf. C.P. 6. 18. 2.

τοπον δὲ ἔχουσι πλείον ἢ τριάκοντα σταδίων. οὐκ ὄζουσι δὲ χλωροὶ ἀλλὰ ξηρανθέντες, τἢ προσόψει δὲ οὐδὲν διαφέρουσι τῶν ἄλλων· εἰσ-βάλλοντι δ' εἰς τὸν τόπον εὐθὺς ὀσμὴ προσ-2 βάλλει· οὐ μὴν πορρωτέρω γε ἡ ἀποπνοὴ γίνεται, καθάπερ τινές φασι, ταῖς προσφερομέναις ναυσὶ πρὸς τὴν χώραν· καὶ γὰρ ὁ τόπος οὖτος ἀπὸ θαλάττης ἀπέχει πλείους ἢ ἐκατὸν πεντήκοντα σταδίους· ἀλλ ἐν τῆ ᾿Αραβία τὴν ἀποπνοὴν

είναί φασι την άπο της χώρας εύοσμον.

Έν μὲν οὖν Συρία τὰ περιττὰ τῆ ὀσμῆ σχεδὸν ταῦτ' ἐστίν ἡ γὰρ χαλβάνη βαρύτερον καὶ μᾶλλον φαρμακῶδες ἐπεὶ καὶ αὕτη γίνεται περὶ Συρίαν ἐκ τοῦ πανάκους καλουμένου. τὰ δὲ ἄλλα πάντα τὰ εὕοσμα οἷς πρὸς τὰ ἀρώματα χρῶνται, τὰ μὲν ἐξ Ἰνδῶν κομίζεται κἀκεῦθεν ἐπὶ θάλατταν καταπέμπεται, τὰ δ' ἐξ Ἰλραβίας, οἷον πρὸς τῷ κιναμώμῳ καὶ τῆ κασία καὶ κώμακον ἔτερον δ' εἶναι τὸ κώμακον καρπόν τὸ δ' ἔτερον παραμίσγουσιν εἰς τὰ σπουδαιότατα τῶν μύρων. τὸ δὲ καρδάμωμον καὶ ἄμωμον οἱ μὲν ἐκ Μηδείας, οἱ δ' ἐξ Ἰνδῶν καὶ ταῦτα καὶ τὴν νάρδον καὶ τὰ ἄλλα ἡ τὰ πλεῖστα.

΄ Οἱς μὲν οὖν εἰς τὰ ἀρώματα χρῶνται σχεδὸν τάδε ἐστί· κασία κινάμωμον καρδάμωμον νάρδος ναῖρον βάλσαμον ἀσπάλαθος στύραξ ἰρις νάρτη

¹ οὐκ ὄζουσι conj. Guilandinus, cf. C.P. 6. 14. 8; οὐ δοκοῦσι P₂Ald.H.

 $^{^{\}circ 2}$ cf. C.P. 6. 18. 1. των άλλων: sc. καλάμων καὶ σχοίνων. $^{\circ 3}$ προσφερομέναις ναυσὶ πρὸς conj. Scal.; προσφ. είναι πρὸς P_2 Ald.

⁴ cf. 9. 9. 2 n. 'The plant,' i.e. one of the plants so called.
⁵ Plin. 12. 135; cf. 13. 18.

ENQUIRY INTO PLANTS, IX. vii. 1-3

of more than thirty furlongs. They have no fragrance 1 when they are green, but only when they are dried, and in appearance they do not differ 2 from ordinary reeds and rushes; but, as you approach the spot, immediately a sweet smell strikes you. However it is not true, as some say, that the fragrance is wafted to ships approaching 3 the country; for indeed this district is more than 150 furlongs from the sea. However it is said that in Arabia the breezes wafted from the land are fragrant.

Such then are the plants in Syria which have remarkable fragrance. For that of khalbane is more oppressive and somewhat medicinal; for this perfume also is produced in Syria from the plant called allheal. As to all the other fragrant plants used for aromatic odours, they come partly from India whence they are sent over sea, and partly from Arabia, for instance, komakon 5—as well as cinnamon and cassia. The fruit called komakon is said to be distinct from this; the komakon of which we are speaking is a perfume which they mix with the choicest unguents. Cardamom and Nepaul cardamom some say come from Media; others say that these come from India, as well as spikenard and most, if not all, of the other species.

Now this is a general list of the plants used for perfumes:—cassia cinnamon cardamom spikenard nairon balsam of Mecca aspalathos storax iris narte

⁶ elvai has no government, and W. considers the passage corrupt. Comparison of Plin. l.c. suggests that the original text may have been something like this: τὸ κάμακον καρπὸν δν· τὸν δὲ χυλὸν κ.τ.λ., i.e. 'komakon is of different character from these, being a fruit, whose juice—.' Plin. calls κ. a kind of cinnamon.

κόστος πάνακες κρόκος σμύρνα κύπειρον σχοίνος κάλαμος ἀμάρακον λωτὸς ἄννητος. τούτων δὲ τὰ μὲν ρίζαι τὰ δὲ φλοιοὶ τὰ δὲ κλῶνες τὰ δὲ ξύλα τὰ δὲ σπέρματα τὰ δὲ δάκρυα τὰ δὲ ἄνθη. καὶ τὰ μὲν πολλαχοῦ γίνεται, τὰ δὲ περιττότατα καὶ εὐοδμότατα πάντα ἐκ τῆς 'Ασίας καὶ ἐκ τῶν ἀλεεινῶν τόπων. ἐκ γὰρ αὐτῆς Εὐρώπης οὐδέν ἐστιν ἔξω τῆς ἴριδος.

Αύτη δ' ἀρίστη ἐν Ἰλλυριοῖς, οὐκ ἐν τῇ πρὸς θάλατταν χώρα, ἀλλ' ἐν τῇ ἀνακεχωρηκυία κειμένῃ δὲ μᾶλλον πρὸς ἄρκτον. τόποι δὲ τόπων διαφέρουσιν ἐν οἶς ἀμείνων ἐργασία δὲ περὶ αὐτὴν οὐδεμία πλὴν τοῦ περικαθήραντα ἀναξηρᾶναι.

Τὰ γὰρ ἐν τἢ Θράκη φυόμενα ριζία, καθάπερ τό τε τἢ νάρδφ προσεμφερῆ τὴν ὀσμὴν ἔχον καὶ ἔτερ' ἄττα, μικράν τινα καὶ ἀσθενῆ τὴν εὐωδίαν ἔχει. καὶ περὶ μὲν τῶν εὐόσμων ἐπὶ τοσοῦτον εἰρήσθω.

VIII. Περί δὲ τῶν ὀπῶν ὅσα μὴ πρότερον εἴρηται, λέγω δ' οἶον εἴ τινες φαρμακώδεις ἡ καὶ
ἄλλας ἔχουσι δυνάμεις, πειρατέον ὁμοίως εἰπεῖν·
ἄμα δὲ καὶ περὶ ρίζῶν, καὶ γὰρ τῶν ὀπῶν τινες ἐκ
τούτων καὶ χωρὸς αὐταὶ καθ' αὐτὰς πολλὰς καὶ
παντοίας ἔχουσι δυνάμεις, ὅλως δὲ περὶ πάντων
φαρμακωδῶν, οἶον καρποῦ χυλισμοῦ φύλλων ριζῶν πόας· καλοῦσι γὰρ καὶ πόαν ἔνια τῶν φαρμακωδῶν οἱ ρίζοτόμοι.

of. C.P. 6. 14. 8; 6. 18. 12; Plin. 21. 40.
 See Index App. (25).

ENQUIRY INTO PLANTS, IX. vii. 3-viii. 1

kostos all-heal saffron-crocus myrrh kypeiron gingergrass sweet-flag sweet marjoram lotos dill. Of these it is the roots, bark, branches, wood, seeds, gum or flowers which in different cases yield the perfume. Some of them grow in many places, but the most excellent and most fragrant all come from Asia and sunny regions. From Europe itself comes none of them except the iris.

¹ This is best in Illyria, not in the part near the sea, but in that which is further inland and lies more to the north. In different districts it varies in quality; no special attention is needed, except to

scrape the roots clean and dry them.

As for the roots which grow in Thrace, such as one 2 which has a smell like spikenard and certain others, their fragrance is but slight and feeble. Let this suffice for an account of sweet-smelling plants.

Of the medicinal juices of plants and the collection of them:

VIII. Now we must endeavour to speak in like manner of those juices which have not been mentioned already, I mean, such as are medicinal or have other properties; and at the same time we must speak of roots; for some of the juices are derived from roots, and apart from that roots have in themselves divers properties of all kinds; and in general we must discuss medicinal things of all kinds, as fruit, extracted juice, leaves, roots, herbs'; for the herb-diggers call some medicinal things by this name.

^{*} χυλισμοῦ P.Ald.H; καυλοῦ conj. W. The list is of the aspects in which the herbalist would regard the plant, not of the parts of the plant.

Των δε ριζων πλείους μέν είσιν αι δυνάμεις καλ πρὸς πλείω. ζητοῦνται δὲ μάλιστα αἱ φαρμακώδεις ώς χρησιμώταται, διαφέρουσαι τῶ τε μὴ πρὸς ταὐτὰ καὶ τῷ μὴ ἐν τοῖς αὐτοῖς ἔχειν τὴν δύναμιν. ώς δ' οὖν ἐπὶ πᾶν αὶ πλεῖσται μὲν ἐν αύταις έχουσι και τοις καρποις και τοις όποις, ένιαι δε καὶ έν τοῖς φύλλοις τὰς δε φυλλώδεις δυνάμεις τὰς πολλὰς σχεδὸν πόας καλοῦσιν, ώσπερ εξρηται μικρώ πρότερον, οἱ ῥιζοτόμοι.

Ο μεν οθν όπισμος γίνεται των όπιζομένων ώς έπὶ τὸ πολύ τοῦ θέρους, τῶν μὲν ἐνισταμένου τῶν δὲ προεληλυθότος. ή δὲ ριζοτομία γίνεταί τινων καὶ ὑπὸ πυροτομίαν καὶ μικρῷ πρότερον, οὐ μὴν άλλ' ή γε πλείων τοῦ μετοπώρου μετ' Αρκτοῦρον, όταν φυλλορροήσωσιν, όσων δέ καὶ ὁ καρπὸς γρήσιμος, όταν άμερθωσι τον καρπόν. έστι δε δ όπισμὸς ἡ ἀπὸ τῶν καυλῶν, ὥσπερ τοῦ τιθυμάλλου καὶ τῆς θριδακίνης καὶ σγεδὸν τῶν πλείστων, η ἀπὸ τῶν ριζῶν, ἡ τρίτον ἀπὸ τῆς κεφαλης, ὥσπερ της μήκωνος ταύτης γαρ μόνης ούτω καὶ τοῦτ' ίδιον αὐτης. των μεν οὖν καὶ αὐτόματος ὁ ὀπὸς

² ριζων: ρίζα signifies a medicinal plant in general (cf. ριζότομοι) as well as 'root'; the double sense makes translation of this section awkward: I print it 'root' where it has

the wider sense.

¹ From this point to 9. 19. 4, γίνεσθαι, the text is repeated in U, with considerable variations, as a tenth book. Ald. also repeats the first few lines of this passage (down to end of § 1) as a fragment of a tenth book. The two Medicean MSS also repeat 9.8.1, των δε ριζων, down to 9.10.3, βέλτιστοι δε και οίς, as part of a tenth book. The tenth book' readings in each case are distinguished by a *.

ENQUIRY INTO PLANTS, IX. viii. 1-2

¹ The properties of 'roots'² are numerous and they have numerous uses; but those which have medicinal virtues are especially sought after, as being the most useful; and they differ in not all being applied to the same ³ purposes and in not all having their virtue in the same parts of them.⁴ To speak generally, most 'roots' have it in themselves ⁵; or else it is found in the fruits or the juices of the plant; and in some cases in the leaves as well, and it is to the virtues of the leaves in most cases that the herb-diggers refer, when they speak, as has just been said.⁶ of 'herbs.'

The collection of the juice from plants from which it is collected is mostly done in summer, in some cases at the beginning of that season, in others when it is well advanced. The digging of roots is done in some cases at the time of wheat-harvest or a little earlier, but the greater part of it in autumn after the rising of Arcturus when the plants have shed their leaves, and, in the case of those whose fruit is serviceable, when they have lost their fruit. The collection of juice is made either from the stalks, as with tithymallos (spurge) wild lettuce and the majority of plants, or from the roots, or thirdly from the head, as in the case of the poppy; for this is the only plant which is so treated and this is its peculiarity. In some plants the juice collects of its own accord in

³ ταὐτὰ conj. Scal. from G; ταῦτα Ald.

After δύναμιν U*Ald.* add ωσπερ εἴρηται μικρῷ πρότερον; omitted here by Sch.: see below.
5 Sc. in the roots.

⁶ ἄσπερ...πρότερον inserted here by Sch.: see above; ἄσπερ εἴρηται Ald.

⁷ καυλῶν Vin. Vo.Cod. Cas.: so also G; καρπῶν Ald. HM*.
⁸ μόνης οὕτω καὶ conj. W.; μόνης καὶ Ald. H; μόνον οὕτω καὶ M*.

συνίσταται δακρυώδης τις, ὅσπερ καὶ τῆς τραγακάνθης· ταύτην γὰρ οὐδὲ τέμνειν ἐστί· τῶν δὲ πλείστων ἀπὸ τῆς ἐντομῆς. ὧν ἐνίους μὲν εὐθὺς εἰς ἀγγεῖα συνάγουσιν, ὅσπερ καὶ τὸν τοῦ τιθυμάλλου ἡ μηκωνίου, καλοῦσι γὰρ ἀμφοτέρως, καὶ ἀπλῶς ὅσα πολύοπα τυγχάνει· τῶν δὲ μὴ πολυόπων ἐρίῳ λαμβάνουσιν ὥσπερ καὶ τῆς θριδακίνης.

- Ἐνίων δ' οὐδ' ὁπισμὸς ἀλλ' οἶον χυλισμός ἐστιν, ὥσπερ ὅσα κόψαντες ἡ τρίψαντες καὶ ὕδωρ ἐπιχέαντες ἀπηθοῦσι καὶ λαμβάνουσι τὴν ὑπόστασιν ξηρὸς δὲ δῆλον ὅτι καὶ ἐλάττων ὁ χυλὸς τούτων. ἔστι δὲ τῶν μὲν ἄλλων ῥιζῶν τὸ χύλισμα ἀσθενέστερον τοῦ καρποῦ, τοῦ κωνείου δὲ ἰσχυρότερον, καὶ τὴν ἀπαλλαγὴν ῥάω ποιεῖ καὶ θάττω μικρὸν πάνυ καταπότιον δοθέν ἐνεργότερον δὲ καὶ εἰς τὰς ἄλλας χρείας. ἰσχυρὸν δὲ καὶ τὸ τῆς θαψίας. τὰ δὲ ἄλλα πάντα ἀσθενέστερα. οἱ μὲν οὖν ὀπισμοὶ σχεδὸν τοσαυταχῶς γίνονται.
- Τῆς δὲ ριζοτομίας οὐκ ἔστι τοιαύτη διαφορὰ πλὴν ἐν ταῖς ὥραις οἷον θέρους ἢ μετοπώρου, καὶ τῷ τάσδε ἢ τάσδε τῶν ριζῶν· οἷον τοῦ ἐλλεβόρου τὰς κάτω τὰς λεπτάς· τὴν γὰρ ἄνω τὴν παχεῖαν τὴν κεφαλώδη φασὶν ἀγρεῖον εἶναι καὶ διδόναι

² cf. Diosc. 3. 7.

^{· 1} cf. Diosc. 2. 136; Plin. 20. 58.

³ ράω conj. Sch.; ραιο U; ραδίως M*Ald.

ENQUIRY INTO PLANTS, IX. vIII. 2-4

the form of a sort of gum, as with tragacanth; for incision of this plant cannot be made; but in most it is obtained by incision. In some cases the juice is collected straight into vessels, for instance that of tithymallos (spurge) or mekonion (for the plant has both names) and in general the juice of specially juicy plants is so collected. But that of those which do not yield abundant juice is taken with a piece of wool, as also that of wild lettuce.¹

² In some cases there can be no collection of juice, but there is a sort of extraction of it, for instance in the case of plants which are cut down or bruised; they then pour water over them and strain off the fluid, keeping the sediment; but it is plain that in these cases the juice obtained is dry and less copious. In most 'roots' the juice thus extracted is less powerful than that of the fruit, but in hemlock it is stronger and it causes an easier ³ and speedier ⁴ death even when administered in a quite small pill; and it is also more effective for other uses. ⁵ That of thapsia is also powerful, while all the rest are less so. Such then is a general account of the various ways of obtaining the juices of plants.

Of the cutting of roots for medicinal purposes, and of certain superstitions connected therewith.

As to cutting of the roots there is no such diversity of practice, except as to the season, which may be summer or autumn, and as to the particular roots selected. ⁶ Thus in hellebore the slender lower roots are taken, for they say that the thick upper part ⁷ which forms a sort of head is useless, and that

⁵ cf. Plin. 13. 125. ⁶ Plin. 25. 53. ⁷ i.e. rhizome.

⁴ θάττω conj. Sch.; ἐλάττω UM; θᾶττον U*M*Ald.

ταις κυσιν όταν βούλωνται καθαίρειν. και έφ έτέρων δέ τινων τοιαύτας λέγουσι διαφοράς.

- Έτι δὲ ὅσα οἱ φαρμακοπῶλαι καὶ οἱ ῥιζοτόμοι τὰ μὲν ἴσως οἰκείως τὰ δὲ καὶ ἐπιτραγωδοῦντες λέγουσι. κελεύουσι γὰρ τὰς μὲν κατ' ἄνεμον ἰσταμένους τέμνειν, ὥσπερ ἐτέρας τέ τινας καὶ τὴν θαψίαν, ἀλειψάμενον λίπα· τὸ γὰρ σῶμα ἀνοιδεῖν ἐὰν ἐξ ἐναντίας. κατ' ἄνεμον δὲ καὶ τοῦ κυνοσβάτου τὸν καρπὸν συλλέγειν, εἰ δὲ μὴ κίνδυνον εἶναι τῶν ὀφθαλμῶν. τὰς δὲ νύκτωρ τὰς δὲ μεθ' ἡμέραν, ἐνίας δὲ πρὶν τὸν ἤλιον ἐπιβάλλειν, οἶον καὶ τὸ καλούμενον κλύμενον.
- Καὶ ταῦτα μὲν καὶ τὰ παραπλήσια τούτοις τάχ' ἃν οὐκ ἀλλοτρίως δόξειεν λέγειν ἐπισινεῖς γάρ τινων αἱ δυνάμεις ἐξάπτειν γάρ φασιν ῶσπερ πῦρ καὶ κατακαίειν ἐπεὶ καὶ ὁ ἐλλέβορος ταχὺ καρηβαρεῖν ποιεῖ, καὶ οὐ δύνανται πολὺν χρόνον ὀρύττειν, δι' ὁ καὶ προεσθίουσι σκόροδα καὶ ἄκρατον ἐπιπίνουσιν. ἀλλὰ τὰ τοιαῦτα ῶσπερ ἐπίθετα καὶ πόρρωθεν, οἰον τὴν παιωνίαν, οἱ δὲ γλυκυσίδην καλοῦσι, νύκτωρ κελεύουσιν ὀρύττειν ἐὰν γὰρ ἡμέρας καὶ ὀφθῆ τις ὑπὸ δρυοκολάπτου τὸν μὲν καρπὸν ἀπολέγων κινδυνεύειν τοῖς ὀφθαλμοῖς, τὴν δὲ ῥίζαν τέμνων ἐκπίπτειν τὴν ἔδραν.

¹ ἔτι δὲ ὅσα conj. Sch. from G ; ἔτι δ' ώs U*; ἔτι δὲ ώs Ald. Η : ἐστι δὲ ώs M*.

²⁵⁶

ENQUIRY INTO PLANTS, IX. viii. 4-6

it is only given to dogs when it is desired to purge them. And in certain other plants also such differences are mentioned.

Further we may ¹ add statements made by druggists and herb-diggers, which in some cases may be to the point, but in others contain exaggeration. Thus they enjoin that in cutting some roots one should stand to windward,—for instance, in cutting thapsia among others, and that one should first anoint oneself with oil,² for that one's body will swell up if one stands the other way. Also that the fruit of the wild rose must be gathered standing to windward, since otherwise there is danger to the eyes. Also that some roots should be gathered at night, others by day, and some before the sun strikes on them, for instance those of the plant called honevsuckle.³

These and similar remarks may well seem to be not off the point, for the properties of these plants are hurtful; they take hold, it is said, like fire and burn; ⁴ for hellebore too soon makes the head heavy, and men cannot go on digging it up for long; wherefore they first eat garlic and take a draught of neat wine therewith. On the other hand the following ideas may be considered far-fetched and irrelevant; ⁵ for instance they say that the peony, which some call glykyside, should be dug up at night, for, if a man does it in the day-time and is observed by a woodpecker while he is gathering the fruit, he risks the loss of his eyesight; and, if he is cutting the root at the time, he gets prolapsus ani.

257

VOL. II.

² Plin. 13. 124; Diosc. 4. 153.

⁸ cf. 9. 18. 6. ⁴ Plin. 25. 50.

⁵ Plin. 27. 85; 25. 29.

- Φυλάττεσθαι δὲ καὶ τὴν κενταυρίδα τέμνοντα τριόρχην, ὅπως αν ἄτρωτος ἀπέλθη. καὶ ἄλλας δέ τινας αιτίας. τὸ δ' ἐπευχόμενον τέμνειν οὐθὲν ἴσως ἄτοπον· ἀλλ' εἴ τι καὶ ἄλλο προστιθέασιν. οίον όταν τὸ πάνακες τὸ 'Ασκληπίειον καλούμενον' άντεμβάλλειν γάρ τη γη παγκαρπίαν <καί> μελιττοῦταν όταν δὲ τὴν ξίριν, τριμήνου μελιττούτας αντεμβάλλειν μισθόν τέμνειν δε αμφήκει Είφει περιγράψαντα είς τρίς καὶ ὅτι ἄν πρῶτον τμηθή μετέωρον έχειν είθ' ουτω τὸ έτερον τέμνειν.
 - Καὶ ἄλλα δὲ τοιαῦτα πλείω. περιγράφειν δὲ καὶ τὸν μανδραγόραν εἰς τρὶς ξίφει, τέμνειν δὲ πρὸς έσπέραν βλέποντα. τὸν δ' ἔτερον κύκλω περιοργείσθαι καὶ λέγειν ώς πλείστα περὶ ἀφροδισίων. τοῦτο δ' ὅμοιον ἔοικε τῷ περὶ τοῦ κυμίνου λεγομένω κατά την βλασφημίαν δταν σπείρωσι. περιγράφειν δὲ καὶ τὸν ἐλλέβορον τον μέλανα και τέμνειν ιστάμενον προς έω και κατευχόμενον ἀετὸν δὲ φυλάττεσθαι καὶ ἐκ δεξιάς καὶ έξ ἀριστεράς κίνδυνον γὰρ είναι τοῖς τέμνουσιν, εάνπερ εγγύς επιγένηται ο άετος, άπο-

⁸ Plin. 25. 30 and 31.

¹ Plin. 25. 69 adds that this plant was therefore also called τρίορχις. cf. Diosc. 4. 162.
² και . . . αἰτίας U*M*; ? και ἄλλα δὲ τοιαῦτα W.

ENQUIRY INTO PLANTS, IX. viii. 7-8

It is also said that, while cutting feverwort 1 one must beware of the buzzard-hawk, if one wishes to come off unhurt; and other reasons for caution 2 are also given. That one should be bidden to pray while cutting is not perhaps unreasonable, but the additions made to this injunction are absurd; for instance as to cutting the kind of all-heal which is called that of Asklepios; 3 for then it is said that one should put in the ground in its place an offering made of all kinds of fruits and a cake; and that, when one is cutting gladwyn,4 one should put in its place to pay for it cakes of meal from spring-sown wheat.5 and that one should cut it with a two-edged sword. first making a circle round it three times, and that the piece first cut must be held up in the air while the rest is being cut.

And many similar notions are mentioned. Thus it is said that one should draw three circles round mandrake with a sword, and cut it with one's face towards the west; and at the cutting of the second piece one should dance round the plant and say as many things as possible about the mysteries of love. (This seems to be like the direction given about cummin,⁷ that one should utter curses at the time of sowing.) One should also, it is said, draw a circle round the black hellebore and cut it standing towards the east and saying prayers, and one should look out for an eagle both on the right and on the left; for that there is danger to those that cut, if your eagle should come near, that

5 τριμήνου conj. Salm.; τριμήνους M*Ald.H.

⁴ cf. Plin. 21. 42, who read Iριν. cf. Diosc. 4. 22, where ξυρίs is called a kind of Iριs; so also Plin. 21. 142.

τρὶs conj. Sch.; τρεῖs U[‡]M[‡]P₂Ald. So also in next section.
 cf. 7. 3. 3.

θνήσκειν ενιαυτφ. ταῦτα μεν οὖν επιθέτοις ἔοικεν, ὥσπερ εἴρηται. τρόποι δ' οὐκ εἰσὶ τῶν

ριζοτομιών πλην οθς εἴπομεν.

τρόποι δ΄ οὐκ εἰσὶ τῶν ομεν.

ΙΧ. "Εστι δέ, ὥσπερ ἐλέχθη, τῶν μὲν πάντα χρήσιμα καὶ ἡ ρίζα καὶ ὁ καρπὸς καὶ ὁ ὀπός, ιοσπερ ἄλλων τε καὶ τοῦ πανάκους τῶν δὲ ἡ ρίζα καὶ ὁ ἀπός, οἰον τῆς σκαμμωνίας καὶ τοῦ κυκλαμίνου καὶ τῆς θαψίας καὶ ἐτέρων, καθάπερ καὶ τοῦ μανδραγόρου τοῦ γὰρ μανδραγόρου τὸ φύλλον χρήσιμον εἰναί φασι πρὸς τὰ ἔλκη μετ ἀλφίτου, τὴν δὲ ρίζαν πρὸς ἐρυσίπελας ξυσθεῖσάν τε καὶ ὄξει δευθεῖσαν καὶ πρὸς τὰ ποδαγρικὰ καὶ πρὸς ὕπνον καὶ πρὸς φίλτρα διδόασι δ' ἐν οἰνω ἡ ὄξει τέμνουσι δὲ τροχίσκους ισπερ ραφανίδος καὶ ἐνείραντες ὑπὲρ γλεύκους ἐκρέμασαν ἐπὶ καπνῷ.

'Ο δὲ ἐλλέβορος ἐπὶ ταὐτὰ τῆ τε ῥίζη καὶ τῷ καρπῷ χρήσιμος, εἴπερ οἱ ἐν 'Αντικύρᾳ, καθάπερ φασί, τῷ καρπῷ καθαίρουσιν ἔχει δὲ <τὸν>

σησαμώδη τοῦτον.

Πλείω δὲ καὶ τοῦ πανάκους τὰ χρήσιμα καὶ οὐ πάντα πρὸς τὰ αὐτά· ἀλλ' ὁ μὲν καρπὸς πρὸς τὰς ἐξαμβλώσεις καὶ τὰς δυσουρίας, ὁ δὲ ἀπὸς ἡ χαλβάνη καλουμένη πρός τε τὰς ἐξαμβλώσεις καὶ πρὸς τὰ σπάσματα καὶ τοὺς τοιούτους πόνους, ἔτι δὲ πρὸς τὰ ὧτα καὶ τὰς φωνασκίας·

4 cf. 9. 12. 1.

 $^{^1}$ έδνπερ έγγὺς ἐπιγένηται conj.W.; ἐδν δὲ δ ἐγγὺς μὴ ἀποτέμνη δ αἐτὸς ἀποθνήσκει ἐνιαυτῷ UMAld. Similar confusion with variations in U*M*PP $_2$: restoration a makeshift. cf. Plin. 25. 50. 2 9. 8. 1.

³ Diosc. 4. 75; Plin. 26. 104 and 121.

ENQUIRY INTO PLANTS, IX. vIII. 8-1X. 2

they may die 1 within the year. These notions then seem to be irrelevant, as has been said. There are however no methods of root-cutting besides those which we have mentioned.

Of the medicinal uses of divers parts of plants.

IX. As was said,2 of some plants the root, fruit and juice are all serviceable, as of all-heal among others; of some the root and the juice, as of scammony 3 cyclamen thapsia and others, such as mandrake; for the leaf of this, they say, used with meal, is useful for wounds, and the root for erysipelas, when scraped and steeped in vinegar, and also for gout, for sleeplessness, and for love potions. It is administered in wine or vinegar; they cut little balls of it, as of4 radishes, and making a string of them hang them up in the smoke over must.

⁵ Of hellebore both root and fruit are useful for the same purposes,6—if it is true, as is said, that the people of Anticyra use the fruit as a purge; this fruit contains the well-known 7 drug called sesamodes.

Various parts of all-heal are also useful, and not all for the same purposes; the fruit is used in cases of miscarriage and for disorders of the bladder, while the juice, 8 which is called khalbane, is used in cases of miscarriage and also for sprains and such-like troubles; also for the ears, and to strengthen the

6 ταὐτὰ conj. Sch. from G; ταῦτα U*M*Ald.

Plin. 12, 126.



⁵ Diosc. 4. 108, 109 and 162.

⁷ I have inserted τδν. cf. 9. 14. 4; Plin. 22. 133; 25. 52 and 64; Diosc. l.c. The drug was actually called σησαμοειδές or σησαμοειδήs. For the sense of τοῦτον cf. 3. 7. 3; 3. 8. 3 and reff.

8 This seems to be a mistake. cf. 9. 7. 2; Diosc. 3. 83;

ή δὲ ρίζα πρός τε τοὺς τόκους καὶ τὰ γυναικεῖα καὶ πρὸς ὑποζυγίων φύσας χρησίμη δὲ καὶ πρὸς τὸ ἴρινον μύρον διὰ τὴν εὐωδίαν ἰσχυρότερον δὲ τὸ σπέρμα τῆς ρίζης. γίνεται δὲ περὶ Συρίαν καὶ τέμνεται περὶ πυραμητόν.

Τοῦ δὲ κυκλαμίνου ἡ μὲν ρίζα πρός τε τὰς έκπυήσεις τῶν φλεγμονῶν καὶ πρόσθετον γυναιξί καὶ πρὸς τὰ ἔλκη ἐν μέλιτι ὁ δὲ ὀπὸς πρὸς τὰς άπὸ κεφαλής καθάρσεις ἐν μέλιτι ἐγχεόμενος, καὶ πρὸς τὸ μεθύσκειν, ἐὰν ἐν οἴνω διάβρέχων διδῶ τις πίνειν. ἀγαθὴν δὲ τὴν ρίζαν καὶ ὧκυτόκιοι περίαπτον καὶ εἰς φίλτρα· ὅταν δὲ ὀρύξωσι, κατακαίουσιν είτ' οίνω δεύσαντες τροχίσκους ποιοῦσιν, ωσπερ της τρυγός ή ρυπτόμεθα.

Καὶ τοῦ σικύου δὲ τοῦ ἀγρίου τὴν μὲν ρίζαν <πρὸς> ἀλφοὺς καὶ ψώρας βοσκημάτων τὸ δὲ σπέρμα χυλισθέν ποιεί τὸ έλατήριον. συλλέγεται δὲ τοῦ φθινοπώρου τότε γὰρ βέλτιστον.

Της δε χαμαίδρυος τὰ μεν φύλλα πρὸς τὰ ρήγματα και πρὸς τὰ τραύματα ἐν ἐλαίω τριβόμενα καὶ πρὸς τὰ νεμόμενα έλκη· τὸν δὲ καρπὸν καθαίρειν χολήν· ἀγαθὸν δὲ καὶ ὀφθαλμοῖς· πρὸς δὲ τὰ ἄργεμα προσάγειν τὸ φύλλον τρίψαντα ἐν έλαίω. έχει δε φύλλα μεν οξάπερ δρύς, το δε ανάστημα της όλης όσον σπιθαμιαίον εύοσμον δὲ καὶ ἡδύ.

Τὸ μέν οὖν μὴ πρὸς ταὐτὸ πάντα τὰ μέρη χρήσιμα τυγγάνειν οὐκ ἴσως ἄτοπον τὸ δὲ τῆς

³ cf. Plin. 23. 63.

¹ enunhaeis conj. Coraës from Plin. 26. 120, eruptiones; ἐκπνεύσεις M*Ald.

² Diosc. 2, 164; Plin. 25, 133; 26, 149.

ENQUIRY INTO PLANTS, IX. 1X. 2-5

voice. The root is used in childbirth, for diseases of women, and for flatulence in beasts of burden. It is also useful in making the iris-perfume because of its fragrance; but the seed is stronger than the root. It grows in Syria and is cut at the time of wheat-harvest.

Of cyclamen the root is used for suppurating 1 boils; also as a pessary for women and, mixed with honey, for dressing wounds; the juice for purgings of the head, 2 for which purpose it is mixed with honey and poured in; it also conduces to drunkenness, if one is given a draught of wine in which it has been steeped. They say also that the root is a good charm for inducing rapid delivery and as a love potion; 3 when they have dug it up, they burn it, and then, having steeped the ashes in wine, make little balls like those made of wine-lees which we use as soap.

⁴ Of 'wild cucumber' (squirting cucumber) the root is used for ⁵ white leprosy and for mange in sheep, while the extracted juice makes the drug called 'the driver.' ⁶ It is collected in autumn, for then it is best.

is best.

Of germander the leaves pounded up in olive-oil are used for fractures and wounds and for spreading sores; the fruit purges bile, and is good also for the eyes; for ulcers in the eye they pound up the leaf in olive-oil before applying it. It has leaves like the oak, but its entire growth is only about a palm high; and it is sweet both to smell and taste.

Now that all parts are not serviceable for the same purpose is perhaps not strange; it is more

Digitized by Google

⁴ Diosc. 4. 150; Plin. 20. 3. ⁵ πρδs add. St. ⁶ cf. 9. 14. 1 and 2. ⁷ cf. 7. 6. 2; Diosc. 3. 98.

αὐτῆς ρίζης τὸ μὲν ἄνω τὸ δὲ κάτω καθαίρειν θαυμασιώτερον, οἶον καὶ τῆς θαψίας καὶ τῆς ἰσχάδος, οἱ δ' ἄπιον καλοῦσι, καὶ τῆς λιβανωτίδος. ὅτι γὰρ αὖ καὶ κάτω καὶ ἄνω ταὐτὰ δύναται [καθαίρειν], καθάπερ τὸ ἐλατήριον, οὐθὲν ἄτοπον.

β Έχει δὲ ή θαψία φύλλον μὲν ὅμοιον τῷ μαράθῳ πλὴν πλατύτερον καυλὸν δὲ ναρθηκώδη

ρίζαν δὲ λευκήν.

'Η δ' ισχὰς ἡ ἄπιος φύλλον μὲν ἔχει πηγανῶδες βραχὺ καυλοὺς δ' ἐπιγείους τρεῖς ἡ
τέτταρας ρίζαν δὲ οἴανπερ ὁ ἀσφόδελος πλὴν
λεπυριώδη· φιλεῖ δὲ ὀρεινὰ χωρία καὶ κοχλακώδη. συλλέγεται δὲ τοῦ ἦρος. τοῦτο μὲν
οὖν ἴδιον τῶν εἰρημένων.

Χ. Ὁ δὲ ἐλλέβορος ὅ τε μέλας καὶ ὁ λευκὸς ὅσπερ ὁμώνυμοι φαίνονται περὶ δὲ τῆς ὄψεως διαφωνοῦσιν οἱ μὲν γὰρ ὁμοίους εἶναι, πλὴν τῷ χρώματι μόνον διαφέρειν τὴν ῥίζαν τοῦ μὲν λευκὴν τοῦ δὲ μέλαιναν οἱ δὲ τοῦ μὲν μέλανος τὸ φύλλον δαφνῶδες τοῦ δὲ λευκοῦ πρασῶδες, τὰς δὲ ῥίζας ὁμοίας πλὴν τῶν χρωμάτων. οἱ δ΄ οὖν ὁμοίους λέγοντες τοιάνδε φασὶν εἶναι τὴν μορφήν καυλὸν μὲν ἀνθερικώδη βραχὺν σφόδρα τῷ τοῦ νάρθηκος, μῆκος δ΄ ἔχον εὐθὺ δ΄ ἐκ τῆς ῥίζης

δτι γὰρ conj. W.; δσα γὰρ UU*M*; τὰ γὰρ M; τὸ γὰρ Ald.
 Diosc. 4. 153; Plin. 13. 124.
 Diosc. 4. 175.

ENQUIRY INTO PLANTS, IX. 1x. 5-x. 1

surprising that part of the same 'root' should purge upwards and another part downwards, as is the case with thapsia and iskhas—which some call apios (spurge)—and with libanotis; for 1 it is not strange that on the other hand the same parts should purge both upwards and downwards, as is the case with 'the driver.'

² Thapsia has a leaf like fennel, but broader, a

stalk like that of ferula, and a white root.

⁸ Iskhas (or apios) has a leaf like rue and short, three or four prostrate stems, and a root like that of asphodel, except that it is composed of scales; ⁴ it loves mountain districts with a gravelly soil. It is collected in spring. Now this account applies only to the above-mentioned plants.

Of hellebores, the white and the black: their uses and distribution.

X. ⁵ The white and the black hellebore appear to have nothing in common except the name. But accounts differ as to the appearance of the plants; some say that the two are alike and differ only in colour, the root of the one being white, of the other black; some however say that the leaf of the 'black' is like that of bay, that of the white like that of the leek, but that the roots are alike except for their respective colours. Now those who say that the two plants are alike describe the appearance ⁶ as follows:—the stem is like that of asphodel and very short; the leaf has broad divisions, and is extremely like that of ferula, but is long; it is closely attached to the root and

⁵ Plin. 25. 47-61. See Index. cf. 9. 11. 5 n.

⁴ cf. Diosc. 3. 134.

⁶ i.e. of the two plants regarded as one; but the text of the following description seems to be hopelessly confused.

ηρτημένον καλ επιγειόφυλλον πολύρριζον δ' εὖ μάλα ταις λεπταις καλ χρησίμοις.

'Αναιρεῖν δὲ τὸν μὲν μέλανα καὶ ἴππους καὶ βοῦς καὶ ὖς, δι' ὁ καὶ οὐδὲν νέμεσθαι τούτων· τὸν δὲ λευκὸν νέμεσθαι τὰ πρόβατα καὶ ἐκ τούτου πρῶτον συνοφθῆναι τὴν δύναμιν καθαιρομένων ἐκείνων· ὡραῖος δὲ μετοπώρου, τοῦ δ' ἢρος ἄωρος· ἀλλὰ πρὸς τὴν πυλαίαν οἱ ἐκ τῆς Οἴτης συλλέγουσι· πλεῖστος γὰρ ἐνταῦθα φύεται καὶ ἄριστος· μοναχοῦ δὲ φύεται τῆς Οἴτης περὶ τὴν Πυράν.

Μίσγεται δὲ πρὸς τὴν πόσιν, ὅπως εὐεμὲς ἢ, τὸ τῆς ἐλλεβορίνης σπέρμα· τοῦτο δ' ἐστὶ ποάριον.

- Φύεται δὲ ὁ μὲν μέλας πανταχοῦ· καὶ γὰρ ἐν τῆ Βοιωτίᾳ καὶ ἐν Εὐβοίᾳ καὶ παρ' ἄλλοις πολλοῖς ἄριστος δὲ ὁ ἐκ τοῦ Ἑλικῶνος, καὶ ὅλως τὸ ὅρος εὐφάρμακον. ὁ δὲ λευκὸς ὀλιγαχοῦ· βέλτιστοι δὲ καὶ οἶς χρῶνται μάλιστα τέτταρες ὁ Οἰταῖος ὁ Ποντικὸς ὁ Ἐλεάτης ὁ Μαλιώτης. φασὶ δὲ τὸν Ἐλεάτην ἐν τοῖς ἀμπελῶσι φύεσθαι καὶ ποιεῖν τὸν οἶνον οὕτω διουρητικὸν ὥστε λαγαροὺς εἶναι πάνυ τοὺς πίνοντας.
- 4 *Αριστος δὲ πάντων καὶ τούτων καὶ τῶν ἄλλων ὁ Οἰταῖος. ὁ δὲ Παρνάσιος καὶ ὁ Αἰτωλικός, γίνεται γὰρ καὶ ἐνταῦθα καὶ πολλοὶ καὶ ἀνοῦνται καὶ πωλοῦσιν οὐκ εἰδότες, [οὐχ ὅτε] σκληροὶ καὶ

¹ Which were held apparently at Thermopylae regularly in autumn and sometimes in spring: the meeting would give opportunities for sale. ἀλλὰ implies a spring meeting.

ENQUIRY INTO PLANTS, IX. x. 1-4

creeps on the ground; the plant has numerous roots, to wit, the slender roots which are serviceable.

Also they say that the black is fatal to horses oxen and pigs, wherefore none of these animals eat it; while the white is eaten by sheep, and from this circumstance the virtue of the plant was first observed, since it purges them; it is at its prime in autumn, and past its prime when spring comes. However the people of Mount Oeta gather it for the meetings 1 of the Amphictyons; for it grows there in greatest abundance and best, though at only one place in the district of Oeta, namely about Pyra.

(The seed of rupture-wort is mixed with the potion given to promote easy vomiting; this plant

is a small herb).

The black kind of hellebore grows everywhere; it is found in Boeotia, in Euboea and in many other places; but best is that from Mount Helicon, which mountain is in general rich in medicinal herbs. The white occurs in few places; the best and that which is most used comes from one of four places, Oeta,² Pontus, Elea, and Malea.³ They say that that of Elea grows in the vineyards and makes the wine so diuretic that those who drink it become quite emaciated.

But best of all these and better than that found anywhere else is that of Mount Oeta, while that of Parnassus and that of Aetolia (for the plant is common in these parts too and men buy and sell it, not knowing 4 the difference) are tough and ex-

² δ before Oiτaios add. Sch.

The words οὐχ ὅτε may have arisen from οὐκ εἰδότες.

² Μαλιώτης conj. Hahnemann, cf. Strabo 9. 3. 3; Μασσαλιώτης Ald. Plin. l.c. gives Parnassus as the fourth locality: cf. § 4.

ἄγαν περισκελεῖς. ταῦτα μὲν οὖν ὅμοια ταῖς

μορφαίς όντα ταίς δυνάμεσι διαφέροντα.

Καλοῦσι δὲ τὸν μέλανά τινες ἔκτομον Μελαμπόδιον, ὡς ἐκείνου πρῶτον τεμόντος καὶ ἀνευρόντος. καθαίρουσι δὲ καὶ οἰκίας αὐτῷ καὶ πρόβατα συνεπάδοντές τινα ἐπφδὴν καὶ εἰς ἄλλα

δὲ πλείω χρῶνται.

ΧΙ. Πολλά δέ ἐστι καὶ τὰ πανάκη καὶ οἱ τιθύμαλλοι καὶ ἔτερ' ἄττα. πάνακες γὰρ καλοῦσι πρῶτον μὲν τὸ ἐν Συρία, περὶ οὖ μικρῷ πρότερον εἰρηται. ἄλλα δὲ τὰ τρία, τὸ μὲν Χειρώνειον καλούμενον τὸ δ' ᾿Ασκληπίειον τὸ δ' Ἡράκλειον. ἔχει δὲ τὸ μὲν Χειρώνειον φύλλον μὲν ὅμοιον λαπάθφ μεῖζον δὲ καὶ δασύτερον, ἄνθος δὲ χρυσοειδές, ρίζαν δὲ μικράν φιλεῖ δὲ μάλιστα τὰ χωρία τὰ πίονα· χρῶνται δὲ πρός τε τοὺς ἔχεις καὶ τὰ φαλάγγια καὶ τοὺς σῆπας καὶ τὰ ἄλλα ἐρπετὰ διδόντες ἐν οἴνφ καὶ ἀλείφοντες μετ' ἐλαίου· τοῦ δ' ἔχεως τὸ δῆγμα καὶ καταπλάττοντες καὶ ἐν ὀξίνη πιεῖν διδόντες· ἀγαθὴν δὲ φασι καὶ ἐλκῶν ἐν οἴνφ καὶ ἐλαίφ καὶ φυμάτων ἐν μέλιτι.

ε΄ Τὸ δ' ᾿Ασκληπίειον τὴν ῥίζαν μῆκος μὲν ὡς σπιθαμὴν λευκὴν δὲ καὶ παχεῖαν σφόδρα, καὶ φλοιὸν παχὺν καὶ ἁλυκώδη· καυλὸν δὲ ἔχει γονατώδη πανταχόθεν, φύλλον δὲ οἶόνπερ ἡ θαψία πλὴν παχύτερον· ἀγαθὸν δὲ εἶναί φασι ἑρπετῶν

² 9. 9. 2. ³ Plin. 25. 32; 26. 139.

¹ From this phrase ξκτομον came to be used as a synonym for 'black hellebore.' of. Plin. 25. 47; Diosc. 4. 149; Hesych. and Galen, Lex. Hipp. s.v.

⁴ μικράν conj. H. from Plin. 25. 32. radix varva; μακράν U* Ald.

ENQUIRY INTO PLANTS, IX. x. 4-x1. 2

ceeding harsh. These plants then, while resembling the best form in appearance, differ in their virtues.

Some call the black the 'hellebore of Melampus,' saying that he first cut and discovered it. Men also purify horses and sheep with it, at the same time chanting an incantation; and they put it to several other uses.

Of the various kinds of all-heal.

XI. There are also several kinds of all-heal tithymallos (spurge) and other herbs. To begin with, one plant called all-heal is the one found in Syria, of which we have recently spoken.² ³ Then come the three other kinds, known as that of Chaeronea, that of Asclepios, and that of Heracles. That of Chaeronea has a leaf like monk's rhubarb, but larger and rougher, a golden flower, and a small 4 root; and it specially loves rich ground; they use it for the bites of snakes, spiders, vipers 5 and other reptiles, administering it in wine or anointing the place with it mixed with olive-oil. In treating a snake-bite they use a plaster of it, and also give a draught of it mixed with vinegar 6; and they also say that it is good for sores 7 when mixed with wine and olive-oil, and for tumours when mixed with honey.

⁸The kind called after Asklepios has a white and very stout root about a span long and a thick bark which is crusted with salt ⁹; its stem is jointed all the way up, its leaf like that of *thapsia*, but thicker; it is said that it is good to scrape and drink

καὶ ἐν δξίνη conj. Sch., cf. 9. 13. 3; ἐν δξίνη καὶ PAld.
 For the genitive cf. §§ 2, 3; Xen. Mem. 3. 8. 3.

⁸ Plin. 25. 30; Diosc. 3. 49.

⁵ σῆπας conj. Scal., cf. Arist. Mir. Ausc. 164; σῆτας Ald.

⁹ άλυκώδη: ? 'has a briny taste.'

τε ξύοντα πίνειν, καὶ σπληνὸς ὅταν αἶμα περὶ αὐτὸν ἐν μελικράτφ, καὶ κεφαλαίας τρίβοντα ἐν ἐλαίφ ἀλείφειν καὶ ἄλλο τι ἐὰν πονἢ τις ἀφανές, καὶ γαστρὸς ὀδύνης ἐν οἴνφ ξύοντα. δύνασθαι δὲ καὶ τὰς μακρὰς ἀρρωστίας ἐκκλίνειν. ἔπειτα τῶν ἑλκῶν τῶν μὲν ὑγρῶν ξηρὸν ἐπιπάττοντα προκατακλύζοντα ἐν οἴνφ θερμῷ, τῶν δὲ ξηρῶν ἐν οἴνφ δεῦσαι καὶ καταπλάττειν.

- Τὸ δ' Ἡράκλειον φύλλον μὲν ἔχει μέγα καὶ πλατὰ καὶ τρισπίθαμον πανταχῆ, ρίζαν δὲ ὡς δακτύλου τὸ πάχος δίκραν ἡ τρίκραν, τῆ γεύσει μὲν ὑπόπικρον τῆ δ' ὀσμῆ καθάπερ λιβανωτοῦ καθαροῦ· ἀγαθὴν δὲ τῆς ἱερᾶς νόσου μιγνυμένην φώκης πιτύα ὅσον τεταρτημόριον πίνειν, καὶ ὀδύνης κατὰ γαστέρα ἐν οἴνῷ γλυκεῖ, καὶ ἐλκῶν τῶν μὲν ὑγρῶν ξηρὰν τῶν δὲ ξηρῶν ἐν μέλιτι. αὐται μὲν οὖν ταύτας ἔχουσι διαφοράς τε καὶ δυνάμεις.
- 4 "Αλλα δὲ πανάκη τὸ μὲν λεπτόφυλλον τὸ δὲ οὔ αἱ δὲ δυνάμεις ἀμφοῖν αἱ αὖταί, πρόσθετόν τε γυναιξὶ καὶ κατάπλασμα μετ' ἀλφίτου καὶ πρὸς τὰ ἔλκη τὰ ἄλλα καὶ πρὸς τὰ νεμόμενα.
- 5 Συνώνυμοι δέ καὶ οί στρύχνοι καὶ οί τιθύμαλ-

¹ κεφαλαίαs conj. Sch.; κεφαλη̂s Ald.

² τῶν ἐλκῶν conj. Sch.; τῶν δὲ ἐλκώδῶν(sic) U*; τῶν ἐλκωδῶν Ald H. cf. S 3.

Ald. H. cf. § 3.

8 Plin. 25. 32; Diosc. 3. 48.

δίκραν ἡ τρίκραν conj. Sch.; δίκραν ἡ τρίκαν UM; δικρανῆ πικράν U*; δικρανῆ ἡ τρικρανῆ Ald.

ENQUIRY INTO PLANTS, IX. xi. 2-5

it against bites of reptiles, to take it in a posset of honey for disorders of the spleen, when the blood collects about it, and against headache 1 to pound it up in olive-oil and anoint the head; that it is of use also in other obscure troubles, and against stomachache, if scraped and taken in wine. It is said also to be able to prevent long periods of sickness. Again for running sores 2 one may sprinkle it on in hot wine, first washing the place, while for dry sores one may soak it in wine and apply a plaster.

The kind named after Herakles has a large broad leaf, three spans each way, a root as thick as a man's finger, forking in two or three 4; in taste it is somewhat bitter, in smell like pure frankincense 5; 6 it is good to drink it against epilepsy, mixed with the rennet of a seal in the proportion of one to four, or in sweet wine against pain 7 in the stomach; it may be used dry 8 for running sores, and mixed with honey for dry ones. Such are the special features about these plants and their respective virtues.

⁹ There are also other kinds of all-heal, of which one has a fine leaf, the other not; the properties of both kinds are the same; namely they are used as a pessary for women, and a plaster may be made of them mixed with meal for spreading sores as well as

for ordinary sores.

Of the various plants called strykhnos.

As to strykhnos again and tithymallos (spurge) there is in either case more than one form of the plant

7 δδύνης conj. W.; δδύναι UMU*Ald.

δ λιβανωτοῦ καθαροῦ conj. Sch.; λιβανωτὸν καθαρὸν UM; λιβανωτοῦ U*; λιβανωτοῦ καθαρῶν Ald.H καθαροῦ perhaps due to καθάπερ. 6 cf. Fr. 175; Diosc. 2. 75.

⁸ ξηράν conj. Sch.; ξηρά U*Ald.; ξηραί Μ. Plin. 35. 33.

λοι. των γάρ στρύχνων ό μεν ύπνώδης ό δε μανικός, και ό μεν ύπνώδης ερυθράν έγων την ρίζαν ώσπερ αίμα Επραινομένην, ορυττομένην δέ . λευκήν. καὶ καρπὸν ἐρυθρότερον κρόκου, φύλλον δὲ τιθυμάλλω ὅμοιον ἡ μηλέα τῆ γλυκεία καὶ αὐτὸ δασὺ καὶ πυγμὴν μέγα. τούτου τῆς ῥίζης τον φλοιον κόπτοντες λίαν και βρέχοντες έν οίνω άκράτω διδόασι πιείν καὶ ποιεί καθεύδειν. Φύεται δὲ ἐν χαράδραις καὶ τοῖς μνήμασιν.

Ο δε μανικός, οί δε θρύορον καλοῦσιν αὐτὸν οί δὲ περιττόν, λευκὴν ἔχει τὴν ρίζαν καὶ μακράν ώς πήγεως καὶ κοίλην. δίδοται δ' αὐτης, ἐὰν μὲν ώστε παίζειν και δοκείν έαυτω κάλλιστον είναι. δραχμή σταθμώ εάν δε μάλλον μαίνεσθαι καί φαντασίας τινάς φαίνεσθαι, δύο δραγμαί έαν δ' ώστε μη παύεσθαι μαινόμενον τρείς, καλ συμπαραμιγνύναι φασίν όπον κενταυρίου εάν δέ **ωστε** ἀποκτείναι, τέτταρες. ἔχει δὲ τὸ μὲν φύλλον δμοιον εὐζώμω πλην μεῖζον, τὸν δὲ καυλὸν ὥσπερ οργυίας, κεφαλην δε ωσπερ γηθύου μείζω δε καί δασυτέραν εοικε δε καὶ πλατάνου καρπώ.

Cοντα: κόκκου MSS.

⁴ Plin. 21. 177-179; Diosc. 4. 73.

¹ cf. 7. 15. 4, where a third στρύχνος is mentioned, which is δμώνυμος, not συνώνυμος, i.e. which has nothing in common with these two στρύχνοι except the name. cf. also 9. 15. 5.
² κρόκου conj. Dalec. from Diosc. 4. 72, καρπόν . . . κροκί-

^{*} πυγμήν μέγα U; πυθμήν μέγας U*Ald. H.; W. adopts Bod.'s conjecture σπιθαμήν μέγα.

⁵ θρύορον Ald. Η.; θρυόρον U*; βρυόρον U; βρύορον MmBas.;

ENQUIRY INTO PLANTS, IX. x1. 5-6

denoted by the name. ¹Of the plants called strykhnos one induces sleep, the other (thorn-apple) causes madness. The first-mentioned has a root which becomes red like blood as it dries, but when first dug up it is white; its fruit is a deeper orange than saffron,² its leaf like that of tithymallos or the sweet apple; and it is itself rough, and about a foot high.³ The 'bark' of the root of this they bruise severely, and soaking it in neat wine give it as a draught, and it induces sleep. It grows in water-courses and on tombs.

⁴ The kind which produces madness (which some call thryoron ⁵ and some peritton ⁶) has a white hollow root about a cubit long. Of this three twentieths of an ounce in weight is given, if the patient is to become merely sportive and to think himself a fine fellow; twice this dose ⁷ if he is to go mad outright and have delusions ⁸; thrice the dose, if he is to be permanently insane; (and then they say that the juice of centaury is mixed with it); four ⁹ times the dose is given, if the man is to be killed. The leaf is like that of rocket, but larger, the stem about a fathom long; the 'head' ¹⁰ is like that of a long onion, but larger and rougher. And it also resembles the fruit of the plane-tree.

briorem G. Plin. l.c. seems to have read $\epsilon \rho \nu \theta \rho \delta \nu$; Diosc. l.c. $\beta \rho \delta \nu \nu$.

⁶ περιττὸν Ald.H., i.e. 'violent'; pissum G; Plin l.c. perisson; Diosc. l.c. πέρσων.

⁷ δραχμαί conj. Sch.; δραχμάς Ald.

8 καl... φαίνεσθαι om. UM: ungrammatical, and possibly a gloss; but cf. Diosc. and Plin. l.c.

τέτταρεs conj. Sch.; τέτταραs Ald ; τέσσαραs U*.

 10 7. 4. 10 it was said that $\gamma h\theta vov$ has no 'head,' i.e. bulb; here the 'head' seems to be the inflorescence. cf. Diosc. and Plin. l.c.

273

VOL. II.

Digitized by Google

T

Τῶν δὲ τιθυμάλλων ὁ μὲν παράλιος καλούμενος κόκκινον φύλλον ἔχει περιφερές, καυλὸν δὲ
καὶ τὸ ὅλον μέγεθος ὡς σπιθαμῆς τὸν δὲ καρπὸν
λευκόν. ἀμᾶται δὲ ὅταν ἄρτι περκάζη σταφυλή,
καὶ ξηρανθεὶς ὁ καρπὸς δίδοται πίνειν τριφθεὶς
ὅσον τρίτον μέρος ὀξυβάφου.

΄ ΄ Ο δ΄ ἄρρην καλούμενος τὸ μὲν φύλλον ἐλαιῶδες ἔχει, τὸ δὲ ὅλον μέγεθος πηχυαῖον. τοῦτον ὀπίζουσιν ἄμα τρυγητῷ καὶ θεραπεύσαντες οὕτως

ώς δεί χρώνται καθαίρει δὲ κάτω μάλλον.

Ο δε μυρτίτης καλούμενος τιθύμαλλος λευκός. τὸ μὲν φύλλον ἔχει καθάπερ ὁ μύρρινος, πλην ἀκανθῶδες ἀπ' ἄκρου· κλήματα δ' ἀφίησιν ἐπὶ την γην ώς σπιθαμιαία, ταθτα δ' ούχ άμα φέρει τον καρπον άλλα παρ' έτος, τα μεν νυν τα δ' είς νέωτα, πεφυκότα ἀπὸ της αὐτης ρίζης. φιλεί δὲ όρεινα χωρία. ὁ δὲ καρπὸς αὐτοῦ καλεῖται κάρυον. άμῶσι δ' ὅταν ἀδρύνωνται αἱ κριθαὶ καὶ ξηραίνοντες και άποκαθαίροντες αὐτὸν τὸν καρπου πλύναντες εν ύδατι και πάλιν ξηράναντες διδόασι πιείν συμμιγνύντες δύο μέρη της μελαίνης μήκωνος, τὸ δὲ συναμφότερον ὅσον ὀξύβαφον καθαίρει δὲ φλέγμα κάτω ἐὰν δὲ τὸ κάρυον αὐτὸ διδώσι, τρίψαντες έν οἴνω γλυκει διδόασιν ή έν σησάμφ πεφρυγμένφ κατατραγείν. ταῦτα μὲν οὖν τοῖς τε φύλλοις καὶ τοῖς όποῖς καὶ τοῖς καρποις γρήσιμα.

¹ Plin. 26. 68.

 ² κόκκινον conj. W.; κόκκος MSS. cf. Plin. l.c. ramis rubentibus.
 3 Diosc. 4. 164; Plin. 26. 62-65.

δύτως ώς δεί χρ. conj. Sch.; σύτως ώς δή χρ. U*; ούτως χρ. Ald.

ENQUIRY INTO PLANTS, IX. xi. 7-9

Of the various kinds of tithymallos.

1 Of the various plants called *tithymallos* (spurge) that which is called sea-spurge has a round scarlet ² leaf; the stem (and the size of the plant generally) is about a span long, and the fruit is white. It is gathered when the grape is just turning, and the dried fruit is given in a draught, the dose being the twenty-fourth part of a pint.

³ That which is called the 'male' has a leaf like the olive, and the height of the whole plant is a cubit. Of this they collect the juice at the time of vintage, and, after preparing it, use it as occasion demands 4; and it purges chiefly downwards.

⁵The kind of tithymallos called 'myrtle-like' is white; it has a leaf like the myrtle, but spinous at the tip; it puts out earthward twigs about a span long, and these bear the fruit of not all at the same time but in alternate years, so that from the same root grow fruits partly this and partly next year. It loves hill-country. The fruit of it is called a 'nut.' They gather it when the barley is ripening and dry and clean it; (it is the actual fruit which they clean); they wash it in water and, after drying it again, give it in a draught, mixing with it two parts of 'black poppy'; and the whole dose amounts to about an eighth of a pint. It purges phlegm downwards. If they administer the 'nut' itself, they first pound it up in sweet wine, or give it in parched sesame to bite up. These plants then have leaves juices or fruits which are serviceable.

⁵ Diosc. 4, 164; Plin. 26, 66, ⁶ cf. C.P. 4, 6, 9.

⁷ W. adds δὲ after αὐτὸν. The treatment of the leaves has perhaps dropped out. cf. Plin. l.c. (3's version is even shorter. 8 μέλαινα must here mean 'dark,' i.e. red. See Index.

Τῶν δὲ λιβανωτίδων, δύο γάρ εἰσιν, ἡ μὲν ἄκαρπος ἡ δὲ κάρπιμος, ἡ μὲν καὶ τῷ καρπῷ καὶ τῷ φύλλῷ χρησίμη ἡ δὲ μόνον τἢ ῥίζη. καλεῖται δὲ ὁ καρπὸς κάχρυ. ἔχει δὲ αὕτη τὸ μὲν φύλλον ἐοικὸς σελίνῷ ἑλείφ μείζον δὲ πολύ, καυλὸν δὲ μέγεθος πήχεως ἡ μείζω, ῥίζαν δὲ μεγάλην καὶ παχεῖαν λευκὴν ὄζουσαν ὅσπερ λιβανωτοῦ, καρπὸν δὲ λευκὸν τραχὸν προμήκη· φύεται δὲ μάλιστα ὅπου ἀν αὐχμηρὰ χωρία ἢ καὶ πετρώδη· χρησίμη δὲ ἡ μὲν ῥίζα πρός τε τὰ ἔλκη καὶ πρὸς τὰ γυναικεῖα πινομένη ἐν οἴνῷ αὐστηρῷ μέλανι· ὁ δὲ καρπὸς πρός τε τὰς στραγγουρίας καὶ πρὸς τὰ ὅτα καὶ ἄργεμα καὶ πρὸς ὀφθαλμίας καὶ ὥστε γάλα γυναιξὶν ἐμποιεῖν.

'Η δὲ ἄκαρπος ἔχει τὸ φύλλον ὅμοιον θριδακίνης τῆς πικρᾶς τραχύτερον δὲ καὶ λευκότερον, ρίζαν δὲ βραχεῖαν. φύεται δὲ ὅπουπερ ἐρείκη πλείστη. δύναται δὲ ἡ ρίζα καθαίρειν καὶ ἄνω καὶ κάτω τὸ μὲν γὰρ πρὸς τὴν βλάστην ἄνω, τὸ δὲ πρὸς τὴν γῆν κάτω κωλύει δὲ καὶ εἰς ἱμάτια τιθεμένη τοὺς σῆτας. συλλέγεται δὲ περὶ πυρο-

τομίας.

ΧΙΙ. Χαμαιλέων δὲ ὁ μὲν λευκὸς ὁ δὲ μέλας αἱ δὲ δυνάμεις τῶν ριζῶν καὶ αὐταὶ δὲ αἱ ρίζαι τοῖς εἴδεσι διάφοροι. τοῦ μὲν γὰρ λευκὴ καὶ παχεῖα καὶ γλυκεῖα καὶ ὀσμὴν ἔχουσα βαρεῖαν χρήσιμον δέ φασι πρός τε τοὺς ροῦς, ὅταν ἐψηθῆ

* cf. 7. 6. 2; 9. 9. 5.

Diosc. 3. 74; Plin 19. 187.

² cf. Plin. 24. 99 and 101.

⁴ δπουπερ έρείκη conj. Dalec. etc. from Diosc. l.c.; δπουπερ εγρηται Ald.H.; δπότε έρείκη U*.

ENQUIRY INTO PLANTS, IX. xi. 10-xii. 1

Of the two herbs called libanotis.

¹ Of the plants called libanotis, (for there are two) one is barren, the other fruitful, the latter having both fruit and leaves that are serviceable, the former only a serviceable root. The fruit is called kakhry.² This plant has a leaf like marsh celery, but much larger, a stem a cubit long or more, a large stout white root, which smells like frankincense, and a white rough elongated fruit. It grows chiefly wherever there is parched and rocky soil; the root is serviceable for sores, and for diseases of women when given in a draught of dry black wine. The fruit is good for strangury, for the ears, for ulcers ³ on the eye, for ophthalmia and for producing milk in women.

The barren kind has a leaf like that of the bitter

The barren kind has a leaf like that of the bitter lettuce, but rougher and paler; the root is short. It grows where there is abundance of heather.⁴ The root can purge both upwards and downwards, the upper part being used for the former, that nearer the ground for the latter purpose. Also, if it is put among clothes, it prevents moth. It is gathered at

the time of wheat-harvest.

Of the two kinds of chamaeleon.

XII. ⁵Of chamaeleon there is the white kind and the dark; the properties of the roots are different, and the roots also differ ⁶ in appearance. In the one case the root is white stout and sweet, and it has a heavy smell; they say that when cooked it is serviceable against flux; it is chopped up like

΄ ⁶ διαφέρουσαι conj. W.; διαφέρει U; διαφέρουσι δὲ M; διάφορον Ald.



⁵ Diosc. 3. 8; Plin. 22. 45 and 46, who explains the name (mutat cum terra colores). See Index.

κατατμηθείσα καθάπερ ραφανὶς ἐνειρομένη ἐφ όλοσχοίνου, καὶ πρὸς τὴν ἔλμινθα τὴν πλατείαν, ὅταν ἀσταφίδα προφάγῃ πίνειν ἐπιξύοντα ταύτην ὅσον ὀξύβαφον ἐν οἴνῷ αὐστηρῷ. ἀναιρεῖ δὲ καὶ κύνα καὶ σῦν· κύνα μὲν ἐν ἀλφίτοις ἀναφυραθεῖσα μετὰ ἐλαίου καὶ ὕδατος, σῦν δὲ μετὰ ραφάνων μεμιγμένη τῶν ὀρείων. γυναικὶ δὲ δίδοται ἐν τρυγὶ γλυκεία ἢ ἐν οἴνῷ γλυκεί. καὶ ἐὰν βούληταί τις ἀσθενοῦντος ἀνθρώπου διαπειρασθαι εἰ βιώσιμος, λούειν κελεύουσι τρεῖς ἡμέρας, κὰν περιενέγκῃ βιώσιμος. φύεται δὲ ὁμοίως πανταχοῦ, καὶ ἔχει τὸ φύλλον ὅμοιον σκολύμῷ μεῖζον δέ· αὐτὸ δὲ πρὸς τῇ γῇ τινα κεφαλὴν ἔχει ἀκανοειδῆ μεγάλην, οἱ δὲ καὶ ἄκανον καλοῦσιν.

΄Ο δὲ μέλας τῷ μὲν φύλλῳ παρόμοιος, σκολυμώδες γὰρ ἔχει πλὴν ἔλαττον καὶ λειότερον, αὐτὸς δ΄ ὅλος ἐστὶν ὥσπερ σκιάδιον, ἡ δὲ ρίζα παχεῖα καὶ μέλαινα διαρραγεῖσα δὲ ὑπόξανθος. χωρία δὲ φιλεῖ ψυχρὰ καὶ ἀργά· δύναται δὲ λέπραν τε ἐξελαύνειν ἐν ὅξει τριβόμενος καὶ ξυσθει ἐπαλειφόμενος καὶ ἀλφὸν ὡσαύτως· ἀναιρεῖ

δὲ καὶ τοὺς κύνας.

s Μήκωνες δ' εἰσὶν ἄγριαι πλείους· ή μὲν κερατῖτις καλουμένη μέλαινα· ταύτης τὸ φύλλον ὥσ-

² cf. Pseudo-Diosc. 4. 175 and Index.

* δέ after την om. Sch.; ? τινα κεφαλην W.

¹ cf. 9. 9. 1.

³ ἀκανοειδή conj. Sch.; κονοειδή U*; κωνοειδή mP; δμοίαν ἀκανώ PAld.

²⁷⁸

ENQUIRY INTO PLANTS, IX. XII. 1-3

radishes 1 and the pieces strung on a rush; it is also good against the broad maw-worm; the patient first eats a bunch of raisins and then drinks about an eighth of a pint of this scraped up in a draught of dry wine. It is fatal to dogs and pigs; to kill a dog it is well mixed up in a meal paste with oil and water, to kill a pig it is mixed with 'mountain cabbage (spurge).2 It is given to a woman in sweet winelees or sweet wine. And if one wishes to discover whether a man that is sick will recover, they say that he should be washed with this for three days, and, if he survives the experience, he will recover. It grows everywhere alike and has a leaf like the golden thistle, but larger; the plant itself has a large thistle-like 3 head 4 close to the ground; some actually 5 call it the thistle.

⁶ The dark kind resembles the other in leaf, which is like that of the golden thistle but smaller and smoother; the plant itself is in general appearance like a sunshade; the root is stout and black, and when broken is yellowish. It likes cold uncultivated soil: it has the property of expelling leprosy; for this it is given pounded up in vinegar, or else scrapings of it are made into a plaster; and it is also used for the white leprosy. This plant is also fatal

to dogs.7

Of the various plants called 'poppy.'

⁸ There are several kinds of wild poppy: the one called the horned poppy is black: the leaf of this is

6 Diose. 3. 9; Plin. l.c.

⁸ Diosc. 4. 64; Plin. 20. 205 and 206.

⁵ δè καὶ ἄκανον I conj. ; δ' ἄκανθαν U*mPar., so also Diosc. l.c.; δ' ἄκανον PAld.G.

⁷ κύνας: κυνορραίστας, dog-ticks, conj. Reinesius from Plin. ricinos canum.

περ φλόμου της μελαίνης ήττον δὲ μέλαν, τοῦ δὲ καυλοῦ τὸ ὕψος ὡς πηχυαῖον, ῥίζα δὲ παχεῖα καὶ ἐπιπόλαιος, ὁ δὲ καρπὸς καμπύλος ὥσπερ κεράτιον συλλέγεται δὲ περὶ πυροτομίας. δύναται δὲ καθαίρειν κοιλίαν, τὸ δὲ φύλλον ἄργεμα προβάτοις ἀφαιρεῖν. φύεται δὲ παρὰ θάλατταν, οῦ ἄν ἢ πετρώδη χωρία.

Έτέρα δὲ μήκων ροιὰς καλουμένη παρομοία κιχορίω τῷ ἀγρίω, δι δ καὶ ἐσθίεται· ἐν τοῖς ἀρουραίοις δὲ φύεται, μάλιστα ἐν ταῖς κριθαῖς· ἄνθος δ' ἔχει ἐρυθρὸν κωδύαν δ' ὅσην ὅνυχα τοῦ δακτύλου. συλλέγεται δὲ πρὸ τοῦ θερισμοῦ τῶν κριθῶν, ἐγχλωροτέρα δὲ μᾶλλον. καθαίρει δὲ

κάτω.

΄Ετέρα δὲ μήκων Ἡρακλεία καλεῖται τὸ μὲν φύλλον ἔχουσα οἶον στρουθός, ῷ τὰ ὀθόνια λευκαίνουσι, ῥίζαν δὲ λεπτὴν ἐπιπόλαιον, τὸν δὲ καρπὸν λευκόν. ταύτης ἡ ῥίζα καθαίρει ἄνω χρῶνται δὲ τινες πρὸς τοὺς ἐπιλήπτους ἐν μελικράτω. ταῦτα μὲν οὖν ὥσπερ ὁμωνυμία τινὶ συνείληπται.

ΧΙΙΙ΄. Των δὲ ριζων καὶ ἐν τοῖς χυμοῖς αἱ διαφοραὶ καὶ ἐν ταῖς ὀσμαῖς· αἱ μὲν γάρ εἰσι δριμεῖαι αἱ δὲ πικραὶ αἱ δὲ γλυκεῖαι, καὶ αἱ μὲν εὐοδμοι αἱ δὲ βαρεῖαι. γλυκεῖα μὲν ἢ τε νυμφαία καλουμένη· φύεται δ' ἐν ταῖς λίμναις καὶ ἐν τοῖς ἑλώδεσιν, οῖον ἔν τε τἢ 'Ορχομενία καὶ Μαραθωνι

 $^{^1}$ ἄσπερ κεράτιον conj. W.; ἄσπερ κέρας UM; ἄσπερ τῶν κερατίων U*Ald.

Diosc. l.c.; Plin. 19. 167-169.
 Diosc. 4. 66; Plin. 20. 207, cf. 19. 21.

²⁸⁰

ENQUIRY INTO PLANTS, IX. XII. 3-XIII. I

like that of the black mullein, but it is not so black; the stem grows about a cubit high, the root is stout and shallow, the fruit is twisted like a little horn1: it is gathered at the time of wheat harvest. It has the property of purging the belly, and the leaf is used for removing ulcers on sheep's eyes. It grows by the sea, wherever there is rocky ground.

²·Another kind of poppy is that called rhoias, which is like wild chicory, wherefore it is even eaten: it grows in cultivated fields and especially among barley. It has a red flower, and a head as large as a man's finger-nail. It is gathered before the barley-harvest, when it is still somewhat green.

It purges downwards.

§ Another kind of poppy is called Herakleia: it has a leaf like soap-wort, with which 4 they bleach linen: the root is slender and does not run deep, and the fruit is white. The root of this plant purges upwards: and some use it in a posset of mead for epileptics.

⁵ These kinds then are distinct plants, though they

come under one name.

Of roots possessing remarkable taste or smell.

XIII. The differences between roots are shown in their tastes 6 and in their smells: some are pungent, some bitter, some sweet: some again have a pleasant, others a disagreeable smell. The plant called yellow water-lily 7 is sweet: it grows in lakes and marshy places, as in the district of Orchomenus, at

⁴ This appears to refer to στρουθός, not to 'Ηρακλεία, as

⁶ χυμοῖς conj. Sch.; χυλοῖς Ald, ⁷ Plin. 25, 75.

28 T

Plin. takes it. cf. 6. 4. 3 and Index, στρούθιον (2).

i.e. ροιάς and Ἡρακλεία are popularly called 'poppies.' δμωνυμία τινι conj. W.; δμώνυμα τινα Ald.; δμώνυμα τινι U*. cf. 7. 15. 4.

καὶ περὶ Κρήτην καλοῦσι δ' αὐτὴν οἱ Βοιωτοὶ μαδωνάϊν και τον καρπον έσθίουσιν. έχει δε το φύλλον μέγα έπι του υδατος είναι δέ φασιν ίσχαιμου, εαν τρίψας τις επί την πληγην έπιθη.

χρησίμη δὲ καὶ πρὸς δυσεντερίαν πινομένη.

Γλυκεία δε και ή Σκυθική και ένιοι δε καλοῦσιν εὐθὺς γλυκεῖαν αὐτήν· γίνεται δὲ περὶ τὴν Μαιῶτιν· χρησίμη δὲ πρός τε τὰ ἄσθματα καὶ πρὸς τὴν βῆχα ξηρὰν καὶ ὅλως τοὺς περὶ τὸν θώρακα πόνους έτι δε προς τὰ έλκη έν μέλιτι δύναται δὲ καὶ τὴν δίψαν παύειν, ἐάν τις ἐν τῷ στόματι έχη· δι' δ ταύτη τε καὶ τῆ ἱππακῆ διάγειν φάσι τους Σκύθας ημέρας και ένδεκα και δώδεκα.

[Ή δὲ ἀριστολοχία τῆ ὀσφρήσει μὲν εὔοδμος τῆ δὲ γεύσει πικρὰ σφόδρα τῆ χροιᾶ δὲ μέλαινα. φύεται δε εν τοις δρεσιν ή βελτίστη φύλλον δε έχει προσεμφερές τῆ ἀλσίνη πλὴν στρογγυλό-τερον χρησίμη δὲ πρὸς πολλά, καὶ ἀρίστη πρὸς κεφαλής άγαθή δὲ καὶ πρὸς τὰ άλλα έλκη, καὶ πρός τὰ έρπετὰ καὶ πρὸς ὕπνον καὶ πρὸς ὑστέραν. τὰ μὲν οὖν προσάγειν κελεύουσιν ἐν ὕδατι ἀναδεύσαντα καὶ καταπλάττοντα, τὰ δὲ ἄλλα εἰς μέλι ἐνξύσαντα καὶ ἔλαιον πρὸς δὲ τὰ τῶν . έρπετῶν ἐν οἴνφ ὀξίνη πίνειν και ἐπι τὸ δῆγμα ἐπιπλάττειν εἰς ὕπνον δὲ ἐν οἴνφ μέλανι αὐστηρφ κνίσαι έαν δε αι μητραι προπέσωσι, τῷ ὕδατι ἀποκλύζειν.]

¹ Diose. 3, 5; Plin, 25, 82.

² γλυκείαν: γλυκύρριζαν conj. Dalec., whence 'liquorice.' cf. Plin. 11. 284.

^{*} cf. Plin. l.c., who took immany to be a plant. 282

ENQUIRY INTO PLANTS, IX. XIII. 1-3

Marathon and in parts of Crete: the Boeotians, who eat the fruit, call it madonais. It has a large leaf which lies on the water: and it is said that it acts as a styptic if it is pounded up and put on the wound: it is also serviceable in the form of a draught for

dvsenterv.

1 'Scythian root' (liquorice) is also sweet; some indeed call it simply 'sweet-root.' It is found about Lake Maeotis: it is useful against asthma or a dry cough and in general for troubles in the chest: also, administered in honey, for wounds: also it has the property of quenching thirst, if one holds it in the mouth: wherefore they say that the Scythians, with the help of this and mares' milk cheese s can go eleven or twelve days without drinking.

⁴[Birthwort is fragrant to the smell but in taste is very bitter: in colour it is black. The best grows on the mountains: it has a leaf like alsine, but rounder: it is useful for many purposes, and is best for sores on the head ⁵ and other sores, also for bites of reptiles, for inducing sleep and for disorders of the womb.⁶ It is directed that it should be applied as a plaster, steeped in water, and for the other purposes should be given shredded into honey and olive-oil: for snake-bites it should be taken in sour wine and also used as a plaster on the bite: to induce sleep it should be scraped up ⁷ and administered in black dry wine; in cases of prolapsus uteri a lotion of it mixed with water should be applied.]

κνίσαι conj. W.; κνίσας U*Ald.

⁴ Diosc. 3. 4; Plin. 25. 95. This section is repeated 9. 20. 4. with considerable variations: that seems to be its proper place.

^{*} κεφαλῆς conj. W.; κεφαλὴν Ald. cf. § 20, κεφαλάθλαστα.
* ὑστέραν conj. W., cf. below, ἐὰν δὲ αἰ μῆτραι κ.τ.λ. and the duplicate passage § 20; ἔτερα MSS.

Αὐται μὲν οὖν γλυκείαι. ἄλλαι δὲ πικραί, αί δὲ βαρεῖαι τῆ γεύσει. γίνονται δέ τινες τῶν γλυκειῶν αἱ μὲν ἐκστατικαί, καθάπερ ἡ ὁμοία τῷ σκολύμφ περὶ Τέγεαν, ἢν καὶ Πάνδειος ὁ ἀνδριαντοποιὸς φαγὼν ἐργαζόμενος ἐν τῷ ἱερῷ ἐξέστη. αἱ δὲ θανατηφόροι, καθάπερ ἡ περὶ τὰ μέταλλα ἐν τοῖς ἔργοις τοῖς ἐν Θράκη· κούφη δὲ καὶ ἡδεῖα πάνυ τῆ γεύσει καὶ τὸν θάνατον ὑπνώδη τινὰ ποιοῦσα καὶ ἐλαφρόν. ἔχουσι δὲ καὶ τοῖς χρώμασι διαφορὰς οὐ τῷ λευκῷ καὶ μέλανι καὶ ξανθῷ μόνον, ἀλλ ἔνιαι καὶ οἰνοχρῶτες, αἱ δ' ἐρυθραί, καθάπερ ἡ τοῦ ἐρευθεδανοῦ.

Ή δὲ τοῦ πενταφύλλου ἡ πενταπετοῦς, καλοῦσι γὰρ ἀμφοτέρως, ὀρυττομένη ἐρυθρὰ ξηραινομένη δὲ μέλαινα γίνεται καὶ τετράγωνος ἔχει δὲ τὸ φύλλον ὥσπερ οἴναρον μικρὸν δὲ καὶ τὴν χροιὰν ὅμοιον καὶ αὐξάνεται καὶ φθίνει ἄμα τῆ ἀμπέλω πάντα δὲ πέντε τὰ φύλλα, δι δ καὶ ἡ προσηγορία καυλοὺς δὲ ἐπὶ γῆν ἵησι λεπτοὺς καὶ κνήμας ἔχει.

Τὸ δὲ ἐρευθεδανὸν φύλλον ὅμοιον κιττῷ πλὴν στρογγυλότερον φύεται δ' ἐπὶ γῆς ὥσπερ ἄγρωστις, φιλεῖ δὲ παλίσκια χωρία. οὐρητικὴ δέ, δί δ καὶ χρῶνται πρὸς τὰ τῆς ὀσφύος ἀλγήματα καὶ πρὸς τὰς ἰσχιάδας.

Ένιαι δὲ ἰδιόμορφοί τινες, ὅσπερ ἤ τε τοῦ σκορπίου καλουμένου καὶ ἡ τοῦ πολυποδίου. ἡ

² cf. C.P. 6.4.5. ³ $\dot{\eta}$ conj. Sch.; ai U*Ald.

⁵ Plin. 25. 139.

¹ These words shew that § 3 is out of place.

^{*} ἐν τοῖς ἔργοις τοῖς W. from U*. ? a gloss on μέταλλα, τὰ μετ. τὰ ἐν Θ. Ald. H.

ENQUIRY INTO PLANTS, IX. xIII. 4-6

¹ These then are sweet: other roots are bitter, and some unpleasant to the taste. Of those that are sweet ² there are some that cause mental derangement, as the plant like the golden thistle which grows near Tegea: of this Pandeios the sculptor ate, and went mad while he was working in the temple. Others have fatal effects, as that ³ which grows near the mines in the fields of ⁴ Thrace: this however is inoffensive and quite sweet to the taste, and the death which it causes is easy and like falling asleep. There are also differences in colour, not merely as to being black or white or yellow, but some are quite wine-coloured and some are red, as the root of madder.

⁵ The root of pentaphyllon or pentapetes (cinquefoil) ⁶ (for the plant bears both names) is red when it is dug up, but as it dries it becomes black and square: its leaf is like a vine-leaf, and it is small and like it in colour: it grows and fades along with the vine. It only has five leaves in all, whence its name: it sends out long slender stems on the ground, and it has

joints.7

⁸ Madder has a leaf like ivy, but it is rounder: it grows along the ground like dog's-tooth grass and loves shady spots. It has diuretic properties, wherefore it is used for pains in the loins or hip-disease.

Some roots are of peculiar shape, as that of the plant called 'scorpion-plant' (leopard's bane) 9 and that of polypody. For the former is like a scorpion

6 πενταπετοῦς conj. Sch.; πενταπέτου UAld.; πεντεπέτου M U*. cf. Diosc. 4. 42.

⁸ Diosc. 3. 143; Plin. 19. 47.

⁹ cf. 9. 18. 2.

⁷ καὶ κνήμας έχει U*; και κν. ἐ. πυκνάς Ald.; καὶ κνίκας έχει πυκνάς UM. cf. πολύκνημος, Diosc. 3. 94. Text probably defective, as nothing is said of the plant's medicinal use.

μὲν γὰρ ὁμοία σκορπίφ καὶ χρησίμη δὲ πρὸς τὴν πληγὴν αὐτοῦ καὶ πρὸς ἄλλ ἄττα. ἡ δὲ τοῦ πολυποδίου δασεῖα καὶ ἔχουσα κοτυληδόνας, ὥσπερ αἱ τοῦ πολύποδος πλεκτάναι. καθαίρει δὲ κάτω κὰν περιάψηταί τις οῦ φασιν ἐμφύεσθαι πολύπουν. ἔχει δὲ φύλλον ὅμοιον τῆ πτερίδι τῆ

μεγάλη καὶ φύεται ἐν ταῖς πέτραις.

ΧΙΥ. Πασῶν δὲ τῶν ριζῶν αἱ μὲν πλείω γρόνον αί δὲ ἐλάττω διαμένουσιν. ὁ μὲν γὰρ έλλέβορος καὶ τριάκοντα έτη χρήσιμος, ή δὲ άριστολοχία πέντε ή έξ, χαμαιλέων δε ό μέλας τετταράκοντα, κενταυρίς δὲ δέκα ἡ δώδεκα πίειρα δὲ ἡ ρίζα καὶ πυκνή· πευκέδανον δὲ πέντε ή έξ, άμπέλου δὲ ἀγρίας ἐνιαυτόν, ἐὰν ἐν σκιᾶ ἦ καὶ ἄπληκτος, εἰ δὲ μή, σαπρὰ καὶ σομφώδης. άλλαι δὲ άλλους ἔχουσαι χρόνους. πάντων δὲ όλως των φαρμάκων πλείστον διαμένει χρόνον τὸ έλατήριον, καὶ τὸ παλαιότατον ἄριστον. ἰατρὸς δ' οὖν τις ἔλεγεν οὖκ ἀλαζών οὖδὲ ψεύστης ώς εἴη παρ' αὐτῶ καὶ διακοσίων ἐτῶν θαυμαστὸν δὲ τῆ 2 ἀρετή, δοῦναι δὲ αὐτῷ τινα δῶρον. αἰτία δὲ τής χρονιότητος ή ύγρότης. διὰ γὰρ ταύτην καὶ ὅταν κόψωσι τιθέασι είς τέφραν ύγρόν, καὶ οὐδ' ως γίνεται ξηρόν, άλλ' ἄχρι πεντήκοντα έτῶν σβέννυσι προσαγομενον τούς λύχνους. φασί δε μόνον

² Diosc. 4. 186.

¹ cf. the mediaeval doctrine of 'signatures.'

³ τις οὐ conj. Sch.; τις ὡς Ald.; τίς UM; τίς ὥς U*.
⁴ Plin. 27. 143.
⁵ cf. 9. 8. 7.
⁶ cf. 9. 20. 3.

⁷ ἄπληκτος: ? by worms. cf. ἄκοπος.

ENQUIRY INTO PLANTS, IX. xIII. 6-XIV. 2

and is also useful 1 against the sting of that creature and for certain other purposes. ² The root of polypody is rough and has suckers like the tentacles of the polyp. It purges downwards: and, if one wears it as an amulet, they say that one 3 does not get a It has a leaf like the great fern, and it grows on rocks.

Of the time for which roots can be kept without losing their virtue.

XIV. 4 Some roots keep a longer, some a shorter time. Hellebore retains its usefulness for as much as thirty years, birthwort five or six, the black chamaeleon for forty, feverwort ⁵ (whose root is thick and compact) for ten or twelve. Sulphur-wort keeps five or six years, the root of the 'wild vine' (bryony) for a year, if it be kept in the shade and not damaged: 7 otherwise it rots and becomes spongy.8 Others keep for various periods. But, to speak generally, of all plants used as drugs the 'driver' keeps longest, and, the older it is, the better it is. At least a certain physician, who was no boaster nor liar, said that he had some which was 200 years old and of marvellous virtue, and that it was a present to him from some one. The cause of its keeping so long is its moisture: 10 for to secure this, as soon as they have cut it, they put it among ashes without drying it, and not even so does it become dry, but up to fifty years it will put the lamp 11 out if it is brought near it. And they say that alone of all

⁸ σομφώδης conj. Sch.; σογκώδης Ald. Η. ⁹ A manufactured drug. cf. 9. 9. 4.

Diosc. 4. 150; Plin. 20. 5.
 λύχνους conj. Sch.: so Vin.Cod.Cas.GPlin. l.c.; αὐχμοὺς U*Ald.: Yodvous UM.

η μάλιστα υπέρινον ανω ποιείν των φαρμάκων αυτη μεν ουν ιδιότης τις δυνάμεως.

- Τῶν δὲ ῥιζῶν ὅσαι μὲν γλυκύτητά τινα ἔχουσι ξυμβαίνει θριπηδέστους γίνεσθαι χρονιζομένας, ὅσαι δὲ δριμεῖαι, τοῦτο μὲν μὴ πάσχειν ἀμαυροῦσθαι δ΄ αὐτῶν τὰς δυνάμεις μανουμένων καὶ κενουμένων. τῶν δ΄ ἔξω θηρίων ἄλλο μὲν οὐδὲν ἄπτεται ῥίζης δριμείας, ἡ δὲ σφονδύλη πασῶν τοῦτο μὲν οὖν ἴδιον τῆς τοῦ ζώου φύσεως.
- Πασαν δὲ χείρω γίνεσθαι ῥίζαν, ἐὰν ἐάση τις τελειωθηναι καὶ άδρυνθηναι τὸν καρπόν ὡσαύτως δὲ καὶ τὸν καρπόν, ἐὰν ὀπίσης τὴν ῥίζαν ὡς ἐπὶ τὸ πολὺ δὲ αὶ φαρμακώδεις οὐκ ὀπίζονται, ὧν δ' ἀν τὰ σπέρματα φαρμακώδη, αὖται δὴ ὀπίζονται χρησθαι δέ τινές φασι μᾶλλον ταις ῥίζαις, ὅτι ἰσχυρότερος ὁ καρπὸς ὥσθ' ὑπενεγκεῦν τὸ σῶμα. φαίνεται δὲ οὐ καθ' ὅλου τοῦτο ἀληθές ἐπεὶ καὶ οἱ ἐν 'Αντικύρα τοῦ σησαμώδους [ἐλλεβόρου] διδόασιν, [ὅτι ὁ καρπὸς ὅμοιος σησάμω].

ΧV. Φαρμακώδεις δὲ δοκοῦσιν εἶναι τόποι μάλιστα τῶν μὲν ἔξω τῆς Ἑλλάδος οἱ περὶ τὴν Τυρρηνίαν καὶ τὴν Λατίνην, ἐν ἢ καὶ τὴν Κίρκην εἶναι λέγουσιν· καὶ ἔτι μᾶλλόν γε, ὡς "Ομηρός

¹ Plin. 27. 143.

² i.e. not engendered in the root.

<sup>A beetle? cf. Arist. H.A. 5. 8.
This section is omitted in U*. Plin. 27. 144.</sup>

ENQUIRY INTO PLANTS, IX. xiv. 2-xv. 1

drugs, or to a greater degree than any, it effects a thorough purge upwards: this then is a virtue peculiar to it.

Those roots which contain any sweetness become worm-eaten in course of time, but those that are pungent are not so affected, though their virtues diminish as they become flabby and waste away.

No creature coming from without 2 touches a pungent root, but the sphondyle 3 attacks them all;

this then is a peculiarity of this creature.

⁴Any root, they say, deteriorates if one lets the fruit grow to maturity and ripen: and so in like manner does the fruit, if you drain the root of its juice: and in general roots with medicinal properties do not have the juice of their roots taken, and only those whose seeds are medicinal are thus treated. But some say that they use the roots for choice, because the fruit is too powerful for the human body to be able to bear it. However this does not appear to be true as a universal rule, seeing that the people of Anticyra administer ⁵ doses of the drug ⁶ sesamodes made from hellebore, which is so called because its fruit is like sesame.

Of the localities which specially produce medicinal herbs.

XV. The places outside Hellas which specially produce medicinal herbs seem to be the parts of Tyrrhenia and Latium (where they say that Circe dwelt), and still more parts of Egypt, as Homer says:

⁵ i.e. and it is in this case the fruit which is used. The drug in question, as well as the plant, was called σησαμοειδές or σησαμοειδής. cf. 9. 9. 2 n.; Diosc. 4. 149.

⁶ Or (if $\epsilon \lambda \lambda \epsilon \beta \delta \rho \rho \nu$ is sound) of the sesame-like hellebore, i.e. he black. δτι... σησάμφ I have bracketed, as a gloss on σησαμώδους: $\epsilon \lambda \lambda \epsilon \beta \delta \rho \rho \nu$ is probably also a gloss.

289

VOI., II.

T7

φησι, τὰ περὶ Αἴγυπτον ἐκείθεν γὰρ τὴν Ἑλένην φησὶ λαβεῖν "ἐσθλὰ τά οἱ Πολύδαμνα πόρεν Θῶνος παράκοιτις Αἰγυπτίη τόθι πλεῖστα φύει ζείδωρος ἄρουρα φάρμακα, πολλὰ μὲν ἐσθλὰ τετυγμένα πολλὰ δὲ λυγρά." ὧν δὴ καὶ τὸ νηπενθὲς ἐκεῖνό φησιν εἰναι καὶ ἄχολον, ὥστε λήθην ποιεῖν καὶ ἀπάθειαν τῶν κακῶν. καὶ σχεδὸν αὖται μὲν ἐοίκασιν ὥσπερ ὑπὸ τῶν ποιητῶν ὑποδεδεῖχθαι. καὶ γὰρ Αἰσχύλος ἐν ταῖς ἐλεγείαις ὡς πολυφάρμακον λέγει τὴν Τυρρηνίαν "Τυρρηνὸν γενεάν, φαρμακοποιὸν ἔθνος."

Οἱ δὲ τόποι πάντες πως φαίνονται μετέχειν τῶν φαρμάκων, ἀλλὰ ἡτῷ μᾶλλον καὶ ἡττον διαφέρειν καὶ γὰρ οἱ πρὸς ἄρκτον καὶ μεσημβρίαν καὶ οἱ πρὸς ἀνατολὰς ἔχουσι θαυμαστὰς δυνάμεις. ἐν Αἰθιοπία γὰρ ἡ τοὺς ὀϊστοὺς χρίουσι ῥίζα τίς ἐστι θανατηφόρος. ἐν δὲ Σκύθαις αὕτη τε καὶ ἔτεραι πλείους, αἱ μὲν παραχρῆμα ἀπαλλάττουσαι τοὺς προσενεγκαμένους, αἱ δ' ἐν χρόνοις αἱ μὲν ἐλάττοσιν αἱ δ' ἐν πλείοσιν, ὥστ' ἐνίους καταφθίνειν. ἐν Ἰνδοῖς δὲ καὶ ἔτερα γένη πλείω, περιττότατα δέ, εἴπερ ἀληθῆ λέγουσιν, ἡ τε δυναμένη τὸ αἶμα διαχεῖν καὶ οἶον ὑποφεύγειν, καὶ πάλιν ἡ συνάγουσα καὶ πρὸς ἑαυτὴν ἐπισπωμένη, ἃ δή φασιν εὐρῆσθαι πρὸς τὰ τῶν ὀφιδίων τῶν θανατηφόρων δήγματα.

Περὶ δὲ τὴν Θράκην εἶναι μὲν καὶ ἐτέρας οὐκ ὀλίγας, ἰσχυροτάτην δὲ ὡς εἰπεῖν τὴν ἴσχαιμον, ἣν δὴ λέγουσιν οἱ μὲν κεντηθείσης τῆς Φλεβὸς

¹ Od. 4. 221 foll.

² ων δη conj. Sch.; ωί δη U*; ἐν οίς δη PAld.

ENQUIRY INTO PLANTS, IX, xv, 1-3

for thence he says 1 that Helen brought "things of virtue which Polydamna, the Egyptian wife of Thon, gave her; there the grain-bearing earth produces most drugs, many that are good, and many baneful." Among these 2 he says was nepenthes, the famous drug which cures sorrow and passion, so that it causes forgetfulness and indifference to ills. So these lands seem to have been pointed out, as it were, by the For Aeschylus too in his elegies speaks of Tyrrhenia as rich in drugs, for he tells of the "Tyrr-

henian stock, a nation that makes drugs."

It seems that almost all places take their share in producing drugs, but that they differ in the extent to which they do so; for the regions of the North, South, and East have herbs of marvellous virtue. Thus in Ethiopia there is a certain deadly root 3 with which they smear their arrows. And in Scythia there is this and there are also others, some of which kill at once those who eat them, some after an interval, shorter or longer, so that in the latter case men have a lingering death. In India there are many other kinds.4 but the most extraordinary,5 if they tell the truth, are these: there is one which has the power to make the blood disperse and as it were to put it to flight,6 and another which collects it and draws it to itself; these they say were discovered as remedies for the bites of deadly serpents.

In Thrace it is said there are fairly numerous other kinds, but that about the most powerful is 'blood-stancher,' 7 which stops and prevents the flow

* γένη conj. Dalec.; μέρη Ald.

29 I

³ Somali arrow-poison. Index App. (27).

^{*} γένη conj. Dalec.; μερη ωι...
* περιττότατα conj. W.; περιττοτάτη Ald.
* Plin. 25. 83.

οί δὲ καὶ σφοδροτέρως διατμηθείσης ἴσχειν καὶ κωλύειν την χύσιν. [ταῦτα μεν οὖν, ὥσπερ εἴποτῶν μὲν οὖν ἔξω μεν. ξοικε δηλοῦν τὸ κοινόν.]

τόπων οἱ φαρμακωδέστατοι οὖτοι. Τῶν δὲ περὶ τὴν Ἑλλάδα τόπων φαρμακωδέστατον τό τε Πήλιον τὸ ἐν Θετταλία καὶ τὸ Τελέθριον τὸ ἐν Εὐβοία καὶ ὁ Παρνασός, ἔτι δὲ καὶ ἡ ᾿Αρκαδία καὶ ἡ ἹΛακωνική καὶ γὰρ αὖται φαρμακώδεις άμφότεραι δι' δ και οί γε 'Αρκάδες είωθασιν άντι του φαρμακοποτείν γαλακτοποτείν περί τὸ ἔαρ, ὅταν οἱ ὁποὶ μάλιστα τῶν τοιούτων φύλλων ακμάζωσι τότε γαρ φαρμακωδέστατον τὸ γάλα πίνουσι δὲ βόειον δοκεί γὰρ πολυνομώτατον καὶ παμφαγώτατον είναι πάντων ό Βοῦς.

Φύεται δὲ παρ' αὐτοῖς ὅ τε ἐλλέβορος ἀμφότερος καὶ ὁ λευκὸς καὶ ὁ μέλας ἔτι δὲ δαῦκον δαφνοειδές κροκόεν, καὶ ην ἐκεῖνοι μὲν ράφανον άγρίαν καλουσι των δ' ιατρών τινες κεράϊν, καλ ην οι μεν άλθαίαν εκείνοι δε μαλάχην άγρίαν, καὶ ή αριστολοχία καὶ τὸ σέσελι καὶ τὸ ἱπποσέλινον καὶ τὸ πευκέδανον καὶ ἡ Ἡράκλεια καὶ ὁ στρύχνος άμφότερος ὅ τε φοινικοῦν ἔχων τὸν καρπὸν και ὁ μέλανα.

Φύεται δὲ καὶ ὁ σίκυος ὁ ἄγριος, ἐξ οὖ τὸ έλατήριον συντίθεται καὶ ὁ τιθύμαλλος, έξ οὖ τὸ ίπποφάες· ἄριστον δὲ τοῦτο περί Τεγέαν κἀκεῖνο μάλιστα σπουδάζεται· φύεται δ' ἐκεῖ ἐπὶ πλέον·

292

¹ I omit ταῦτα . . . κοινόν as apparently out of place and a duplicate of the last sentence of § 8. ³ Plin. 25, 110. Plin. 25. 94; cf. 4. 5. 2.

ENQUIRY INTO PLANTS, IX. xv. 3-6

of blood, some say if the vein is merely pricked, others even if it is deeply cut into. These then of the places outside Hellas are those that are most

productive of drugs.

² Of places in Hellas those most productive of drugs are Pelion in Thessaly, Telethrion in Euboea, Parnassus, and also Arcadia and Laconia, for both these states produce medicinal herbs; wherefore the Arcadians are accustomed, instead of drinking medicine, to drink milk in spring when the juices of such plants are at their best, for then the milk has most medicinal virtue. It is cows' milk that they drink, since it appears that the cow eats more than any other animal and is more impartial as to what she eats.

⁸ Both kinds of hellebore, the white and the black, grow in their country, and also carrot, ⁴ a saffroncoloured plant like bay, and a plant which the Arcadians call 'wild cabbage' ⁵ (spurge) but some physicians keraïs; also a plant called by some marsh mallow, ⁶ also birthwort hartwort alexanders sulphurwort Herakleia, and both kinds of strykhnos, ⁷ that which has a scarlet and that which has a black fruit.

There also grow there the 'wild cucumber' (squirting cucumber), of which the drug 'driver' is compounded, and the tithymallos (spurge) of which hippophaës is made; this is best about Tegea, and that kind is much sought after; it grows there in

⁵ cf. 9. 12. 1. and Index.

⁴ δαῦκον. This name recurs § 8 and 9. 20. 2. Text must be defective here: the epithets are unintelligible, and perhaps belong to another plant whose name has dropped out. See Index.

άλθαίαν conj. Sch., cf. 9. 18. 1; ἀλθέαν Ald. cf. Plin. 20. 222.
 cf. 9. 11. 5.
 cf. 9. 9. 4; 9. 14. 1.

ξ, 9, 11, 5.
 lπποφdes is elsewhere the name of a plant: cf. Diosc. 4, 159,
 δ ω may be corrupt, or the text defective.

πλείστον δε καὶ κάλλιστον φύεται περὶ τὴν Κλειτορίαν.

7 'Η δέ πανάκεια γίνεται κατά τὸ πετραĵον περὶ Ψωφίδα καὶ πλείστη καὶ ἀρίστη. τὸ δὲ μῶλυ περί Φενεον και έν τη Κυλλήνη. φασι δ΄ είναι καὶ ὅμοιον οἱ ὁ "Ομηρος εἴρηκε, τὴν μεν ρίζαν έγον στρογγύλην προσεμφερή κρομύω τὸ δὲ φύλλον δμοιον σκίλλη γρησθαι δε αὐτῷ πρός τε τὰ άλεξιφάρμακα και τάς μαγείας ου μην ορύττειν γ' είναι χαλεπόν, ώς "Ομηρός φησι.

Τὸ δὲ κώνειον ἄριστον περί Σοῦσα καὶ ἐν τοῖς ψυχροτάτοις τόποις. γίνεται δὲ καὶ ἐν τῆ Λακωνική τὰ πολλὰ τούτων καὶ γὰρ αὕτη πολυφάρμακος. ἐν ᾿Αχαία δὲ ἥ τε τραγάκανθα πολλή καὶ οὐδὲν χείρων ὡς οἴονται τῆς Κρητικῆς ἀλλά καὶ τῆ ὄψει καλλίων καὶ δαῦκον περὶ τὴν Πατραϊκήν διαφέρον τοῦτο δὲ θερμαντικὸν φύσει, ρίζαν δὲ ἔχει μέλαιναν. Φύεται δὲ τὰ πολλά τούτων καλ έν τῷ Παρνασῷ καλ περὶ τὸ Τελέκαὶ ταῦτα μέν κοινὰ πλειόνων χώρων.

ΧVΙ. Τὸ δὲ δίκταμνον ἴδιον τῆς Κρήτης, θαυμαστον δε τη δυνάμει και προς πλείω χρήσιμον μάλιστα δὲ πρὸς τοὺς τόκους τῶν γυναικὧν. ἔστι δὲ τὸ μὲν φύλλον παρόμοιον τῆ βληχοῖ, ἔχει δέ τι καί κατά τὸν χυλὸν ἐμφερές τὰ δὲ κλωνία λεπτότερα. χρῶνται δὲ τοῖς φύλλοις, οὐ τοῖς κλωσὶν οὐδὲ τῷ καρπῷ· χρήσιμον δὲ πρὸς πολλὰ μεν καὶ ἄλλα, μάλιστα δέ, ὅσπερ ελέχθη, πρὸς

294

Plin. 25, 30-33. ² κατὰ coni. St.: καὶ Ald. H.

³ Plin. 25, 151.

⁴ Σοῦσα: cf. 9. 16. 8; Λοῦσα (a town in Arcadia) conj. Sch. (usually Λοῦσοι), the other places mentioned being all in

ENQUIRY INTO PLANTS, IX. xv. 6-xvi, i

considerable abundance, but in greatest abundance and best about Kleitoria.

¹ All-heal grows in great abundance and best in ² the rocky ground about Psophis, moly about Pheneos and on Mount Kyllene. They say that this plant is like the moly mentioned by Homer, that it has a round root like an onion and a leaf like squill, and that it is used against spells and magic arts, but that

it is not, as Homer says, difficult to dig up.

3 Hemlock is best about Susa 4 and in the coldest spots. Most of these plants occur also in Laconia, for this too is a land rich in medicinal herbs. In Achaia tragacanth 5 is abundant and is as good as that of Crete, it is believed, and even fairer in appearance. Daukon 6 again is excellent in the country about Patrai 7; this is by nature healing, and it has a black root. Most of these grow also on Mount Parnassus So these plants are comand about Telethrion. mon to several lands.

Of the medicinal herbs peculiar to Crete.

XVI. 8But dittany is peculiar to Crete. This plant is marvellous in virtue and is useful for many purposes, but especially for women in child-birth. Its leaf is like pennyroyal, to which it also bears some resemblance in taste; but the twigs are slenderer. They use the leaves, not the twigs nor the fruit: and the leaf is useful for many other purposes, but above all,

But Plin. 25. 154 has Susa: it can hardly be the Hellas. Persian town.

⁵ Plin. 13, 115.

 Repeated 9. 20. 2; cf. 9. 15. 5 and Index.
 Πατραϊκήν conj. Sch., cf. 9. 20. 2; πατρικήν Ald.; σπαρτιαкhv U*; отартину MP; Patrensi agro G.

8 Plin. 25, 92,

τὰς δυστοκίας τῶν γυναικῶν ἡ γὰρ εὐτοκεῖν φασι ποιεῖν ἡ παύειν γε τοὺς πόνους ὁμολογουμένως δίδοται δὲ πίνειν ἐν ὕδατι. σπάνιον δέ ἐστι καὶ γὰρ ὀλίγος ὁ τόπος ὁ φέρων, καὶ τοῦτον αἱ αἰγες ἐκνέμονται διὰ τὸ φιληδεῖν. ἀληθὲς δέ φασιν εἰναι καὶ τὸ περὶ τῶν βελῶν, ὅτι φαγούσαις ὅταν τοξευθῶσι ἐκβάλλει. τὸ μὲν οὖν δίκταμνον τοιοῦτόν τε καὶ τοιαύτας ἔχει τὰς δυνάμεις.

Τὸ δὲ ψευδοδίκταμνον τῷ μὲν φύλλῳ ὅμοιον τοῖς κλωνίοις δ' ἔλαττον τῷ δυνάμει δὲ πολὺ λειπόμενον. βοηθεῖ μὲν γὰρ καὶ ταὐτά, χεῖρον δὲ πολλῷ καὶ ἀσθενέστερον. ἔστι δὲ εὐθὺς ἐν τῷ στόματι φανερὰ τοῦ δικτάμνου ἡ δύναμις· διαθερμαίνει γὰρ ἀπὸ μικροῦ σφόδρα. τιθέασι δὲ τὰς δεσμίδας ἐν νάρθηκι ἡ καλάμῳ πρὸς τὸ μὴ ἀποπνεῖν· ἀσθενέστερον γὰρ ἀποπνεῦσαν. λέγουσι δέ τινες ὡς ἡ μὲν φύσις μία ἡ τοῦ δικτάμνου καὶ ἡ τοῦ ψευδοδικτάμνου, διὰ δὲ τὸ ἐν εὐγειοτέροις φύεσθαι τόποις χεῖρον γίνεσθαι, καθάπερ καὶ ἄλλα πολλὰ πλείω τούτων κατὰ τὰς δυνάμεις. τὸ γὰρ δίκταμνον φιλεῖ χώραν τραχεῖαν.

"Έστι δὲ καὶ ἔτερον δίκταμνον ὥσπερ ὁμώνυμον, οὔτε τὴν ὄψιν οὔτε τὴν δύναμιν ἔχον τὴν αὐτήν φύλλον γὰρ ἔχει ὅμοιον σισυμβρίφ τοὺς δὲ κλῶνας μείζους ἔτι δὲ τὴν χρείαν καὶ τὴν δύναμιν οὖκ ἐν τοῖς αὐτοῖς. τοῦτο μὲν οὖν, ὥσπερ ἐλέχθη, θαυμαστὸν ἄμα καὶ ἴδιον τῆς νήσου. φασὶ δέ

¹ ἐκβάλλει conj. Sch.; ἐκβάλλειν Ald.

²¿Plin. 25. 93.

 $^{^3}$ νάρθηκι ή conj. Sch.; ναρθηκίδη ή U; ναρθηκίδι ή M; νάρθηκι καὶ Ald.

ENQUIRY INTO PLANTS, IX. xvi. 1-3

as was said, against difficult labour in women; for it is said that either it makes labour quite easy or at least it confessedly makes the pains to cease: it is given as a draught in water. It is a scarce plant: for the region which bears it is not extensive, and the goats graze it down because they are fond of it. The story of the arrows is also said to be true,—that, if goats eat it when they have been shot, it rids them 1 of the arrow. Such then is dittany and

such its properties.

2'False dittany' is like it in leaf, but has smaller twigs, and in virtue is far inferior. For it is of service in the same ways, but is feebler and not nearly so powerful. The virtue of dittany is perceived directly it is taken into the mouth: for a small piece of it has a very warming effect. The bunches of it are put in the hollow stem of ferula 3 or a reed, so that it may not exhale its virtue: for, if it does so, it is less effective. Some say that dittany and 'false dittany' are essentially the same plant, but that the latter is an inferior form produced by growing in places with richer soil; just as many other things become inferior in their properties for the same cause. For dittany loves rough ground.

⁵ There is also another plant called 'dittany,' though it has nothing in common with these except the name. This has neither the same appearance nor the same virtue; for its leaf is like bergamot-mint and its twigs are larger, and further its use and virtue are differently shewn. The true plant is, as was said, marvellous, and is also peculiar to the island of Crete. Indeed some say that the plants

⁴ πλείω τούτων Ald., probably a duplicate of ἄλλα πολλὰ; not represented in G; ἀλλοιοῦται conj. W, Plin.25. 94.

τινες όλως των φύλλων καὶ των ὀροδάμνων καὶ άπλως των ὑπὲρ γης τὰ ἐν Κρήτη διαφέρειν, των δὲ ἄλλων των γε πλείστων τὰ ἐν τῷ Παρνασῷ.

Τὸ δ' ἀκόνιτον γίνεται μὲν καὶ ἐν Κρήτη καὶ ἐν Ζακύνθω, πλείστον δε καὶ ἄριστον εν Ἡρακλεία τη εν Πόντω. έχει δε φύλλον μεν κιχοριώδες, ρίζαν δε δμοίαν τῷ σχήματι καὶ τῷ χρώματι καρίδι, την δε δύναμιν την θανατηφόρον εν ταύτη τὸ δὲ Φύλλον καὶ τὸν καρπὸν οὐθέν φασι ποιείν. καρπὸς δέ ἐστι πόας οὐχ ὑλήματος. βραχεῖα δὲ ή πόα καὶ οὐδὲν ἔχουσα περιττόν, ἀλλὰ παρομοία τῶ σίτω τὸ δὲ σπέρμα οὐ σταχυηρόν. Φύεται δὲ πανταγού και ούκ έν ταις 'Ακόναις μόνον, ἀφ' ὧν έχει την προσηγορίαν αυτη δέ έστι κώμη τις των Μαριανδυνών φιλεί δε μάλιστα τους πετρώδεις τόπους οὐ νέμεται δὲ οὔτε πρόβατον οὔτ' ἄλλο 5 ζωον οὐδέν. συντίθεσθαι δὲ τρόπον τινὰ πρὸς τὸ έργάζεσθαι καὶ οὐ παντὸς είναι δι' δ καὶ τοὺς ιατρούς οὐκ ἐπισταμένους συντιθέναι σηπτικώ τε γρησθαι καὶ πρὸς ἄλλα ἄττα· πινόμενον δ' ούδεμίαν αϊσθησιν ποιείν ουτ' έν οινω ουτ' έν μελικράτω συντίθεσθαι δε ώστε κατά γρόνους τακτούς άναιρείν, οίον δίμηνον τρίμηνον έξάμηνον ένιαυτόν, τους δε και δύο έτη γείριστα δε άπαλ-

² Diosc. 4. 76 and 77; Plin. 27. 9 and 10.

¹ δροδάμνων: this word seems to occur only here in T.

³ καρίδι conj. W.; καρίαι U; καρύα Ald. cf. Diosc. l.c. + cf. 9. 8. 1.

⁵ Plin. 6. 4, portus Acone veneno aconito dirus. But in 27. 10. he apparently did not recognise 'Ακόναις as a proper name, 208

ENQUIRY INTO PLANTS, IX. xvi. 3-5

of Crete are superior in leaves boughs 1 and in general all the parts above ground to those of other places; while those of Parnassus are superior to most of those found elsewhere.

Of wolf's-bane and its habitat, and of meadow-saffron.

² Wolf's-bane grows in Crete and in Zakynthos, but is most abundant and best at Herakleia in Pontus. It has a leaf like chicory, a root like in shape and colour to a prawn,3 and in this root resides its deadly property, whereas they say that the leaf and the fruit produce no effects. The fruit is that of a herb.4 not that of a shrub or tree. It is a lowgrowing herb and shows no special feature, but is like corn, except that the seed is not in an ear. It grows everywhere and not only at Akonai, from whence it gets its name (this is a village of the Mariandynoi) 6: and it specially likes rocky ground. Neither sheep nor any other animals eat it.7 In order to be effective it is said that it must be compounded in a certain manner, and that not everyone can do this: and so that physicians, not knowing how to compound it, use it as a septic and for other purposes: and 8 that, if drunk mixed in wine or a honey-posset, it produces no sensation: but that it can be so compounded as to prove fatal at a certain moment which may be in two three or six months, or in a year, or even in two years: and that the

and translates it in nudis cautibus, misled perhaps by τοὺς πετρώδεις τόπους below.

6 Μαριανδυνών conj. Meurs.; περιανδύνων U*Ald.H.

⁷ U* adds here βοηθεῖαι δὲ τοῖς ἐνεγκαμένοις εἰσί and omits §§ 5, 6... εἰδέναι, continuing πολλάκις γὰρ φασὶ τὰ ἀνδράποδα. δὲ add. Sch.

λάττειν τοὺς ἐν πλείστφ χρόνφ καταφθίνοντος τοῦ σώματος, ἡᾶστα δὲ τοὺς παραχρῆμα. λυτικὸν δὲ φάρμακον οὐχ εὑρῆσθαι, καθάπερ ἀκούομεν ἐτέρων τι φύεσθαι. ἀλλὰ τοὺς ἐγχωρίους ἀνασώζειν τινὰς μέλιτι καὶ οἴνφ καὶ τοιούτοις τισί, σπανίως δὲ καὶ τούτους καὶ ἐργωδῶς.

'Αλλὰ τοῦ ἐφημέρου τὸ φάρμακον εὐρῆσθαι· ἔτερον χάρ τι ρίζιον εἶναι ὁ ἐφήμερον ἀπαλλάττει· τοῦτο δὲ φύλλον ὅμοιον ἔχειν τῷ ἐλλεβόρῳ ἡ τῷ λειρίῳ· καὶ τοῦτο πάντας εἰδέναι· δι' ὁ καὶ τὰ ἀνδράποδά φασι πολλάκις παροργισθέντα χρῆσθαι, κἄπειτα ἰατρεύειν αὐτὰ πρὸς τοῦτο ὁρμῶντα, καὶ γὰρ οὐδὲ ταχεῖαν ποιεῖσθαι τὴν ἀπαλλαγὴν οὐδὲ ἐλαφρὰν ἀλλὰ δυσχερῆ καὶ χρόνιον· εἰ μὴ ἄρα διὰ τὸ εὐθεράπευτον εἶναι καὶ ἀκατασκεύαστον ὡς δεῖ. φασὶ γοῦν καὶ παραχρῆμα ἀπαλλάττεσθαι καὶ ὕστερον χρόνῳ τοὺς δὲ καὶ εἰς ἐνιαυτὸν ἄγειν, καὶ τὰς δόσεις ἀβοηθήτους εἶναι.

² ἐτέρων conj. Sch.; ἔτερόν τι φύεσθαι UAld. Η.; ἔτερόν τι φυόμενον conj. W. G seems to have had a fuller text.

3 άλλα τούς έγχ. UM; άλλα τινες τοις ένεγκαμένοις βοήθειαι εὔρηνται· τους γαρ έγχ. Ald.H., which the indicative εὔρηνται

shews to be a gloss.

¹ i.e. no herb having that effect.

⁴ τοῦ ἐφημέρου U; τὰ ἐφ' ἡμέρου M; καὶ τὸ ἐφήμερου Ald. The passage about ἐφήμερου, which interrupts the account of ἀκόνιτου, is confused, and the text probably defective; translation a makeshift. The sense of εἰ μὴ... ὡς δεῖ being perhaps irrecoverable, the connexion of what follows is obscured. W. gives up the passage.

ENQUIRY INTO PLANTS, IX. xvi. 5-6

longer the time the more painful the death, since the body then wastes away, while, if it acts at once, death is quite painless. And it is said that no antidote 1 which can counteract it has been discovered, like the natural antidotes to other 2 poisonous herbs of which we are told: though the country-folk 3 can sometimes save a man with honey and wine and such like things, only however occasionally and with difficulty.

(On the other hand they say that for meadow-saffron 4 the antidote has been found: for that there is another root which counteracts that herb: 5 and that it 6 has a leaf like hellebore 7 or the madonna lily: 8 and that this 9 is generally known. Wherefore they say that slaves often take meadow-saffron when greatly provoked, and then themselves have recourse 10 to the antidote and effect a cure,—seeing that the poison does not cause a speedy and easy death, but 11 one that is lingering and slow,—unless indeed, merely because the cure is so easy, 12 the antidote has not been properly prepared. 13 At least they say that though death may ensue at once, sometimes it only occurs after a considerable interval, which in some cases extends to a year, and that in these latter cases the dose given has incurable effects: and that

 $^{^{\}mathbf{6}}$ δ έφήμερον PH.; δ έφημεραΐον U; δ έφ' ήμεραΐον M; δ οὐκ έφήμερον Ald.

⁶ τοῦτο δὲ Ald.; τόνδε δὲ καὶ U; τῶνδε δὲ καὶ M.

i.e. the 'black': see Index.

δ λειρίφ conj. Guilandinus from Diosc. 4. 84 (κρίνφ); αἰρίφ Ald. Η.
• τοῦτο Ald.; τοῦτο μὲν UM.

After δρμῶντα UM add καὶ τοὺς οἰκέτας ἐπὶ τοῦτο δρμῶν and omit καὶ γὰρ... θανατηφόρων.
¹¹ ἀλλὰ Ald.; οὐδὲ U*.

¹² ευθεράπευτον Ald.; αθεράπευτον U*P.

¹³ In which case apparently the slave outwits himself as well as his master by 'dying on him.'

ταῦτα δὲ ἐξακριβωθῆναι μάλιστα παρὰ τοῖς Τυρ7 ρηνοῖς τοῖς ἐν Ἡρακλείᾳ. τοῦτο μὲν <οὖν> οὐδὲν ἄτοπον, εἰ τρόπον μέν τινα ἀβοήθητον ἄλλως δὲ βοηθήσιμον, ὥσπερ καὶ ἔτερα τῶν θανατηφόρων.

Τὸ δὲ ἀκόνιτον ἄχρηστον, ὥσπερ εἴρηται, τοῖς μὴ ἐπισταμένοις· οὐδὲ κεκτῆσθαι δὲ ἐξεῖναι, ἀλλὰ θάνατον τὴν ζημίαν· τὴν δὲ τῶν χρόνων διαφορὰν ἀκολουθεῖν κατὰ τὰς συλλογάς· ἰσοχρόνους γὰρ τοὺς θανάτους γίνεσθαι τοῖς ἀπὸ τῆς συλλογῆς χρόνοις.

Θρασύας δ' ο Μαντινεὺς εὐρήκει τι τοιοῦτον, ὅσπερ ἔλεγεν, ὅστε ῥαδίαν ποιεῖν καὶ ἄπονον τὴν ἀπόλυσιν τοῖς ὀποῖς χρώμενος κωνείου τε καὶ μήκωνος καὶ ἐτέρων τοιούτων, ὅστε εὔογκον εἶναι σφόδρα καὶ μικρὸν ὅσον εἰς δραχμῆς ὁλκήν. ἀβοήθητον δὲ πάντη καὶ δυνάμενον διαμένειν ὁποσονοῦν χρόνον καὶ οὐδὲν ἀλλοιούμενον. ἐλάμβανε δὲ τὸ κώνειον οὐχ ὅθεν ἐτύγχανεν ἀλλ' ἐκ Σούσων καὶ εἴ τις ἄλλος τόπος ψυχρὸς καὶ παλίσκιος· ὡσαύτως δὲ καὶ τἄλλα. συνετίθει δὲ καὶ ἔτερα φάρμακα πολλὰ καὶ ἐκ πολλῶν. δεινὸς δὲ καὶ ᾿Αλεξίας ὁ μαθητὴς αὐτοῦ καὶ οὐχ ἤττον ἔντεχνος ἐκείνου· καὶ γὰρ τῆς ἄλλης ἰατρικῆς ἔμπειρος.

¹ οδν add. W.

² Δκολουθείν κατά conj. W.; ἀκούειν είναι κατά Ald. Η.; ἀκουείν είναι καὶ Μ.

ENQUIRY INTO PLANTS, IX. xvi. 6-8

these facts have been most carefully ascertained among the Tyrrhenians of Herakleia. Now 1 it is not surprising that in some circumstances the effects of the poison should be incurable, and yet in others that a cure should be possible, this being also the case with other deadly poisons.)

To return—wolf's bane, as has been said, is useless to those who do not understand it; in fact it is said that it is not lawful even to have it in one's possession, under pain of death; also that the length of time which it takes to produce its effects depends on 2 the time when it is gathered; for that the time which it takes to kill is equal to that which has elapsed since it was gathered.

Of two famous druggists and of the virtues of hemlock.

Thrasyas of Mantineia had discovered, as he said, a poison which produces an easy and painless end; he used the juices of hemlock poppy and other such herbs, so compounded as to make a dose of conveniently small size, weighing only somewhat less than a quarter of an ounce. For the effects of this compound there is absolutely no cure, and it will keep any length of time without losing its virtue at all. He used to gather his hemlock, not just anywhere, but at Susa³ or some other cold and shady spot; and so too with the other ingredients; he also used to compound many other poisons, using many ingredients. His pupil Alexias was also clever and no less skilful than his master, being also versed in the science of medicine generally.

Digitized by Google

³ Σούσων MSS.; Λούσων conj. Sch. cf. 9. 15. 8 n. The mention of Mantineia makes it likely that a place in Arcadia is intended.

Ταῦτα μὲν οὖν εὑρῆσθαι δοκεῖ πολλῷ μᾶλλον νῦν ἡ πρότερον. ὅτι δὲ διαφέρει τὸ χρῆσθαί πως ἐκάστφ φανερὸν ἐκ πολλῶν ἐπεὶ καὶ Κεῖοι τῷ κωνείφ πρότερον οὐχ οὕτω ἀλλὰ τρίβοντες ἐχρῶντο, καθάπερ οἱ ἄλλοι νῦν δ' οὐδ' ἄν εἶς τρίψειεν, ἀλλὰ περιπτίσαντες καὶ ἀφελόντες τὸ κέλυφος, τοῦτο γὰρ τὸ τὴν δυσχέρειαν παρέχον δυσκατέργαστον ὄν, μετὰ ταῦτα κόπτουσιν ἐν τῷ ὅλμῷ καὶ διαττήσαντες λεπτὰ ἐπιπάττοντες ἐφ' ὕδωρ πίνουσιν, ὥστε ταχεῖαν καὶ ἐλαφρὰν γίνεσθαι τὴν ἀπαλλαγήν.

XVII. 'Απάντων δὲ τῶν φαρμάκων αἱ δυνάμεις ἀσθενέστεραι τοῖς συνειθισμένοις τοῖς δὲ καὶ ἀνενεργεῖς τὸ ὅλον. ἔνιοι γὰρ ἐλλέβορον ἐσθίοντες πολὺν ὥστε ἀναλίσκειν δέσμας ὅλας οὐδὲν πάσχουσιν· ὅπερ ἐποίει καὶ Θρασύας δεινότατος ῶν ὡς ἐδόκει περὶ τὰς ῥίζας. ποιοῦσι δὲ τοῦθ' ὡς ἔοικε καὶ τῶν νομέων τινές· δι' δ καὶ πρὸς τὸν φαρμακοπώλην τὸν θαυμαζόμενον ὡς κατήσθιε ῥίζαν μίαν ἡ δύο παραγενόμενος ὁ ποιμὴν καὶ ἀναλώσας ὅλην τὴν δέσμην ἐποίησεν ἀδόκιμον· ἐλέγετο δ' ὅτι καθ' ἐκάστην ἡμέραν τοῦτο ποιεῖ καὶ αὐτὸς καὶ ἔτεροι.

 Κινδυνεύει γὰρ ἔνια τῶν φαρμάκων τῆ ἀσυνηθεία φάρμακα γίνεσθαι, τάχα δὲ ἀληθέστερον

¹ δυσκατέργαστον: cf. C.P. 1 14. 4.
2 διαττήσαντες conj. Hoffmann from G; διαπτήσαντες Ald. H.; διαθήσαντες U*mP.

ENQUIRY INTO PLANTS, IX. xvi. 9-xvit. 2

Now these things seem to have been ascertained far better in recent than in former times. And many things go to shew that the method of using the various drugs makes a difference; thus the people of Ceos formerly did not use hemlock in the way described, but just shredded it up for use, as did other people; but now not one of them would think of shredding it, but they first strip off the outside and take off the husk, since this is what causes the difficulty, as it is not easily assimilated ¹; then they bruise it in the mortar, and, after putting it through a fine sieve, ² sprinkle it on water and so drink it; and then death is made swift and easy.

How use diminishes the efficacy of drugs, and how they have not the same effect on all constitutions.

XVII. ³The virtues of all drugs become weaker to those who are accustomed to them, and in some cases become entirely ineffective. Thus some eat enough hellebore to consume whole bundles and yet suffer no hurt; this is what Thrasyas did, who, as it appeared, was very cunning in the use of herbs. And it appears that shepherds sometimes do the like; wherefore the shepherd who came before the vendor of drugs (at whom men marvelled because he ate one or two roots) and himself consumed the whole bundle, destroyed the vendor's reputation: it was said that both this man and others did this every day.

For it seems that some poisons become poisonous because they are unfamiliar, or perhaps it is a more accurate way of putting it to say that familiarity makes

⁸ Plin. 27, 144.

305

VOL. II.

X

είπειν ώς τη συνηθεία ου φάρμακα προσδεξαμένης γάρ της φύσεως καὶ κατακρατούσης οὐκέτι φάρμακα, καθάπερ καὶ Θρασύας έλεγεν έκεινος γὰρ ἔφη τὸ αὐτὸ τοῖς μὲν φάρμακον είναι τοῖς δ' ου φάρμακον, διαιρών τὰς φύσεις έκάστων ώετο γαρ δείν καὶ ην δεινὸς διαγνώναι. ποιεί δέ τι δηλον ὅτι πρὸς τῆ φύσει καὶ τὸ ἔθος. Εὔδημος γοῦν ο φαρμακοπώλης εὐδοκιμῶν σφόδρα κατὰ την τέχνην συνθέμενος μηδέν πείσεσθαι πρὸ ήλίου δύναντος κατέφαγε μέτριον πάνυ καὶ οὐ 8 κατέσγεν οὐδ' ἐκράτησεν. ὁ δὲ Χίος Εὔδημος πίνων ελλέβορον οὐκ εκαθαίρετο, καί ποτε εφη πιείν εν μια ήμερα δύο και είκοσι πόσεις εν τή άγορα καθήμενος έπι των σκευών και οὐκ έξαναστηναι πρό τοῦ δείλην γενέσθαι τότε δ' έλθων καὶ λούσασθαι καὶ δειπνεῖν ὥσπερ εἰώθει καὶ οὐκ έξεμέσαι πλην οὐτός γε βοήθειάν τινα παρασκευασάμενος κατέσχε κίσσηριν γαρ έπιπάττων έπ' όξος δριμύ πιείν έφη μετά την έβδόμην πόσιν, καὶ πάλιν ὕστερον ἐν οἴνφ τὸν αὐτὸν τρόπον τὴν δὲ τῆς κισσήριδος οὕτως ἰσχυρὰν εἶναι δύναμιν ωστ' εάν τις είς πίθον ζέοντα <οίνου> εμβάλη παύειν την ζέσιν ου παραχρημα μόνον άλλα καί όλως καταξηραίνουσάν τε δηλον ότι και άναδεχομένην τὸ πνεῦμα καὶ τοῦτο διϊείσαν, οῦτος μὲν οὖν τό γε πληθος ταύτη τη βοηθεία κατέσχεν. 4 "Οτι δὲ καὶ τὸ ἔθος ἰσχυρὸν φανερὸν ἐκ πολλῶν"

¹ This story is quoted by Apollonius, *Hist. Mirab.* 50. 306

ENQUIRY INTO PLANTS, IX. xvii. 2-4

poisons non-poisonous; for, when the constitution has accepted them and prevails over them, they cease to be poisons, as Thrasyas also remarked; for he said that the same thing was a poison to one and not to another; thus he distinguished between different constitutions, as he thought was right; and he was clever at observing the differences. Also, besides the constitution, it is plain that use has something to do with it. At least Eudemus, the vendor of drugs, who had a high reputation in his business, after making a wager that he would experience no effect before sunset, drank a quite moderate dose, and it proved too strong for his power of resistance: 1 while the Chian Eudemus took a draught of hellebore and was not purged. And on one occasion he said that in a single day he took two and twenty draughts in the market-place as he sat at his stall, and did not leave the place till it was evening, and then he went home and had a bath and dined, and was not sick. However this man was able to hold out because he had provided himself with an antidote; for he said that after the seventh dose he took a draught of tart vinegar with pumice-stone dust in it, and later on took a draught of the same in wine in like manner: and that the virtue of the pumice-stone dust is so great that, if one puts it into a boiling pot of wine,2 it causes it to cease to boil, not merely for the moment, but altogether, clearly because it has a drying effect and it catches the vapour and passes it off. It was then by this antidote that Eudemus was able to contain himself in spite of the large quantity of hellebore which he took.

However many things go to show that use makes

307

² oĭvou add. Sch., cf. Plin. 36. 42; 14, 138.

έπεὶ καὶ τὸ ἀψίνθιον τὰ μὲν ἐνταῦθα πρόβατα οὕ φασί τινες νέμεσθαι, τὰ δ' ἐν τῷ Πόντῷ νέμεται καὶ γίνεται πιότερα καὶ καλλίω καί, ὡς δή τινες λέγουσιν, οὖκ ἔχοντα χολήν. ἀλλὰ γὰρ ταῦτα μὲν ἑτέρας ἄν τινος εἴη θεωρίας.

ΧVIII. Αἱ δὲ ῥίζαι καὶ τὰ ὑλήματα, καθάπερ εἴρηται, πολλὰς ἔχουσι δυνάμεις οὐ πρὸς τὰ ἔμ-ψυχα σώματα μόνον ἀλλὰ καὶ πρὸς τὰ ἄψυχα. λέγουσι γὰρ ἄκανθάν τινα εἶναι ἢ πήγνυσι τὸ ὕδωρ ἐμβαλλομένη· πηγνύναι δὲ καὶ τὴν τῆς ἀλθαίας ῥίζαν, ἐάν τις τρίψας ἐμβάλη καὶ θῆ ὑπαίθριον· ἔχει δὲ ἡ ἀλθαία φύλλον μὲν ὅμοιον τῆ μαλάχη πλὴν μεῖζον καὶ δασύτερον, τοὺς δὲ καυλοὺς μαλακούς, ἄνθος δὲ μήλινον, καρπὸν δ΄ ὅμοιον τῆ μαλάχη, ῥίζαν δὲ ἰνώδη λευκὴν τῆ γεύσει δὲ ὥσπερ τῆς μαλάχης ὁ καυλός· χρῶνται δὲ αὐτῆ πρός τε τὰ ῥήγματα καὶ τὰς βῆχας ἐν οἴνω γλυκεῖ καὶ ἐπὶ τὰ ἔλκη ἐν ἐλαίω.

Έτέραν δέ τινα συνεψομένην τοις κρέασι συνάπτειν εἰς ταὐτὸ καὶ οίον πηγνύναι τὰς δὲ καὶ ἔλκειν, ὥσπερ ἡ λίθος καὶ τὸ ἤλεκτρον. καὶ ταῦτα μὲν ἐν τοις ἀψύγοις.

Τὸ δὲ θηλύφονον, οἱ δὲ σκορπίον καλοῦσι διὰ τὸ τὴν ῥίζαν ὁμοίαν ἔχειν τῷ σκορπίῳ, ἐπιξυό-

¹ cf. Plin. 27. 45.

² δλήματα: here a general term for shrubs and under-shrubs. cf. 9. 20. 6.

³ Diosc. 3. 146; Plin. 20. 84.

ENQUIRY INTO PLANTS, IX. xvii. 4-xviii. 2

much difference; 1 thus some say that the sheep of some places do not eat wormwood; yet those of Pontus not only eat it but become fatter and fairer and, as some say, have no bile. But these things may be said to belong to a different enquiry.

Of plants that possess properties affecting lifeless objects.

XVIII. Herbs and shrubs,² as has been said, have many virtues which are shown in their effects not only on living bodies but on lifeless ones. Thus they say that there is a kind of akantha (gum arabic) which thickens water, when it is put in it; ³ and that so also does the root of marsh-mallow if one shreds it and puts it in and stands the water in the open air. Marsh-mallow has a leaf like mallow, but larger and rougher; the stems are soft, the flower yellow, the fruit like that of mallow, the root fibrous and white, with a taste like that of the stem of mallow. They use it for fractures and for coughs in sweet wine, and for sores in olive-oil.

⁴ They say that there is another kind which, if cooked with meat, combines with it and as it were sets it hard; and there are others that attract things to them, like the magnet or amber. So much for effects produced on lifeless things.

Of plants whose properties affect animals other than man.

^b Wolf's bane, which some call 'scorpion-plant because it has a root like a scorpion, kills that animal

4 Referred to by Apollon. Hist. Mirab. 41. cf. Diosc. 3. 147; Plin. 27, 42: 25, 67.

⁶ Referred to by Ael. H.A. 9. 27; Apollon. Hist. Mirab. 41. cf. Plin. 25. 122 (cf. 27. 6); Diosc. 4. 76. This is evidently a different plant to the σκορπίος mentioned 9. 13. 6. See Index.

309

μενον ἀποκτείνει τὸν σκορπίον ἐὰν δέ τις έλλέβορον λευκὸν καταπάση, πάλιν ἀνίστασθαί φασιν ἀπόλλυσι δὲ καὶ βοῦς καὶ πρόβατα καὶ ύποζύγια καὶ άπλῶς πᾶν τετράπουν ἐὰν εἰς τὰ αίδοια τεθή ή ρίζα ή τὰ φύλλα αὐθήμερον χρήσιμον δὲ πρὸς σκορπίου πληγὴν πινόμενον. έχει δὲ τὸ μὲν φύλλον ὅμοιον κυκλαμίνω τὴν δὲ ρίζαν, ωσπερ ελέχθη, σκορπίφ. φύεται δε ωσπερ ή ἄγρωστις καὶ γόνατα ἔχει φιλεῖ δὲ χωρία σκιώδη. εί δὲ ἀληθη τὰ περί τὸν σκορπίον ήδη καὶ τάλλα, οὐκ ἀπίθανα τὰ τοιαῦτα, καὶ τὰ 3 μυθώδη δὲ οὐκ ἀλόγως συγκεῖται. ἐν δὲ τοῖς ήμετέροις σώμασι χωρίς τῶν πρὸς ὑγείαν καὶ νόσον καὶ θάνατον καὶ πρὸς ἄλλα δυνάμεις ἔγειν φασίν οὐ μόνον τῶν σωματικῶν ἀλλά καὶ τῶν The Vuxne. .

ΧΙΧ. Πρός δε την ψυχην τον μεν στρύχνον ώστε παρακινείν καὶ εξιστάναι, καθάπερ ελέχθη πρότερον, ή δε τοῦ ονοθήρα ρίζα δοθείσα εν οἴνφ πραότερον καὶ ἱλαρώτερον ποιεῖ το ἡθος. ἔχει δε ὁ μεν ονοθήρας το μεν φύλλον ὅμοιον ἀμυγδαλή μικρότερον δε, τὸ δε ἄνθος ερυθρὸν ὥσπερ ρόδον αὐτὸς δε μέγας θάμνος ρίζα δε ερυθρὰ καὶ μεγάλη, ὅζει δε αὐανθείσης ὥσπερ οἴνου φιλει δε ορεινὰ χωρία. φαίνεται δε οὐ τοῦτο ἄτοπον οἶον γὰρ προσφορά τις γίνεται δύναμιν ἔχοντος

οινώδη.

1 σκορπίφ conj. W.; σκορπίου Ald.

§ 9. 11. 6.

² 18. 3, λέγω δὲ σωματικῶν . . . 18. 11 (the account of the physical effects) is here omitted.

ENQUIRY INTO PLANTS, IX. XVIII. 2-XIX. I

if it is shredded over him; while if one then sprinkles him with white hellebore, they say that he comes to life again. It is also fatal to oxen sheep beasts of burden and in general to any fourfooted animal, and kills them the same day if the root or leaf is put on the genitals; and it is also useful as a draught against a scorpion's sting. It has a leaf like cyclamen, and a root, as was said, like a scorpion. It grows like dog's-tooth grass, and is jointed, and it loves shady places. Now if what has been told already about the scorpion be true, then other similar tales are not incredible. (Indeed fabulous tales are not composed without some reason).2 And in relation to our own persons, apart from their effects in regard to health disease and death, it is said that herbs have also other properties affecting not only the bodily but also the mental powers.

Of plants possessing properties which affect the mental powers.

XIX. As to those which affect the mind, strykhnos, as was said before,³ is said to upset the mental powers and make one mad; ⁴ while the root of onotheras (oleander) administered in wine makes the temper gentler and more cheerful. This plant has a leaf like the almond, but smaller,⁵ and the flower is red like a rose. The plant itself (which loves hilly country) forms a large bush; the root is red and large, and, if this is dried, it gives off a fragrance like wine. And this does not seem surprising, since there is a sort of 'bouquet' given off by a thing which has the peculiar quality of wine.

4 Diosc. 4. 117; Plin. 26. 111.

⁶ μικρότερον conj. W.; πικρότερον UM; πλατύτερον Ald. (so also Diosc. l.c.). G seems to have read μακρότερον.

'Αλλὰ τάδε εὐηθέστερα καὶ ἀπιθανώτερα τά τε των περιάπτων καὶ όλως των άλεξιφαρμάκων λεγομένων τοις τε σώμασι και ταις οικίαις. και ώς δή φασι τὸ τριπόλιον καθ' Ἡσίοδον καὶ Μουσαίον είς παν πράγμα σπουδαίον χρήσιμον είναι, δι' δ και δρύττουσιν αὐτὸ νύκτωρ σκηνην πηξάμενοι, και τὰ περί τῆς εὐκλείας δὲ καὶ εὐδοξίας ὁμοίως ἡ καὶ μᾶλλον εὔκλειαν γάρ φασι ποιείν τὸ ἀντίρρινον καλούμενον τοῦτο δ' ὅμοιόν έστι τη ἀπαρίνη ρίζα δὲ οὐχ ὕπεστιν ὁ δὲ καρπός ώσπερ μόσχου ρίνας έχει. τον δ' άπο 8 τούτου άλειφόμενον εὐδοξεῖν, εὐδοξεῖν δὲ καὶ έάν τις του έλειοχρύσου τῷ ἄνθει στεφανῶται μύρω ραίνων εκ χρυσίου απύρου. έχει δε ό έλειόγρυσος τὸ μὲν ἄνθος γρυσοειδές, φύλλον δὲ λευκὸν καὶ τὸν καυλὸν δὲ λεπτὸν καὶ σκληρὸν ρίζαν δὲ ἐπιπόλαιον καὶ λεπτήν. χρῶνται δὲ αὐτῷ πρὸς τὰ δακετὰ ἐν οἴνω καὶ πρὸς τὰ πυρίκαυστα κατακαύσαντες καὶ μίξαντες μέλιτι. οὖν τοιαῦτα, καθάπερ καὶ πρότερον ἐλέχθη, συναύξειν βουλομένων έστι τὰς ξαυτών τέχνας.

Αί δὲ τῶν ριζῶν καὶ τῶν καρπῶν καὶ τῶν ὀπῶν
 φύσεις ἐπεὶ πολλὰς ἔχουσι καὶ παντοίας δυνάμεις, ὅσαι ταὐτὸ δύνανται καὶ τῶν αὐτῶν αἰτίαι.

1 ἀπιθανώτερα conj. Scal. after G: so also Cod. Cas. Vin. Vo.; πιθανώτερα U*; πιθανότερα Ald.

² τριπόλιον ÚMU*Ald.; G from Plin. 21. 44 has polium. It may be observed that τριπόλιον can hardly have occurred in a hexameter. Hesych., however, gives τρισπόλιον as the name

ENQUIRY INTO PLANTS, IX. xix. 2-4

Of plants said to have magical properties.

On the other hand what is said of amulets and charms in general for the body or the house is somewhat foolish and incredible. Thus they say that tripolion 2 according to Hesiod and Musaeus is useful for every good purpose, wherefore they dig it up by night, camping on the spot. So too what is said of good or fair fame as affected by plants is quite as foolish or more so: for they say that the plant called snapdragon produces fair fame. This plant is like bedstraw but it has no root: and the fruit has what resembles a calf's nostrils. The man who anoints himself with this they say wins fair fame. 4 And they say that the same result follows, if he crowns himself with the flower of gold-flower, sprinkling it with unguent from a vessel of unfired gold. The flower of gold-flower is like gold, the leaf is white. The stem also is white 5 and hard, the root is slender and does not run deep. 6 Men use it in wine against the bites of serpents, and to make a plaster for burns after burning it and mixing the ashes with honey. tales then, as was said before, proceed from men who desire to glorify their own crafts.

A problem as to cause and effect.

Now since the natural qualities of roots fruits and juices have many virtues of all sorts, some having the same virtue and causing the same result, while of a plant. Plin. l.c. seems to combine Diosc.'s account of πόλιον (3. 110) with his account of τριπόλιον (4. 132).

3 το ἀντίρρινον conj. St. from Diosc. 4. 130; Plin. 25. 129;

τὸ ἀντίρριζων Ald. H.; τὸν ἀντ. UM; τὸ ἀντίριζον U*.
4 Diosc. 4. 57; Plin. 21. 66. Cited also by Athen. 15. 27.

* λευκόν conj. Sch.; λεπτόν UMU*Ald.G.

⁶ Diosc. l.c.; Plin. 21. 168 and 169.

καὶ πάλιν ὅσαι τὰ ἐναντία, διαπορήσειεν ἄν τις κοινον ἴσως ἀπόρημα καὶ ἐφ' ἐτέρων ἀπόρων, πότερον όσα τῶν αὐτῶν αἴτια κατὰ μίαν τινὰ δύναμίν έστιν, ή καὶ ἀφ' έτέρων ἐνδέχεται ταὐτὸ γίνεσθαι, τοῦτο μεν οὖν ταύτη ἡπορήσθω εἰ δέ τινων καὶ ἄλλων τὰς φύσεις ἡ τὰς δυνάμεις έγομεν είπειν, ταθτα ρητέον.

ΧΧ. Τὸ δὴ πέπερι καρπὸς μέν ἐστι διττὸν δὲ αὐτοῦ τὸ γένος τὸ μὲν γὰρ στρογγύλον ὧσπερ όροβος, κέλυφος έγον καὶ σάρκα καθάπερ αί δαφνίδες, ὑπέρυθρου τὸ δὲ πρόμηκες μέλαν σπερμάτια μηκωνικά έχον ισχυρότερον δέ πολύ τοῦτο θατέρου θερμαντικά δὲ ἄμφω δι' δ καὶ πρὸς τὸ κώνειον βοηθεῖ ταῦτά τε καὶ ὁ λιβανωτός.

Ο δε Κυίδιος κόκκος στρογγύλον ερυθρούν τη χροιά μείζον δε τοῦ πεπέριος ἰσχυρότερον δε πολύ τη θερμότητι, δι' δ καὶ ὅταν δίδωσι κατάποτον. διδόασι γὰρ πρὸς κοιλίας λύσιν, ἐν ἄρτω ἡ στέατι περιπλάττοντες κάει γάρ ἄλλως τὸν φάρυγγα.

Θερμαντικόν δε και ή του πευκεδάνου <ρίζα,> δι' δ καὶ ἄλειμμά τι ποιοῦσιν έξ αὐτῆς ίδρωτικὸν ώσπερ καὶ ἐξ ἄλλων. δίδοται δὲ ἡ τοῦ πευκε-

² Cited by Athen. 2. 73; cf. Diosc. 2. 159. ³ Plin. 27. 70.

¹ ἀφ' conj. Sch.; ἐφ' U*P; Ald. omits the preposition.

ENQUIRY INTO PLANTS, IX. xix. 4-xx. 2

others have opposite virtues, one might raise a question which is perhaps equally perplexing in regard to other matters, to wit, whether those that produce the same effect do so in virtue of some single virtue which is common to them all, or whether the same result may not come about also from ¹ different causes.—Let us be content to put the question thus: but now we must proceed to speak of the natural qualities or virtues of any other plants that we can mention.

Of certain plants, not yet mentioned, which possess special properties.

XX. ² Pepper is a fruit, and there are two kinds: one is round like bitter vetch, having a case and flesh like the berries of bay, and it is reddish: the other is elongated and black and has seeds like those of poppy: and this kind is much stronger than the other. Both however are heating: wherefore these, as well as frankincense, are used as antidotes for poisoning by hemlock.

³ The 'Cnidian berry' is round, red in colour, larger than that of pepper, and far stronger in its heating power; wherefore, when it is given as a pill ⁴ (for it is given to open the bowels) they knead it up in a piece of bread or dough: otherwise it burns the

throat.

⁵ The root ⁶ of sulphur-wort is also heating, wherefore they make of it an ointment to produce a sweat, as with other things so used. This root ⁶ is also

6 bica add. W.

315

⁴ κατάποτον conj. Sch.; κατὰ πότον Ald. cf. καταπότιον 9. 8. 3. ⁵ cf. 9. 14. 1; Plin. 25. 117.

δάνου ρίζα καὶ πρὸς τοὺς σπληνας· τὸ δὲ σπέρμα οὐ χρήσιμον οὐδὲ ὁ ὀπὸς αὐτης· γίνεται δὲ ἐν 'Αρκαδία.

Δαῦκον δὲ περὶ Πατραϊκὴν τῆς 'Αχαΐας διαφέρον, θερμαντικὸν φύσει ἡίζαν δὲ ἔχει μέλαιναν.

Θερμαντικόν δὲ καὶ δριμύ καὶ τῆς ἀμπέλου τῆς ἀγρίας ρίζα· δι' δ καὶ εἰς ψίλωθρον χρήσιμον καὶ ἐφηλίδας ἀπάγειν· τῷ δὲ καρπῷ ψιλοῦσι τὰ δέρματα. τέμνεται δὲ πᾶσαν ὥραν ὀπώρας δὲ μάλιστα.

Δρακουτίου δε ρίζα βηχας εν μελιτι διδομένη παύειν χρησίμη. καυλον δε έχει ποικίλον οφιώδη:

σπέρματι δ' οὐ χρῶνται.

Ἡ δὲ τῆς θαψίας ἐμετική· ἐὰν δέ τις κατάσχη, καθαίρει καὶ ἄνω καὶ κάτω· δύναται δὲ καὶ τὰ πελιώματα ἐξαιρεῖν· ὑπώπια δὲ ποιεῖ ἄλλα ἔκλευκα. ὁ δὲ ὀπὸς ἰσχυρότερος αὐτῆς καθαίρει καὶ ἄνω καὶ κάτω· σπέρματι δ' οὐ χρῶνται· γίνεται δὲ καὶ ἄλλοθι μὲν ἀτὰρ καὶ ἐν τῆ Αττικῆ· καὶ τὰ βοσκήματα ταύτης οὐχ ἄπτεται τὰ ἐγχώρια, τὰ δὲ ξενικὰ βόσκεται καὶ διαρροίᾳ διαφθείρεται.

Τὸ δὲ πολυπόδιον μετὰ τὰ ὕδατα ἀναβλαστεῖ

σπέρμα δὲ οὐ φύει.

Το δε της εβένου ξύλον κατα μεν την πρόσοψιν δμοιον πύξω φλοϊσθεν δε μέλαν γίνεται· χρήσιμον δε προς όφθαλμίας ακόνη τριβόμενον.

cf. 7, 12, 2; Diosc. 2, 167. cf. Plin. 24, 89.
 Diosc. 4, 153; Plin. 13, 125 and 126.

¹ cf. 9. 15. 5. ² cf. 9. 15. 8. n.

³ cf. 9. 14. 1; Diosc. 4. 181-183; Plin. 23. 19 and 21.

ENQUIRY INTO PLANTS, IX. xx. 2-4

given for the spleen: but neither its seed nor its

juice is of use: it grows in Arcadia.1

² Daukon of excellent quality grows in the district of Patrai in Achaia, and is heating by nature: it has a black root.

³ The root of the 'wild vine' (bryony) is also heating and pungent: wherefore it is useful as a depilatory and to remove freckles: and the fruit is used for smoothing hides. It is cut at any season, but especially in autumn.

⁴ The root of edderwort given in milk is useful for stopping a cough. It has a variegated snake-like

stem: the seed is not used.

⁵ The root of thapsia has emetic properties: and, if one retains it, it purges both upwards and downwards. It is also able to remove bruises: and it restores other contusions to a pale colour.⁶ Its juice is stronger and purges both upwards and downwards: the seed is not used. It grows especially in Attica, but also in other places: the cattle of the country do not touch it, but imported cattle feed on it and perish of diarrhoea.⁷

⁸ Polypody springs up ⁹ after rain, and produces no

seed.

¹⁰ The wood of ebony is in appearance like box, but when barked it becomes black: it is useful against ophthalmia, and is rubbed on a whetstone for that use.

6 ὑπώπια . . . ἔκλευκα: text perhaps defective.

 7 διαρροία conj. Sch.; διάρροια \hbar U $\hat{\mathbf{M}}$: διάρροια αὐτοῖ \mathbf{s} γίνεται \hbar Ald.G.

⁸ cf. 9. 13. 6; C. P. 2. 17. 4. The account of the virtues of this plant is evidently missing.

⁹ ἀναβλαστεῖ conj. W.; αἰεὶ βάλλει Ald.

10 Diosc. 1. 98; Plin. 24. 89.

Ή δὲ ἀριστολοχία παχεῖα καὶ ἐσθιομένη πικρὰ τῷ χρώματι μέλαινα καὶ εὔοσμος, τὸ δὲ φύλλον στρογγύλον, οὐ πολὺ δὲ τὸ ὑπὲρ τῆς γῆς. φύεται δὲ καὶ μάλιστα ἐν τοῖς ὅρεσι· καὶ αὕτη βελτίστη. τὴν δὲ χρείαν αὐτῆς εἰς πολλὰ καταριθμοῦσιν· ἀρίστη μὲν πρὸς τὰ κεφαλόθλαστα, ἀγαθὴ δὲ καὶ πρὸς τὰ ἄλλα ἔλκη καὶ πρὸς τὰ ἐρπετὰ καὶ πρὸς ὕπνον καὶ πρὸς ὕστεραν ὡς πεσσός, τὰ μὲν σὺν ὕδατι ἀναδευομένη καὶ καταπλαττομένη, τὰ δ' ἄλλα εἰς μέλι ξυομένη καὶ ἔλαιον· τῶν δὲ ἐρπετῶν ἐν οἴνῷ ὀξίνη πινομένη καὶ ἐπὶ τὸ δῆγμα ἐπιπαττομένη· εἰς ὕπνον δὲ ἐν οἴνῷ μέλανι αὐστηρῷ κνισθεῖσα· ἐὰν δὲ αἱ μῆτραι προπέσωσι, τῷ ὕδατι ἀποκλύζειν. αὕτη μὲν οὖν ἔοικε διαφέρειν τῷ πολυχρηστίᾳ.

Τῆς δὲ σκαμμωνίας ὥσπερ ἐξ ἐναντίας ὁ ὀπὸς

μόνον χρήσιμος ἄλλο δ' οὐδέν.

Η δε της πτερίδος ρίζα μόνον τῷ χυλῷ γλυκύστρυφνος ελμινθα δε πλατείαν εκβάλλει· σπέρμα δε οὐκ εχει οὐδε όπόν· τέμνεσθαι δε ώραίαν μετο-

πώρου φασίν.

'Η δ' έλμις σύμφυτον ἐνίοις ἔθνεσιν· ἔχουσι γὰρ ὡς ἐπὶ πᾶν Αἰγύπτιοι 'Αραβες 'Αρμένιοι Ματαδίδες Σύροι Κίλικες· Θρᾶκες δ' οὐκ ἔχουσιν οὐδὲ Φρύγες· τῶν δὲ 'Ελλήνων Θηβαῖοί τε οἱ περὶ τὰ γυμνάσια καὶ ὅλως Βοιωτοί· 'Αθηναῖοι δ' οὔ.

Πάντων δὲ τῶν φαρμάκων ὡς ἀπλῶς εἰπεῖν βελτίω τὰ ἐκ τῶν χειμερινῶν καὶ προσβόρρων

cf. 9. 13. 3.
 καὶ μάλιστα conj. W.; μάλιστα καὶ Ald.
 αῦτη conj. Scal.; αὐτὴ Ald.

ENQUIRY INTO PLANTS, IX. xx. 4-5

¹ Birthwort is a stout plant and is bitter to the taste: it is black in colour and fragrant; the leaf is round. However there is not much of the plant above ground. It grows especially 2 on mountains, and then 3 it is best. Many uses of it for various purposes are enumerated; it is best for bruises on the head, good also for other wounds, against snake-bites, to produce sleep, for the womb as a pessary: for some purposes it is soaked with water and applied as a plaster, for others it is scraped into honey and oliveoil: against snake-bites it is drunk in sour wine and also sprinkled over the bite; to induce sleep it is given pounded up in black dry wine: 4 in cases of prolapsus uteri it is used in water as a lotion. plant then seems to have a surpassing variety of usefulness.

⁵ Of scammony, as though by contrast, only the

juice is useful and no other part.

Of male-fern no part but the root is useful and it has a sweet astringent taste. It expels the flat worm. It has no seed nor juice: and they say it

is ripe for cutting in autumn.

6 (This worm naturally infests certain races: speaking generally the following are liable to itthe Egyptians, the Arabians, the Armenians, the Matadides, the Syrians, the Cilicians: the Thracians have it not, nor the Phrygians. Among the Hellenes those Thebans who frequent wrestling-schools and the Boeotians generally are liable to it: but not the Athenians.)

Of all drugs, to speak generally, those are better which come from places that are wintry, face the

⁴ Cited by Apollon. *Hist. Mirab.* 29.

⁶ Plin. 27, 145.

καὶ ξηρῶν· δι' δ καὶ τῶν ἐν Εὐβοία τὰ ἐν ταῖς Αἰγαῖς ἡ τὰ ἐν τῷ Τελεθρίῳ φασί· ξηρότερα γάρ· τὸ δὲ Τελέθριον σύσκιον.

6 Περὶ μὲν οὖν τῶν ῥιζῶν ὅσαι φαρμακώδεις καὶ ὁποιασοῦν ἔχουσι δυνάμεις εἴτε ἐν αὐταῖς εἴτε ἐν τοῖς ὀποῖς ἡ καὶ ἄλλῷ τινὶ τῶν μορίων, καὶ τὸ ὅλον εἴ τι φρυγανικὸν ἡ ποῶδες ἔχει τοιαύτας δυνάμεις, καὶ περὶ τῶν χυλῶν τῶν τε εὐόσμων καὶ τῶν ἀόσμων καὶ ὅσας ἔχουσι διαφοράς, αἴπερ οὐθὲν ἡττον φυσικαί εἰσιν, εἴρηται.

ENQUIRY INTO PLANTS, IX. xx. 5-6

north and are dry: wherefore of those which grow in Euboea best, they say, are the drugs of Aigai or Telethrion, these places being dry, while Telethrion is also shady.

¹Thus we have spoken of drugs, those that are medicinal and those that have virtues of whatsoever kind, whether in the root itself, or in the juice, or in any other of their parts, and in general of all the shrubby or herbaceous plants which have such virtues, as well as their tastes, whether they be fragrant or without fragrance, with the differences between them, which are equally part of their essential character.

¹ This section begins a tenth book in UMAld.H.G; cf. 9. 8. 1 n. The concluding words can hardly represent the original text.

321

Y

MINOR WORKS

INTRODUCTION TO THE TREATISES CONCERNING ODOURS AND CONCERNING WEATHER SIGNS

THE text of the two opuscula given here is reprinted from that of Wimmer in the Teubner series, 1862, and in the Didot edition, 1866; the latter is very carelessly printed: a few slight alterations are mentioned in the notes. Both works are included in the Aldine edition (1497), and in that of Camotius (see p. x). For the de odoribus two MSS., Cod. Vaticanus (A) and Cod. Parisiensis (Q) were collated by Brandis. The text of the de signis is considered by Wimmer to be very corrupt and defective: he has admitted some emendations made by Schneider from an old Latin translation published at Bologna in 1516. Schneider's commentary makes frequent reference to an edition of the opuscula of Theophrastus by Turnebus and Daniel Furlanus, printed at Hanau in Prussia in 1605, and reprinted there in 1615.

The de signis was one of Aratus' authorities for his Diosemeia: I have only however made reference to that work where it appears to throw light on the text of Theophrastus. These and most other references for the two fragments I owe chiefly to Schneider.

CONCERNING ODOURS

CONTENTS

SECTIONS.

- 1-3. Introductory: Of odours in general and the classification of them.
 - Of natural odours: Of those of animals and of the effect of odours on animals.
 - 5. Of smell and taste.
 - 6. Of odours in plants.
- 7-13. Of artificial odours in general and their manufacture: especially of the use of perfumes in wine.
- 14-20. Of the oils used as the vehicle of perfumes.
- 21-26. Of the spices used in making perfumes and their treatment.
 - 27-31. Of the various parts of plants used for perfumes, and of the composition of various notable perfumes.
 - 32-35. Of the properties of various spices.
- 35-36. Of the medicinal properties of certain perfumes.
- 37-41. Of rules for the mixture of spices, and of the storing of various perfumes.
- 42-50. Of the properties of certain perfumes.
- 51-56. Of other properties and peculiarities of perfumes.
- 57-59. Of the making of perfume-powders and compound perfumes.
- 61-63. Of the characteristic smells of animals, and of certain curious facts as to the smell of animal and vegetable products.
- 64-69. Of odours as compared with other sense-impressions.

ΠΕΡΙ ΟΣΜΩΝ

Ι. Αἱ ὀσμαὶ τὸ μὲν ὅλον ἐκ μίξεώς εἰσι καθάπερ οἱ χυμοί· τὸ γὰρ ἄμικτον ἄπαν ἄοδμον ὅσπερ ἄχυμον, διὸ καὶ τὰ ἀπλᾶ ἄοδμα, οἱον ὕδωρ ἀὴρ πῦρ· ἡ δὲ γῆ μάλιστ' ἡ μόνη ὀδμὴν

έχει διὸ μάλιστα μικτή.

Των δ' όδμων αί μὲν ὅσπερ ἀειδεῖς καὶ ὑδαρεῖς καθάπερ ἐπὶ των χυμων, αἰ δ' ἔχουσαί τινας ἰδέας. αἱ δ' ἰδέαι δοκοῦσι μὲν ἀκολουθεῖν ταῖς των χυμων, οὐ μὴν ἔχουσί γε πᾶσαι τὰς αὐτὰς προσηγορίας, ὥσπερ ἐν τοῖς πρότερον εἴπομεν, οὐδ' ὅλως οὕτω διωρισμέναι τοῖς εἴδεσιν ὥσπερ οἱ χυμοὶ ἀλλ' ὡς ᾶν τοῖς γένεσιν, ὅτι τὰ μὲν 2 εὕοσμα τὰ δὲ κάκοσμα. τῆς δ' εὐωδίας καὶ κακωδίας οὐκέτι τὰ εἴδη κατωνόμασται καίπερ ἔχοντα διαφορὰς μεγάλας ἐπί γ' αὐτῶν τῶν γλυκέων καὶ πικρῶν, ἀλλὰ δριμεῖα λέγεται καὶ ἰσχυρὰ καὶ μαλακὴ καὶ γλυκεῖα καὶ βαρεῖα ὀδμή·κοιναὶ δ' ἔνιαι τούτων καὶ τῶν κακωδῶν.

¹ i.e. there is not one set of terms applied to the varieties of 'good' and another distinct set applied to the varieties of 'evil' odours, but we get a cross-division, some terms (such as 'strong') being applied to varieties of both classes. cf. 64-66.

CONCERNING ODOURS

Introductory: Of odours in general and the classification of them.

I. Opours in general, like tastes, are due to mixture: for anything which is uncompounded has no smell, just as it has no taste: wherefore simple substances have no smell, such as water air and fire: on the other hand earth is the only elementary substance which has a smell, or at least it has one to a greater extent than the others, because it is of a

more composite character than they.

Of odours some are, as it were, indistinct and insipid, as is the case with tastes, while some have a distinct character. And these characters appear to correspond to those of tastes, yet they have not in all cases the same names, as we said in a former treatise; nor in general are they marked off from one another by such specific differences as are tastes: rather the differences are, one may say, in generic character, some things having a good, some an evil odour.1 But the various kinds of good or evil odour, although they exhibit considerable differences, have not received further distinguishing names, marking off one particular kind of sweetness or of bitterness from another: we speak of an odour as pungent, powerful, faint, sweet, or heavy, though some of these descriptions apply to evil-smelling things as well as to those which have a good odour.

'Η δὲ καθόλου καὶ ὥσπερ ἐπὶ πᾶσι τοῖς δια-Φθειρομένοις σαπρότης. άπαν γάρ τὸ σηπόμενον κακώδες, εί μή τις την όξύτητα λέγει του οίνου 3 σαπρότητα τη όμοιότητι της φθοράς. ἐν ἄπασι δ' έστλν ή τοῦ σαπροῦ κακωδία καλ έν φυτοῖς καὶ ἐν ζώοις καὶ ἐν τοῖς ἀψύγοις ἐν ἄπασι δὲ διαφθειρομένοις ών μη η σύστασις εὐθὺς ἐκ τοιαύτης ύλης έγει γαρ ένια και την της ύλης οσμήν, ου μην έπὶ πάντων γε τοῦτ' ἀκολουθεῖ. πολλά γάρ οὐ κακώδη τὰ ἐκ τῶν σαπρῶν, ὡς ούδ' οἱ μύκητες οἱ ἐκ τῆς κόπρου φυόμενοι τὰ δ' έκ σήψεως φυόμενα καὶ συνιστάμενα κακώδη. εύοσμα μεν ούν ώς άπλως είπειν τα πεπεμμένα καὶ λεπτὰ καὶ ἥκιστα γεώδη τὸ γὰρ τῆς ὀσμῆς έν άναπνοη κακώδη δε δηλονότι τάναντία. πολλά δὲ ὥσπερ τῶν γλυκέων ἐμφαίνει τινὰ πικρότητα, καὶ τῶν εὐωδῶν βαρύτητα ταῖς ὀσμαῖς.

 II. "Εχει δὲ ἔκαστον ὀσμὴν ἰδίαν καὶ ζώων καὶ φυτῶν καὶ τῶν ἀψύχων ὅσα ὀσμώδη· πολλὰ

²Which is not an 'evil' odour.

The sense is apparently that 'lighter' (or less solid) things exhale a lighter and pleasanter odour because in their 328

¹ And so here we have a term which possibly is applied only to the one class of 'evil' odours.

³ i.e. putridity is a quality which things acquire as they decay, and does not necessarily imply that they are themselves formed out of decaying matter. In fact things so produced are not always 'putrid.'

CONCERNING ODOURS, 2-4

Putridity however is a general term, applied, one may say, to anything which is subject to decay 1: for anything which is decomposing has an evil odour,unless indeed the name putridity be extended to sourness 2 in wine because the change in the wine is analogous to decomposition. The evil odour of putridity is found in all things, alike in plants in animals and in inanimate things: it attends the decay of things which are not formed directly out of a substance which is decaying: for some things have also the odour of that substance, though it is not found in every case.3 Thus in many instances things which are produced by decaying matter have no evil odour: for instance, mushrooms which grow from dung have none: but things which grow from decay and are actually formed out of it have such an odour. To speak generally then, things that have been cooked, delicate things, and things which are least of an earthy nature have a good odour,4 (odour being a matter of exhalation), and it is obvious that those of an opposite character have an evil odour. But, even as many things pleasant to the taste present a certain bitterness, so many things that have a good odour have a kind of heavy scent.

Of natural odours; of those of animals and of the effect of odours on animals.

II. Every plant animal or inanimate thing that has an odour has one peculiar to itself: but in many

case exhalation is easier. The sense given to ἀναπνοή requires illustration (the passages cited by LS. are not in point). Sch., construing apparently as W. does, 'since smell depends on breathing' (? inhalation), admits that he does not see the point of this clause.

δ΄ ήμιν οὐ φαίνεται διὰ τὸ χειρίστην ἔχειν τὴν αἴσθησιν ταύτην ὡς εἰπεῖν. ἐπεὶ τοῖς γε ἄλλοις καὶ τὰ παντελῶς ἄοδμα φαινόμενα δίδωσί τινα ὀσμήν, ὥσπερ αἰ κριθαὶ τοῖς ὑποζυγίοις αὶ ἐκ τῆς Κεδροπόλιος, ἃς οὐκ ἐσθίουσιν διὰ τὴν κακωδίαν. ἡμᾶς δὲ καὶ αὶ τῶν ζώων λανθάνουσιν τῶν ὀσμωδῶν δοκούντων. εὐωδία μὲν οῦν οὐθὲν φαίνεται καθ' αὐτὸ χαίρειν ὡς εἰπεῖν, ἀλλ' ὅσα πρὸς τὴν τροφὴν καὶ τὴν ἀπόλαυσιν. πονεῖν δ' ἔνια φαίνεται ταῖς ὀσμαῖς καὶ ταῖς εὐωδίαις, εἴπερ ἀληθὲς τὸ ἐπὶ τῶν γυπῶν καὶ τῶν κανθάρων. τοῦτο δὲ δῆλον ὡς δι' ἐναντίωσιν τῆς ἐν αὐτοῖς φύσεως. ὡς δὲ καθ' ἔκαστον ἄμα δεῖ τὴν γε κρᾶσιν τὴν ἐκάστου καὶ τὴν τῆς ὀσμῆς λαμβάνειν δύναμιν.

Είσι μεν οὖν ἔνιαι τῶν εὐόσμων καὶ ἐν ταῖς τροφαῖς, οἶον αἱ τῶν ἀκροδρύων καὶ ἀπίων καὶ μήλων αἰται γὰρ ἄνευ τῆς προσφορᾶς ἡδεῖαι, καὶ μᾶλλον ὡς εἰπεῖν. οὐ μὴν ἀλλ' ὡς γ' ἀπλῶς διελεῖν αἱ μέν εἰσι καθ' αὐτὰς αἱ δὲ κατὰ συμβεβηκός αἱ μὲν τῶν χυλῶν καὶ τῆς τροφῆς κατὰ συμβεβηκός, αἱ δ' ὥσπερ τῶν ἀνθῶν καθ' αὐτάς. ὡς δ' ἐπίπαν τὰ εὔοσμα, καθάπερ καὶ πρότερον ἐλέχθη, δύσχυμα καὶ στρυφνὰ καὶ ὑπόπικρα.

² εὐωδίαις. ? εὐώδεσι.

¹ In Thrace. cf. Arist. H.A. 9. 36. Turn. quotes an illustration from Scriptor θαυμασίων ἀκουσμάτων 126.

CONCERNING ODOURS, 4-5

cases it is not obvious to us because, one might almost say, our sense of smell is inferior to that of all other animals. Thus things which appear to us to have no odour give forth an odour of which other animals are conscious: for instance beasts of burden can smell the barley of Kedropolis, and refuse to eat it because of its evil odour. Also we are unaware of the odour of animals which appear to possess one. Now no animal appears to take pleasure in a good odour for its own sake, so to speak, but only in the odour of things which conduce to its nurture and enjoyment. Indeed some animals seem to be annoved by odours, even good 2 ones, if what is said of vultures and beetles be true; the explanation is that their natural character is antipathetic to odours. To appreciate this in particular cases one should take into consideration the temperament of the animal in question and also its power of smell.

Of smell and taste.

Now the odour of some things which have a good odour resides in things which are used for food, for instance that of stone-fruits 8 pears and apples, the smell of which is sweet even if one does not eat them; indeed it may be said to be sweeter in that case. However, to make a general distinction, some odours exist independently, while others are incidental; those of juices and things used for food are incidental, those of flowers exist independently. And, as was said above, things which have a good odour are generally of unpleasant, astringent or

 ^{*} ἀκροδρύων here apparently plums, peaches, etc.
 * i.e. the smell is a kind of 'accident,' or by-product of te taste.
 * 1. 3. the taste.

ένια δὲ τῶν εὐχύμων καὶ κακώδη, καθάπερ καὶ τὸ Αἰγύπτιον καλούμενον σῦκον, γλυκὰ ὄν, καὶ εἰ μὴ πανταχοῦ ἀλλ' ἐνιαχοῦ. καὶ ἡ ἄρκευθος ἐμφαίνει τινὰ τῷ μασήσει κακωδίαν γλυκεῖα οὖσα τὸ δ' οὖρον ποιεῖ εὐῶδες.

Έπεὶ δὲ τῶν ὀσμῶν αἱ μὲν ἐν φυτοῖς καὶ τοῖς τούτων μορίοις, οἰον κλωσὶ φύλλοις φλοιοῖς καρποῖς δακρύοις, αἱ δὲ ὅσπερ διείλομεν ἐν ζώοις [καὶ φυτοῖς] καὶ τοῖς ἀψύχοις, αὖται μὲν φανερὸν ὅτι πέψιν ἔκασται λαμβάνουσιν ἐν τοῖς οἰκείοις [αἰς]· καὶ τὸ εὐῶδες καὶ κακῶδες ἀκολουθεῖ κατὰ τὰς οἰκείας φύσεις, ἡ δὲ πέψις τῷ οἰκείφ θερμῷ. ἐν δὲ τοῖς ἀψύχοις ταῖς τῶν ἀπλῶν δυνάμεσι καὶ γίνονται καὶ μεθίστανται καθάπερ οἱ χυμοί.

ΙΙΙ. "Οσαι δὲ δὴ κατὰ τέχνην καὶ ἐπίνοιαν γίνονται περὶ τούτων πειρατέον εἰπεῖν ὥσπερ καὶ περὶ τῶν χυλῶν. ἐν ἀμφοῖν δὲ δῆλον ὡς ἀεὶ πρὸς τὸ βέλτιον [ἦν] ἡμῖν ἡ ἀναφορά· πᾶσα γὰρ τέχνη στοχάζεται τούτου. εἰσὶ μὲν οὖν καὶ τοῖς ἀμίκτοις ὀσμαί τινες πρὸς τις συνεργεῖν πειρῶνται καὶ ταῖς παρα<σκευαῖς, ὡς καὶ> πρὸς τὰς τῶν χυμῶν εὐστομίας. οὐ μὴν ἀλλ' ὡς γ' ἀπλῶς

1 cf. H.P. 1. 11. 2.

4 als I omit; ή και τὸ εὐ. conj. Turn.

² i.e. the berry: Sch. would read ἀρκευθίς. cf. H. P. 3. 12. 4, with which this statement is inconsistent. Sch. suggests punctuating—γλυκὺ δτ. καὶ εἰ μὴ πανταχοῦ ἀλλ' ἐνιαχοῦ καὶ ἡ ἄρκευθος κ.τ.λ.
3 καὶ φυτοῖς om. Turn.

CONCERNING ODOURS, 5-7

somewhat bitter taste. Again some things which have a good taste have also an evil odour, such as the carob, which is sweet (this is true of some regions, if not of all). Again the Phoenician cedar, though it is sweet to the taste, when chewed produces a sort of evil odour, though it makes the water fragrant.

Of odours in plants.

Some odours being found in plants or in their parts—as twig, leaf, bark, fruit, gum—and others, as we distinguished, in animals and in inanimate things, it is plain that the former are matured each of them in the part to which it belongs; and a good or evil odour follows according to the natural character of that part, the maturing being due to the warmth which is found in it. On the other hand in inanimate things the odour, like the taste, is formed and modified by the properties of the simple substances of which the thing is made.

Of artificial odours in general and their manufacture: especially of the use of perfumes in wine.

III. Next we must endeavour to speak of those odours, and also those tastes, which are artificially 5 and deliberately produced. In either case it is clear that improvement is always what we have in view; for that is the aim of every artificial process. Now even uncompounded substances have certain odours, which men endeavour to assist by artificial means,6 even as they try to assist nature in producing palatable tastes. However, to speak generally, the

⁵ The same phrase occurs in similar connexion C.P. 6. 11. 2.

333

Text defective. Ald, Bas. Vo. have marks of omission. W. after Turn. gives καὶ ταῖς παρα < σκευαῖς, ὡς καὶ > ταῖς τῶν χυμῶν εὐστο < μίαις >, which I have slightly altered.

είπειν εν μίξει το πλέον, και οὕτως αι <μίξεις> δυοιν μεν ως τῷ γένει λαβειν, ὑγροῦ και ξηροῦ· τριχῶς δε γί<νονται>, ὅταν ἡ ὁμογενες ὁμογενει, ἡ παράλλαττον τῷ παραλλάττοντι, ἡ ὑγρῷ ὑγρὸν ἡ ξηρῷ ξηρόν>.

Έκ δυοῖν γὰρ τούτων καὶ ἡ τῶν χυλῶν καὶ τῶν ὀσμῶν γένεσις ὡς μὲν οἱ τὰ ἀρώματα καὶ τὰ διαπάσματα συντιθέντες ξηροῖς πρὸς ξηρά ὡς δ' οἱ τὰ μύρα κεραννύντες ἡ τῷ οἴνῷ ἐπιχέοντες ὑγροῖς πρὸς ὑγρά. τὸ δὲ τρίτον, ὁ καὶ πλεῖστόν ἐστιν, ὡς οἱ μυρεψοὶ ξηροῖς πρὸς ὑγρά παντὸς γὰρ μύρου καὶ χρίσματος ἡ σύνθεσις αὕτη. δεῖ δ' εἰδέναι ποῖαι ποίοις εὔμικτοι καὶ ποῖαι ποίοις συνεργοῦσιν εἰς τὸ ποιεῖν μίαν ὡσπερ ἐπὶ τῶν χυμῶν. καὶ γὰρ ἐκεῖ ταὐτὸ τοῦτο ζητοῦσιν οἱ μιγνύντες καὶ οἷον ἀρτύοντες. ταῦτα μὲν οὖν ἐν οἶς καὶ δι' ὧν αἱ τέχναι ποιοῦνται τὰ τέλη.

Μίγνυνται δὲ τὰ μὲν αὐτῆς τῆς ὀσμῆς ἔνεκα καὶ πρὸς ταύτην τὴν αἴσθησιν, τὰ δ' ὥσπερ ἡδύνειν βουλόμενα τὴν γεῦσιν, οἶον ὡς οἰ τὰ μύρα τοῖς οἴνοις ἐπιχέοντες ἡ τὰ ἀρώματα ἐμβάλλοντες.

¹ I have supplied $\mu l \xi \epsilon i s$ to fill the lacuna marked by W. after οδτωs αl : the text to the end of the section is defective, but a makeshift restoration and rendering seem possible: the sense of οδτωs is obscure.

CONCERNING ODOURS, 7-9

result is usually obtained by a mixture, and accordingly 1 such mixtures are of two things (or classes of things), a liquid and a solid: but there are three ways in which the result may be reached (the combination 2 being one either of like with like, or of unlike substances), according as a liquid is compounded with another liquid, a solid with another

solid, or a solid with a liquid.

For tastes and odours alike are derived from these two things: the method of the makers of spices and perfume-powders is to mix solid with solid, that of those who compound unguents or flavour wines is to mix liquid with liquid: but the third method, which is the commonest, is that of the perfumer, who mixes solid with liquid, that being the way in which all perfumes and ointments are compounded. Further one must know which odours will combine well with which, and what combination makes a good blend, just as in the case of tastes: for there too those who make combinations and, as it were, season their dishes, are aiming at this same object. So much for the ingredients and the methods whereby these arts attain their ends.

The object of the mixture is in the one case simply the production of a particular odour and the gratification of the corresponding sense, in the other there is a desire to produce, as it were, a pleasanter taste: this for instance is the object of flavouring wine with perfumes or of putting spices into it.

³ διαπάσματα. cf. Plin. 13. 19; 21. 125.

² i.e. given two components we have three possible combinations, A with A, B with B, or A with B.

⁴ The difference between μύρου and χρίσματοs does not appear; μύρου seems to be loosely used, as just above it was used of an entirely liquid mixture.

αί γὰρ αἰσθήσεις σύνεγγυς οὖσαι ποιοῦσί τινα ἀπόλαυσιν ἀλλήλων, ὅθεν καὶ αὐτοῖς τοῖς ψευστοῖς ζητοῦσι τὰς εὐοσμίας.

'Απορήσειε δ' ἄν τις ἴσως διὰ τί ποτε μύρον καὶ τάλλα εὔοσμα τοὺς μὲν οἴνους ἡδύνει τῶν δὲ Βρωμάτων οὐδέν, ἀλλὰ πάντα λυμαίνεται καὶ άπύρωτα καὶ πεπυρωμένα. τὸ δ' αἴτιον ύποληπτέον ὅτι συμβαίνει τῶν μὲν ξηρῶν ἀφαιρεῖσθαί τε τὸν οἰκεῖον χυλὸν διὰ τὴν ἰσχὺν καὶ άμα συνεπιφαίνειν τὸν αύτοῦ ὄντα στρυφνὸν καὶ ὑπόπικρον ἄπαν γὰρ τὸ εὔοσμον τοιοῦτον. διαμασωμένοις δὲ καὶ μᾶλλον ἐμφανὲς διά τε τὴν 11 θλίψιν καὶ τομὴν καὶ ἔτι τῷ γρονίζεσθαι. τὸν δ' οίνον οὐδέτερον ποιεί· καὶ γὰρ ὁ χυλὸς ἰσχυρότατος καὶ πλείων εἰς τὸ μὴ κρατεῖσθαι καὶ οὐδένα τη γεύσει χρόνον ἐπιδιατρίβων ἀλλ' ὅσον ἐπίθιγγάνων, ώστε τὸ μὲν ἡδὺ ἐνδιδόναι τῆ αἰσθήσει τὸ δὲ πικρὸν καὶ δύσχυμον τῆ γεύσει μὴ ἐμφαινειν, άλλα συμβαίνειν τώ οντι καθάπερ ήδυσμα γίνεσθαι τῷ πόματι τὴν ὀσμήν τῷ μὲν γὰρ γλυκεί και μάλιστα δεομένω διά το μηδεν έχειν, τοίς δ' άλλοις ώσπερ μιας έξ αμφοίν γενομένης

I have restored καl, which Sch. and W. omit, missing the point of the antithesis μέν...δέ.

¹ cf. 67; Arist. de Sens. 5. ² cf. Arist. l.c.

³ As opposed to wine. Sch., misunderstanding this, thinks Enpar corrupt.

CONCERNING ODOURS, 9-11

¹ For the two senses of taste and smell being akin to one another, each provides in a way for the enjoyment of the other: wherefore it is through things which appeal to the taste, as well as those which appeal to the sense of smell, that men try to discover

fragrant odours.

The question may perhaps be raised why perfume and other fragrant things, while they give a pleasant taste to wine,2 yet have not this effect on any other article of food, but in all cases spoil food, whether it be cooked or not. The explanation we must take to be that this is what happens—the perfume if mixed with solid 3 things is in any case powerful enough to deprive them of their proper taste, and at the same time it makes obtrusive its own taste. which is astringent and somewhat bitter.--all perfumeries having that character,-while, if one bites up the food, this effect is even 4 more apparent because the food is crushed and broken up, and also because it remains longer in the mouth. But on wine neither effect is produced, since in this the taste is very strong and too generally diffused to be overpowered: also wine does not linger on the palate for any length of time, but merely touches it, so that, while it makes one conscious of its own pleasant taste, it does not make the palate feel the bitter unpalatable taste of the perfume: in fact the odour of this acts as a sort of relish 5 to the draught. This effect indeed 6 it has on wine which is sweet and specially needs the addition of perfume, because it has no 'relish' of its own; while with other wines the reason is that, as the effect of the mixture, the

337

VOL. II.



z

⁵ Sc. 'bouquet.'

⁶ I have restored γàρ, omitted by Sch. and W.

διὰ τὴν μίξιν. ὁ γὰρ οἶνος, ώσπερ καὶ πρότερον ἐλέχθη, δεινὸς δέξασθαι τὰς ὀσμάς.

"Εχει δ' ἀπόρησιν καὶ τόδε, διὰ τί τὰ μὲν άνθη καὶ τὰ στεφανώματα ἀσθενέστερα ὄντα ταῖς όσμαις και πόρρωθεν όζει, ή δ' ίρις και τὸ νάρδον καὶ τάλλα τὰ εὔοσμα τῶν ξηρῶν ἰσγυρότερα έγγύθεν καὶ ένιά γε προσενεγκαμένοις, ένια δὲ καὶ τρίψεως προσδείται καὶ διαιρέσεως, τὰ δὲ καὶ πυρώσεως, ώσπερ ή σμύρνα καὶ ὁ λιβανωτὸς 18 καὶ πᾶν τὸ θυμιατόν, αἴτιον δ' ὅτι τῶν μὲν ανθών επιπολής το ποιούν την όσμην ατε μανών όντων καὶ οὐκ ἐγόντων βάθος, τῶν δὲ ῥιζῶν καὶ πάντων τῶν στερεῶν ἐν βάθει, τὰ δ' ἔξωθεν άπεξηραμμένα και πεπυκνωμένα διο και άφιασι πόρρω τὰς ἀποπνοίας, τὰ δ' οἱον ἀνοίξεως δέονται τῶν πόρων, ὅθεν διαιρούμενα καὶ κοπτόμενα πάντ' εὐωδέστερα, τὰ δ' ἄνθη κακωδέστερα τριβόμενα τὰ μὲν γὰρ ἐκφαίνει τὸ οἰκεῖον τὰ δὲ προσλαμβάνει τὸ ἀλλότριον, ὁ δὲ λιβανωτὸς καὶ ή σμύρνα πυκνοτέραν ἔτι τὴν φύσιν ἔχοντα προσδέονται πυρώσεως μαλακής, ή κατά μικρόν έκθερμαίνουσα ποιήσει την αναθυμίασιν. γαρ κόπτη τις ή τρίβη ταῦτα, προσοίσονται μέν

338

i.e. of the unadulterated wine and of the perfume.

² C.P. 6. 19. 2. Sch.'s reasons for bracketing this sentence seem inadequate.

³ i.e. fragrant leaves, etc. cf. H.P. 1. 12. 4.

⁴ Made from the rhizomes: cf. H.P. 1. 7. 2, and Index.

CONCERNING ODOURS, 11-13

two 1 odours combine, as it were, to form one. Wine indeed, as was said before, 2 has a special property of

assimilating odours.

Another question also suggests itself.—why it is that, while the smell of flowers and other 3 things used for garlands, though it is not so strong, can be perceived even at a great distance, the iris-perfume,4 spikenard and other fragrant solids smell stronger at a short distance: and of some of these the smell is only perceived when they are eaten, while some need even to be bruised and broken up, and others to be subjected to fire, as myrrh frankincense and anything that is burnt as incense. The explanation is that, whereas in flowers that which causes the smell is on the surface, seeing that the texture of flowers is open and they are not substantial, in all such solid substances as roots the power of producing smell is diffused through a substantial mass, while the exterior parts are dried up and of close texture : and this is why flowers emit the scent which exhales from them to a long distance, while things like roots need an opening of their passages. Hence, when these are broken up or bruised, they are in all cases more fragrant, while, 5 if flowers are crushed, they have a comparatively evil smell: for under such treatment roots give forth the property which belongs to them, but flowers acquire a property which is not their Again frankincense and myrrh, since they are by nature of even closer texture than roots, need a gentle application of fire, which, by gradually warming them, will cause the scent to be exhaled. For, if these substances are bruised or crushed, they will indeed present an odour, but it will not be so

⁵ cf. Arist. Probl. 12. 9; 13. 3 and 11.

όσμην ούχ όμοίως δὲ ήδεῖαν οὐδ' εὐταμίευτον. τούτων μέν οθν τοιαθταί τινες αι αιτίαι.

- ΙΥ. Των δε μύρων ή σύνθεσις και ή κατασκευή τὸ όλον οίον εἰς θησαυρισμόν ἐστι τῶν ὀσμῶν. διόπερ είς τούλαιον τίθενται τοῦτο γὰρ χρονιώτατον καὶ ἄμα πρὸς τὴν χρείαν μάλισθ' άρμόττον. έπει φύσει ηκιστα δεκτικον όσμης δια την πυκνότητα καὶ τὸ λίπος, αὐτῶν δὲ τούτων τὸ λιπαρώτατον, οίον τὸ ἀμυγδάλινον τὸ δὲ σησάμινον καὶ τὸ ἐκ τῶν ἐλαιῶν μάλιστα.
- 15 Χρώνται δὲ μάλιστα τω ἐκ τῆς βαλάνου τῆς Αίγυπτίας καὶ Συρίας, ήκιστα γὰρ λιπαρόν ἐπεὶ καὶ τῶ ἐκ τῶν ἐλαιῶν μάλιστα γρῶνται τῶ ωμοτριβεί της φαυλίας δοκεί γαρ άλιπέστατον έχειν καὶ λεπτότατον καὶ τούτω νέω καὶ μὴ παλαιώ το γαρ υπέρ ενιαυτον άχρειον παχύτερον καὶ λιπαρώτερον γενόμενον. έλαιον μέν οθν τὸ τοιοθτον οἰκειότατον, ἀλιπέστατον γάρ. φασί δέ τινες καὶ <έν> τῷ χρίσματι τὸ ἐκ τῶν πικρών άμυγδάλων πολλά δε γίνεται περί Κιλι-16 κίαν καὶ ποιούσιν έξ αὐτών χρίσμα. φασὶ δὲ καὶ εἰς τὰ σπουδαῖα τῶν μύρων άρμόττειν, ὥσπερ καὶ τὸ ἐκ τῆς βαλάνου καὶ αὐτό ποιεί δὲ <τὰ> κελύφη αὐτῶν εὔοσμον εἰς τὸ ἔλαιον ἐμβαλλό-

See Index.

¹ This passage was misunderstood by Plin. 13. 19. The sense seems to be that the viscous character of oil, though preservative of perfume, is not easily receptive of it.

² cf. H.P. 4. 2. 1; 4. 2. 6. βάλανος, balanites aegyptiaca.

CONCERNING ODOURS, 13-16

sweet nor so lasting as when they have been subjected to fire. Such are the explanations of these difficulties.

Of the oils used as the vehicle of perfumes.

IV. Now the composition and preparation of perfumes aim entirely, one may say, at making the odours last. That is why men make oil the vehicle of them, since it keeps a very long time and also is most convenient for use. ¹ By nature indeed oil is not at all well suited to take in an odour, because of its close and greasy character: and of particular oils this is specially true of the most viscous, such as almond-oil, while sesame-oil and olive-oil are the least receptive of all.

The oil most used is that derived from the Egyptian 2 or Syrian balanos, since this is the least viscous; the olive-oil which is most used is that which is pressed from 'coarse olives's in the raw state, since this is thought to be the least greasy and the least coarse: this is used while it is new. not when it is old, for that which is kept above a year is useless, having become thick and viscous. This then is the kind of olive-oil which is most suitable, since it is the least greasy. Some say that for unguent the oil derived from bitter almonds is best: these are abundant in Cilicia, where an unguent is made from them. It is said that this is suitable for choice perfumes, like the oil of the Egyptian balanos: this is suitable in itself,4 however the shells of the fruit are thrown into the oil to give it a good odour: indeed they are also thrown into

⁴ αὐτὸ conj. Sch.; τοῦτο Vulg.W.

ef. H.P. 2. 2. 12; C.P. 6. 8. 3 and 5.

μενα· έπεὶ καὶ τὸ τῶν πικρῶν. ἤδη δὲ πῶς οὐκ ἐναντίον ἄμα μὲν τὸ ἀοσμότατον ζητεῖν, ὥσπερ καὶ τὸ ἀμοτριβὲς ἐκ τῶν φαυλιῶν, ἄμα δ' ἐν τούτοις ποιεῖν; δριμύτητα γὰρ ἔχει τὸ τῶν ἀμυγδάλων· εἰ μὴ ἄρ' ὅτι τὸ ἔλαιον ἑψόμενον κακῶδες. ταῦτα μὲν οὖν ἐπισκεπτέον.

17 Χρώνται δὲ πρὸς πάντα τοῖς ἀρώμασι, τοῖς μὲν ἐπιστύφοντες τὸ ἔλαιον τοῖς δὲ καὶ τὴν ὀσμὴν ἐκ τούτων ἐμποιοῦντες. ὑποστύφουσι γὰρ πᾶν εἰς τὸ δέξασθαι μᾶλλον τὴν ὀσμήν, ὥσπερ τὰ ἔρια εἰς τὴν βαφήν. ὑποστύφεται δὲ τοῖς ἀσθενεστέροις τῶν ἀρωμάτων, εἶθ ὕστερον ἐμβάλλουσιν ἀφ' οὖ ἄν βούλωνται τὴν ὀσμὴν λαβεῖν ἐπικρατεῖ γὰρ ἀεὶ τὸ ἔσχατον ἐμβαλλόμενον καὶ ἄν ἔλαττον ἢ οἶον ἐὰν εἰς κοτύλην σμύρνης ἐμβληθῆ μνᾶ καὶ ὕστερον ἐμβληθῶσι κιναμώμου δραχμαὶ δύο, κρατοῦσιν αἱ τοῦ κιναμώμου δύο δραχμαί.

18 Θαυμάσειε δ' ἄν τις ἴσως τοῦτό τε καὶ διὰ τί ποτε τὰ ἀρώματα προεμβαλλόμενα δεκτικώ-τερον ποιεῖ τοὕλαιον ὀσμὴν ἔχοντα· δεῖ γὰρ ἀῶδες εἶναι τὸ δεξόμενον, τὸ δὲ κατειλημμένον ὑφ' ἐτέρου οὐκ ἀῶδες, ὥσθ' ἦττον ἐχρῆν εἶναι δεκτικόν. αἴτιον δ' ἀμφοτέρων ἢ πάντων τὸ αὐτό. ξηρὰ γὰρ ὄντα τὸ λίπος ἔλκει πρὸς ἑαυτὰ

¹ τὸ conj. Sch.; τὰ Vulg. W. Sch. also adds ἀμυγδάλων after πικρῶν.

² i.e. those derived from the Egyptian balanos and bitter almonds.

CONCERNING ODOURS, 16-18

that 1 which is made from bitter almonds. Once more, is it not inconsistent to seek the vehicle which has the least odour of its own, such as the oil which is pressed raw from 'coarse olives,' and yet at the same time to use the above-mentioned 2 oils as vehicles? (for oil of almonds has a pungent smell). Possibly the explanation is that it is only by being cooked that oil acquires an evil smell.³ These

matters then are subject for enquiry.

They use spices in the making of all perfumes; some to thicken 4 the oil, some in order to impart their odour. For in all cases they thicken the oil to some extent to make it take the odour better, just as they treat wool for dyeing. The less powerful spices are used for the thickening, and then at a later stage they put in the one whose odour they wish to secure. ⁵ For that which is put in last always dominates, even if it is in small quantity; thus, if a pound of myrrh is put into a half-pint of oil, and at a later stage a third of an ounce of cinnamon is added, this small amount dominates.

At this one may well wonder; and also why it is that the previous addition of spices, which have an odour of their own, renders the oil more receptive: for the vehicle should be scentless, but a substance over which another substance has thus prevailed, cannot be scentless, so that it ought, one would think, to have become *less* receptive. However both facts, or rather all of them, may be accounted for in the same way:—the spices, being solid, attract to

4 i.e. to make it less volatile. 5 cf. Plin. 13. 19.

³ Sc. 'and these oils are used in the raw state' (?). I do not see how Furlanus' explanation, quoted by Sch., is to be found in the text. The following sentence shews that T. does not claim to have settled the question.

καὶ ἀναδέχεται, διὸ καὶ τὴν συνέχειαν ἐξαιρεῖ· μανὸν δὲ γενόμενον [καὶ] τοῦ λίπους ἀφαιρεθέντος ἐν ῷ καὶ ἡ οἰκεία μάλιστα ὀσμή, δεκτικώτερον ἐγένετο τοῦ ἐπιβαλλομένου διὰ τὸ μὴ ἀντιστατεῖν.

- ΥΗ δὲ ἀπὸ τῶν ἀρωμάτων ὀδμὴ καὶ ἀσθενὴς ἄτε εἰς τὸ λιπαρὸν ἀνηλωμένη, καὶ ἔτι κατέχεται τούτφ διὰ τὸ πληρῶσαι τοὺς πόρους. ὅστε κατὰ λόγον κᾶν ἔλαττον ἢ τὸ ἐπιβαλλόμενον ἐπικρατεῖν τὴν τούτου ὀσμήν· εἰς ἀσθενέστατον γὰρ ἐμπίπτει καὶ δεκτικώτερον. ἀνὰ λόγον δ΄ ἔχει καὶ ἡ πολυχρονιότης ἡ ἐν ἑκάστφ καὶ ἡ πρὸς τὴν πύρωσιν εὐσθένεια καὶ τἄλλα τὰ τοιαῦτα. τὸ γὰρ δεκτικώτατον, οἰον τῆς βαλάνου, καὶ χρονιώτατον, καὶ διὰ τὴν αὐτὴν αἰτίαν· μάλιστα γὰρ ὥσπερ ἐν γίνεται καὶ συμφυὲς τὸ μάλιστα δεχόμενον· ἀεὶ γὰρ τὸ τοιοῦτον διαμονώτατον, διὸ καὶ πυρούμενον μάλιστα ἀπαθές.
- 20 'Ωσαύτως δὲ καὶ τῶν ἄλλων τὸ σησάμινον, τοῦτο γὰρ δεκτικώτατον· τὸ δὲ ἀμυγδάλινον παρακμάζει ταχὺ καὶ ὀλιγοχρονιώτατον διὰ τὴν ἐναντίαν αἰτίαν· τὸ γὰρ ἥκιστα δεξάμενον τάχιστα μεθίησι. τοῦ ῥοδίνου δὲ μάλιστα δεκτικὸν τὸ

1 I have bracketed sal.

²⁻² This passage is omitted, apparently by accident, in both W.'s texts, though represented in his Latin version. I 344

CONCERNING ODOURS, 18-20

themselves the viscid part of the oil, and so it attaches itself to them; thus the density of the oil is destroyed: the oil, thus becoming thinner by the removal 1 of its viscid part which chiefly contains the characteristic odour, becomes more receptive of the spice which is added to it, because it does not now offer resistance.

Again that odour which is due to the spices becomes less powerful as it is spent on the viscid part of the oil, while at the same time it is preserved by this because it has entirely filled up its passages. Wherefore it naturally follows that, even if the added spice is in small quantity, its odour predominates, since it passes into a vehicle which is in itself not at all powerful and which is more receptive than itself. A corresponding account may be given of the keeping quality of the several oils, of their power of resisting fire, and other such Thus that oil which is most receptive, for instance, that of the Egyptian balanos, will also keep longest, and for the same reason; namely that that oil which is most receptive unites, more than others, into one single substance, as it were, with the spices. Such a substance will always last longer than others; which also explains why, if exposed to fire, it is less affected than others.

Of the other oils the same applies to that of sesame, this being specially receptive; ² but, for the contrary reason, almond-oil soon loses its virtue and keeps for a shorter time than any other, for that oil which has been least receptive parts soonest with the property received. Sesame-oil however receives rose-perfume better than other oils ² because of its have printed it from Sch.'s text. The omission is evidently due to the double occurrence of τh of squares.

σησάμινον διὰ τὴν λιπαρότητα· πυρούμενον δὲ ἐξόζει σησάμου καθάπερ ἀναλυόμενον. αὶ μὲν οὖν τῶν ἐλαίων φύσεις καὶ δυνάμεις τοιαῦται.

- 21 V. Τὰ δ' ἀρώματα πάντα σχεδὸν καὶ εὕοσμα πλὴν τῶν ἀνθῶν ξηρὰ καὶ θερμὰ καὶ στυπτικὰ καὶ δηκτικά. τὰ δὲ καὶ ἔχοντά τινα πικρότητα, καθάπερ καὶ ἐν τοῦς πρότερον εἴπομεν, ὥσπερ ἔρις σμύρνα λιβανωτός, ὡς δ' ἀπλῶς εἰπεῖν καὶ τὰ μύρα. κοινόταται δὲ τῶν δυνάμεων τό τε στυπτικὸν καὶ τὸ θερμαντικόν, ἃ δὴ καὶ ἐργάζονται.
- 22 'Υποστύφονται μὲν οὖν πάντα πυρούμενα, τὰς δ' ὀσμὰς τὰς κυρίας ἔνια λαμβάνει ψυχρὰ καὶ ἀπύρωτα. καὶ ἔοικεν ὥσπερ τῶν ἀνθῶν τὰ μὲν ψυχροβαφῆ τὰ δὲ θερμοβαφῆ παραπλησίως ἔχειν καὶ ἐπὶ τῶν ὀσμῶν. πάντων δὲ ἡ ἔψησις εἴς τε τὴν ὑπόστυψιν καὶ τὰς κυρίας ὀσμὰς ἐνισταμένων τῶν ἀγγείων ὕδατι γίνεται καὶ οὐκ αὐτῷ τῷ πυρὶ χρωμένων. τοῦτο δέ, ὅτι μαλακὴν εἶναι δεῖ τὴν θερμότητα, καὶ ἀπουσία πολλὴ γένοιτ' ἀν τῆ φλογὶ χρωμένων, καὶ ἔτι καῦσιν ἀν ὄζοι.
- 23 Ποιεί δ' ἐλάττω τὴν ἀπουσίαν ὅσα πυρούμενα λαμβάνει τὰς κυρίας ὀσμὰς μᾶλλον ἢ ὅσα ψυχρὰ διὰ τὸ προφυρᾶσθαι τὰ πυρούμενα, τὰ μὲν οἴνῷ εὐώδει, τὰ δὲ ὕδατι· ἤττον γὰρ ἀναπίνει· τὰ δὲ ψυχρὰ ξηρὰ ὄντα μᾶλλον, καθάπερ ἴρις κοπεῖσα. 346

CONCERNING ODOURS, 20-23

viscid quality; and, when subjected to fire, it gives out a smell of sesame, as though it were being disintegrated. Such are the special characters and properties of the various oils.

Of the spices used in making perfumes and their treatment.

V. Almost all spices and sweet scents except flowers are dry hot astringent and mordant. Some also possess a certain bitterness, as we said above, as iris, myrrh, frankincense, and perfumes in general. However the most universal qualities are astringency and the production of heat; they actually produce these effects.

All spices are given their astringent quality by exposure to fire, but some of them assume their special odours even when cold and not exposed to fire; and it also appears that, just as with vegetable dyes some are applied hot and some cold, so is it with odours. But in all cases the cooking, whether to produce the astringent quality or to impart the proper odour, is done in vessels standing in water and not in actual contact with the fire; the reason being that the heating must be gentle, and there would be considerable waste if these were in actual contact with the flames; and further the perfume would smell of burning.

However there is less waste when the perfume obtains its proper odour by exposure to fire than when it does so in a cold state, since those perfumes which are subjected to fire are first steeped either in fragrant wine or in water: for then they absorb 1 less: while those which are treated in a cold state, being dry, absorb 1 more, for instance bruised iris-root. Thus, if

¹ αναπίνει. So Sch. explains. cf. ἐκπίνωσιν, 24.

λαμβάνοντος γὰρ τοῦ ἀμφορέως ξηρᾶς ἴριδος κεκομμένης μέδιμνον καὶ δύο ἡμίεκτα πολλὴν ποιεῖν φασὶν ἀπουσίαν, ἐὰν δὲ μετρίως φυράση λείπειν ὅσον δύο χόας, τοῖς δὲ πολλοῖς ἔλαττον.

24 Γίνεται δὲ τὸ βέλτιον ἴρινον ἐὰν ἢ ξηρὰ καὶ ἀπύρωτος ἡ ἴρις ἀκρατεστέρα γὰρ ἡ δύναμις ἡ ἐὰν φυραθεῖσα καὶ πυρουμένη. συμβαίνει δὲ ὅσπερ καὶ ἐκθλίβεσθαι μᾶλλον ἐκ τῶν προπεφυραμένων διὰ τὸ ἦττον ἀναδέχεσθαι καὶ ἔλκειν εἰς αὐτό προστύφοντες δὲ οὐ πολὺν χρόνον ἐῶσι τὰ ἀρώματα ἀλλ' ἐξαιροῦσιν, ὅπως μὴ πολὺ ἐκπίνωσιν.

25 Πρὸς ἔκαστον δὲ τῶν μύρων ἐμβάλλουσι τὰ πρόσφορα τῶν ἀρωμάτων, οἶον εἰς μὲν τὴν κύπρον καρδάμωμον ἀσπάλαθον ἀναφυράσαντες τῷ εὐώδει. εἰς δὲ τὸ ρόδινον σχοῖνον ἀσπάλαθον κάλαμον. ἡ δ' ἀναφύρασις ὁμοίως. καὶ τοῖς ἄλλοις ἀεὶ τὰ ἀρμόττοντα. τῷ ροδίνῷ δ' ἐμβάλλονται καὶ ἄλες πολλοὶ καὶ τοῦτ' ἴδιον παρὰ τάλλα, διὸ καὶ πλείστη ἀπουσία γίνεται μίγνυται γὰρ εἰς τὸν ἀμφορέα δύο μέδιμνοι.

26 Τῆς δὲ κύπρου ἡ μὲν ἐργασία παραπλησία τῆ

Το βέλτιον Γρινον W. after Sch.; το βέλτιον το Γρινον vulg. The article must be omitted in one place or the other.

 $\kappa \omega \rho \rho \rho \rho$, called from a tree of that name: not mentioned in H.P. of Plin. 12. 119.

⁴ cf. H.P. 9. 7. 2 and 3. ⁶ cf. H.P. 9. 7. 3.

 $^{^{\}rm 1}$ Dry measure: the equivalents given are, of course, only approximate.

CONCERNING ODOURS, 23-26

into eight and a half gallons of oil we put thirteen gallons of dry and bruised iris-root, they say that much loss is caused, while if one does not steep it too much, only about eleven pints and a half are wasted: and in the case of most perfumes the waste is less.

However the superior 2 iris-perfume is made by using the root dry and not subjecting it to fire: for then its virtue asserts itself more completely than when it is steeped in a liquid or subjected to fire. It also comes to pass that, if the perfumes have been first steeped, their virtues are, as it were, squeezed out of them to a greater extent, because they take in and absorb less: and so, when they are making them astringent, they do not leave the spices in the oil for long, but take them out, so that they should not absorb an excessive amount.

For making each perfume they put in the suitable spices. Thus to make kypros³ they put in cardamom⁴ and aspalathos,⁵ having first steeped them in sweet wine.⁶ To make rose-perfume they put in ginger-grass aspalathos and sweet-flag: and these are steeped as in the case of kypros. So too into each of the others are put the spices which suit them. Into rose-perfume moreover is put a quantity of salt ⁷: this treatment is peculiar to that perfume, and involves a great deal of waste, twenty-three gallons ⁸ of salt being put to eight gallons and a half of the perfume.

The manufacture of kypros resembles that of

 $^{^6}$ τ $\hat{\varphi}$ εὐάδει here evidently means the same as τ $\hat{\varphi}$ γλυκε \hat{i} , 44, where τ $\hat{\varphi}$ οἴν φ τ $\hat{\varphi}$ εὐάδει occurs just above: cf. μελικράτ φ $\hat{\eta}$ γλυκε \hat{i} , C. P. 6. 17. 2.

⁷ To prevent decay, as Diosc. 2. 53 explains.

⁸ Turn. suggests that μέδιμνοι should be μναι, the initial M having been misunderstood by a copyist.

τοῦ ἡοδίνου· πλὴν ἀλλ' ἐάν τις μὴ ταχέως ἐξαίρῃ καὶ ἀποθλίβῃ σῆψις ἐγγινομένη φθείρει τὰ μύρα διὰ τὴν δυσωδίαν· ποιεῖ γὰρ σῆψιν ἀνυγραινομένη. παραπλησία δ' ἐργασία καὶ τοῦ μηλίνου· προστυφέντος γὰρ ἐλαίου καὶ τὰ μῆλα ἐμβάλλουσιν εἰς ψυχρόν, εἰτ' ἐξαιροῦσι πάλιν πρὸ τοῦ μελαίνεσθαι κατὰ πάσας τὰς ἐμβολάς· μελαινομένων γὰρ σῆψις διὰ τὸ ἀνυγραίνεσθαι, καθάπερ καὶ ἐπὶ τῆς κύπρου.

27 VI. "Απαντα δὲ συντίθενται τὰ μύρα τὰ μὲν ἀπ' ἀνθῶν τὰ δὲ ἀπὸ φύλλων τὰ δὲ ἀπὸ κλωνὸς τὰ δ' ἀπὸ ῥίζης τὰ δ' ἀπὸ ξύλων τὰ δ' ἀπὸ καρποῦ τὰ δ' ἀπὸ δακρύων. μικτὰ δὲ πάνθ' ὡς εἰπεῖν. ἀπ' ἀνθῶν μὲν οἰον τὸ ῥόδινον καὶ τὸ λευκόϊνον. καὶ τὸ σούσινον καὶ γὰρ τοῦτο ἐκ τῶν κρίνων ἔτι δὲ τὸ σισύμβρινον καὶ τὸ ἐρπύλλινον καὶ ἢ κύπρος καὶ πρὸς τούτοις τὸ κρόκινον βέλτιστος δ' ἐν Αἰγίνη καὶ Κιλικία. ἀπὸ δὲ τῶν φύλλων οἰον τό τε μύρρινον καὶ τὸ οἰνάνθινον αὕτη δ' ἐν Κύπρφ φύεται ὀρεινὴ καὶ πολύοδμος ἀπὸ δὲ τῆς ἐν τῆ Ἑλλάδι οὐ γίνεται διὰ τὸ ἄοδμον.

28 'Απὸ ῥιζῶν δὲ τό τε ἴρινον καὶ τὸ νάρδινον καὶ τὸ ἀμαράκινον ἐκ τοῦ κόστου· τοῦτο γὰρ ὀνο-

² I have bracketed καl as suggested by Sch.

¹ cf. Diosc. 1. 58.

³ This passage, with some variations, is quoted by Athen. 15. 39. ⁴ cf. Plin. 13. 11.

⁵ cf. H.P. 6. 6. 11. for the plant, and for the perfume Athen, 15. 38.

CONCERNING ODOURS, 26-28

rose-perfume, except that, unless one soon takes out the flowers and squeezes them out, decay sets in and ruins the perfume by giving it a disagreeable smell, since they cause decay as they get soaked. Similar also is the manufacture of quince-perfume 1: the oil is first made astringent, and is cold when the quinces 2 are put into it: then they take them out before they turn black, removing each batch before the next is put in: for, as they turn black, decay ensues because they get soaked through—just as in the case of kypros.

Of the various parts of plants used for perfumes, and of the composition of various notable perfumes.

- VI. ³ Perfumes are compounded from various parts of the plant, flowers leaves twigs root wood fruit and gum: and in most cases the perfume is made from a mixture of several parts. Rose and gilliflower perfumes are made from the flowers: so also is the perfume called susinon,⁴ this too being made from flowers, namely, lilies: also the perfumes named from bergamot-mint and tufted thyme, kypros, and also the saffron-perfume; the crocus which produces this is best in Aegina and Cilicia. Instances of those made from the leaves are the perfumes called from myrtle and drop-wort⁵: this grows in Cyprus on the hills and is very fragrant: that which grows in Hellas yields no perfume, being scentless.
- ⁶ From roots are made the perfumes named from iris spikenard and sweet marjoram,⁷ an ingredient in which is koston; for it is the root to which this name

Digitized by Google

Instances of perfumes made from twigs seem to be missing.
7 cf. 30. Text perhaps defective.

μάζουσι τὴν ῥίζαν. τὸ δὲ χρίσμα τὸ Ἐρετρικὸν ἐκ τοῦ κυπείρου. κομίζεται δὲ ἀπὸ τῶν Κυκλά-δων τὸ κύπειρον. ἀπὸ ξύλου δὲ ὁ φοῖνιξ καλούμενος ἐμβάλλουσι γὰρ τὴν ὀνομαζομένην σπάθην ξηράναντες. ἀπὸ καρπῶν δὲ τό τε μήλινον καὶ τὸ μύρτινον καὶ τὸ δάφνινον τὸ δ' Αἰγύπτιον ἐκ πλειόνων, ἔκ τε τοῦ κιναμώμου καὶ ἐκ σμύρνης καὶ ἐξ ἄλλων.

"Ετι δ' ἐκ πλειόνων τούτου τὸ μεγαλείον καὶ γὰρ ἐκ κιναμώμου . . . καὶ ἐκ τῆς σμύρνης κοπτομένης ἔλαιον ῥεῖ· στακτὴ γὰρ καλεῖται διὰ τὸ <κατὰ> μικρὸν στάζειν. ὁ δὴ μόνον τινές φασιν ἀπλοῦν εἶναι καὶ ἀσύνθετον τῶν μύρων τὰ δ' ἄλλα πάντα σύνθετα, πλὴν τὰ μὲν ἐκ πλειόνων τὰ δ' ἐξ ἐλαττόνων, ἐξ ἐλαχίστων δὲ τὸ ἴρινον. οἱ μὲν οὖν οὕτω λέγουσιν, οἱ δὲ τὴν ἐργασίαν τῆς στακτῆς εἶναι τοιάνδε· τὴν σμύρναν ὅταν κόψωσι καὶ διατήξωσι ἐν ἐλαίω βαλανίνω πυρὶ μαλακῶ ὕδωρ ἐπιχεῖν θερμόν· συνιζάνειν δ' εἰς βυθὸν τὴν σμύρναν καὶ τοὔλαιον καθάπερ ἰλύν· ὅταν δὲ τοῦτο συμβῆ, τὸ μὲν ὕδωρ ἀπηθεῖν τὴν δ' ὑπόστασιν ἀποθλίβειν ὀργάνοις.

Τὸ δὲ μεγαλεῖον ἐκ ἡητίνης κεκαυμένης συντίθεσθαι καὶ ἐλαίου βαλανίνου μίγνυσθαι δὲ κασίαν κινάμωμον σμύρναν. πλείστην δὲ πραγματείαν περὶ τὸ μεγαλεῖον καὶ τὸ Αἰγύπτιον εἰναι,

¹ cf. H.P. 9. 7. 3; C.P. 6. 11. 13.

² cf. H.P. 2. 8, 4. σπάθην appears to be a conj. of W. for vulg. πλάτην: ἐλάτην Turn. cf. LS. s.v.

³ Said to be called after the inventor, one Megallos: cf. Plin. 13. 13.

CONCERNING ODOURS, 28-30

is applied. The Eretrian unguent is made from the root of kypeiron, which is obtained from the Cyclades as well as from Euboea. From wood is made what is called 'palm-perfume': for they put in what is called the 'spathe, having first dried it. From fruits are made the quince-perfume, the myrtle, and the bay. The 'Egyptian' is made from several ingredients, including cinnamon and

myrrh.

Again from several parts of the plant is made the perfume called megaleion,3 which is made from cinnamon and 4 and from the myrrh when it is bruised flows an oil: it is in fact called stakte 5 (in drops) because it comes in drops slowly. Some indeed say that this is the only simple uncompounded perfume, and that all the others are compound, though made from a larger or smaller number of ingredients, and that iris-perfume is made from the smallest number of all. assert this, but others declare that the manufacture of stakte (myrrh-oil) is as follows: having bruised the myrrh and dissolved it in oil of balanos over a gentle fire, they pour hot water on it: and the myrrh and oil sink to the bottom like a deposit; and, as soon as this has occurred, they strain off the water and squeeze the sediment in a press.

Megaleion, these authorities say, is compounded of burnt resin 6 and oil of balanos, with which are mixed cassia cinnamon and myrrh. They add that this perfume and the Egyptian are the most troublesome

⁵ cf. H.P. 9. 4. 10. 6 cf. Plin. 13. 7.

353

VOL. II. A

⁴ The end of the account of megaleion and the beginning of that of myrrh-perfume seem to be missing. ? Supply καὶ κασίας καὶ σμύρνης.

πλείστων γὰρ μίξιν καὶ πολυτελεστάτων. τῷ δὲ μεγαλείω καὶ τὸ ἔλαιον ἔψεσθαι δέχ' ἡμέρας καὶ δέκα νύκτας, εἶτα οὕτως τὴν ἡητίνην ἐμβάλλεσθαι καὶ τάλλα· δεκτικώτερον γὰρ ἀφεψηθέν. τὸ δ' ἀμαράκινον τὸ χρηστὸν ἐκ τῶν βελτίστων ἀρωμάτων συντίθεσθαι χωρὶς ἀμαράκου· τούτω δ' οὐ χρῆσθαι μόνω τῶν ἀρωμάτων τοὺς μυρεψοὺς οὐδ' εἰς ἐν μύρον, ἀλλὰ ψευδώνυμός τις ἡ ἐπίκλησις.

- ποιοῦσι δὲ καὶ τὰ μὲν ἀχρωμάτιστα τὰ δὲ κεχρωματισμένα. χρωματίζουσι δὲ ἀμαράκινον ρόδινον μεγαλεῖον, ἀχρωμάτιστα δὲ τῶν μὲν πολυτελῶν Αἰγύπτιον μήλινον κύπρος, τὰ δ' εὐτελῆ πάντα ταῦτα δὲ ἀχρωμάτιστα διότι τὸ μὲν Αἰγύπτιον καὶ τὴν κύπρον λευκὰ εἶναι βούλονται, τῷ δὲ μηλίνῳ τὴν τῶν μήλων χρόαν, τοῖς δ' εὐτελέσιν οὐ λυσιτελεῖ τὸ χρῶμα προστιθέναι. χρωματίζουσι δὲ τὰ μὲν ἐρυθρὰ τῷ ἀγχούσῃ, τὸ δ' ἀμαράκινον τῷ καλουμένῳ χρώματι τοῦτο δ' ἐστὶ ριζίον ὅ ἄγουσιν ἐκ τῆς Συρίας.
- 82 VII. Συνεργείν δὲ δοκοῦσι πρὸς τὰς γεύσεις οὐχ αἱ ὀδμαὶ μόνον ἀλλὰ καὶ αἱ δριμύτητες καὶ αἱ θερμότητες ἐνίων, διὸ καὶ τῶν οἴνων τισὶ τὰ τοιαῦτα μιγνύντες ὤσπερ κέντρον ἐμποιοῦσιν. ἔστι δὲ ἡ μὲν σμύρνη θερμὴ καὶ δηκτικὴ μετὰ

CONCERNING ODOURS, 30-32

to make, since no others involve the mixture of so many and such costly ingredients. To make megaleion, they say, the oil is boiled for ten days and nights, and not till then do they put in the resin and the other things, since the oil is more receptive when it has been thoroughly boiled. The superior kind of sweet marjoram-perfume, they say, is made of all the best spices except sweet marjoram: in fact this is the only spice which perfumers do not use for any

perfume, and the name is a misnomer.

Some perfumes are made up colourless, some are given a colour. They give a colour to sweet marjoram-perfume, rose-perfume, and megaleion, while among expensive kinds the Egyptian, quince-perfume and kypros are colourless, as well as all the cheaper kinds. The reason why these are made without colour is that it is desired that the Egyptian and kypros should look white and that quince-perfume should have the colour of quinces, while it is not worth while to add colour to the cheaper sorts. The dye used for colouring red perfumes is alkanet; the sweet marjoram-perfume is dyed with the substance called khroma (dye), which is a root imported from Syria.

Of the properties of various spices.

VII. It is thought that not only the smells of perfumes contribute to a pleasant taste, but also the qualities of pungency and heat which are found in some of them: accordingly some of these perfumes are also mixed with certain wines to give, as it were, 'point' to them. Thus myrrh is hot and has a biting quality as well as being astringent, and it also

¹ Clearly distinct from that mentioned in 28.

στύψεως, έχει δὲ καὶ πικρίαν. τὸ δὲ κινάμωμον δριμύτητά τινα μετρίαν μετὰ θερμότητος. παραπλησίως δὲ καὶ τὸ κόστον. ἡ δὲ κασία τούτων ὑπερβάλλει θερμότητι καὶ δριμύτητι καὶ στύψει. θερμὴ δὲ καὶ στυπτικὴ καὶ ἡ ἴρις, καθ' ὑπερβολὴν δὲ καὶ πικρὰ νέα οὖσα καὶ τὸν χρῶτα τῶν ἐργαζομένων ἀφελκοῖ. δηκτικὸν δὲ καὶ τὸ καρδάμωμον μετὰ θερμότητος. τοῦ δὲ βαλσάμου ὁ μὲν ἀπὸς καὶ τὸ καρπίον ἀνδρικώτερα πρὸς ἀμφότερα ταῦτα, τὸ δὲ ξύλον ἀσθενέστερον. παραπλησίαν δ' ἔχει τούτω δύναμιν καὶ τὸ ἄμωμον.

Ὁ δὲ σχοῖνος δηκτικώτερον μὲν τοῦ καλάμου καὶ θερμότερον, στυπτικὰ δὲ ὁμοίως ἄμφω. τούτων δὲ στυπτικώτερον τὸ κύπειρον. στύφει δὲ καὶ ἡ ἀσπάλαθος ἡ εὐώδης. ἡ δὲ νάρδος δηκτικὴ μετὰ θερμότητος. τὸ δὲ μάρον καὶ τὸ χρῶμα τὸ εἰς τὸ ἀμαράκινον ἐμμιγνύμενον θερμαντικά. [συνεργεῖ δὲ καὶ τῆς ἀγχούσης τὸ ρἰζίον εἰς τὴν χρόαν τοῦ ροδίνου καὶ τῆς ἴριδος.]

Νέα μὲν οὖν ὄντα τῶν ἀρωμάτων ἔνια δυνάμεις μὲν εὐθὺς ἔχει βαρείας καὶ δριμείας, παλαιούμενα δὲ μέχρι τῆς ἀκμῆς γλυκαίνεται, εἶτ ἀναλύεται πάλιν. οἰον ἡ ἰρις εἰς μὲν τὴν ἐργασίαν ἀκμάζει μετὰ τὴν συλλογὴν τρία ἔτη, καὶ διαμένει δὲ πλεῖστον χρόνον ἐξ ἔτη. τὸ δὲ μάρον ἔτη δύο. ἡ δὲ σμύρνα δέκα ἔτη διαμένει βελτίων γενομένη. παραπλησίως δὲ τούτοις ἡ τῆς ἀκμῆς διαμονὴ καὶ τοῦ κιναμώμου καὶ τοῦ κόστου καὶ τῆς κασίας. σχοῖνος δὲ καὶ κάλαμος παρακμάζει ταχύ. τῶν δ' ἀνθῶν τὰ μὲν εὐθὺς χλωρὰ ὄντα

cf. Plin. 21. 42.
 cf. Index, σχοῖνος (2).
 cf. Index, κάλαμος ὁ εὐώδης.

CONCERNING ODOURS, 32-34

has a bitter quality. Cinnamon again has a fair amount of pungency as well as heat. So too is it with koston. Cassia exceeds both of these in heat pungency and astringency. Iris-perfume is hot and astringent, and excessively bitter when it is fresh, in which state it causes sores on the skin of those that work on it. Cardamom has also a biting quality as well as heat. The juice and the fruit of balsam of Mecca are more active in producing both these qualities, while the wood is less so. Nepaul cardamom has also a property similar to this.

Ginger-grass² has a more biting quality than sweet-flag,³ and is hotter; but both are equally astringent. Kypeiron is however more astringent than either. The sweet-scented aspalathos also has this quality. Spikenard has a biting quality as well as heat. Maron and the khroma which is mixed with sweet marjoram-perfume are heating. ⁴The root of alkanet also contributes to the colour of rose-perfume

and iris-perfume.

Now some spices when they are fresh have at first heavy and pungent qualities, but in course of time become sweet till they have reached their prime, and then lose their properties again. Thus the iris is at its prime for manufacturing the perfume for three years after it was gathered, and 5 lasts for six years at longest. 5 Maron lasts two years; myrrh ten, and improves with time. Cinnamon koston and cassia keep at their best for about the same periods as these. Ginger-grass and sweet-flag soon get past their prime. Of flowers some, like the rose, possess

4 This sentence seems irrelevant here.

⁵⁻⁵ καl—έτη. These words are omitted, apparently by accident, in both W.'s editions, though represented in his Latin translation.

τὰς δυνάμεις ἔχει, καθάπερ τὸ ῥόδον, τὰ δὲ ξηρανθέντα, καθάπερ ὁ κρόκος καὶ ὁ μελίλωτος χλωρὰ γὰρ ὑγρότερα.

Τὰς μὲν οὖν φύσεις καὶ δυνάμεις τῶν ἀρωμάτων ἐκ τούτων θεωρητέον.

VIII. Δοκεί δὲ τὸ μεγαλείον ἀφλέγμαντον εἶναι παντὸς τραύματος τὸ δὲ ρόδινον ἄριστον πρὸς τὰ ὧτα. ταῦτα δ' οὐκ ἀλόγως. τοῦ μὲν γὰρ ἡ σύνθεσις ἐκ ρητίνης κεκαυμένης, ὥσπερ ἐλέχθη, καὶ κασίας καὶ κιναμώμου καὶ σμύρνης, ἄπαντα δὲ ταῦτα στυπτικὰ καὶ ξηραντικά. τὸ δὲ ρόδινον τοῖς ἀσὶν ἀγαθὸν ὅτι ἐν άλσὶν ἡ ποίησις ἀναξηραίνει γὰρ καὶ ἐκθερμαίνει διὰ τοὺς ἄλας διὸ καὶ ἡ ἀλοσάχνη ἀγαθόν. ἀλλὰ τὸ τῆς στραγγουρίας λόγου δεῖται καὶ γὰρ ταύτη λέγουσι μάλιστα βοηθεῖν. αἴτιον δ' ἄν εἴη διότι πῶν τὸ ὑπεξάγειν μέλλον ἀναλῦσαι δεῖ πρότερον τὸ ὑπεξαχθησόμενον τοῦτο δὲ οἱ ἄλες ποιοῦσιν, ἡ δ' εὐωδία τὴν ὁρμὴν ἀπέδωκε.

86 Διὰ τί δὲ τὸ ἴρινον εὕοσμον μὲν οὐ ποιεῖ δὲ τὴν όρμήν; ἢ διότι στυπτικὸν καὶ συνάγει τοὺς πόρους, ὥστε συγκλείσει κωλύειν τὴν δίοδον; ἀλλὰ καὶ κοιλίας λυτικὴ διά τε τὴν θερμότητα καὶ διὰ τὸ ἀποστύφειν τοὺς ἐπὶ τὴν κύστιν πόρους· ἀποκλειομένων γὰρ τούτων εἰς τὴν κοιλίαν ἡ συρροή. τὸ δὲ ὅλον φαρμακῶδες καὶ

^{1.} cf. C.P. 6. 14. 8 and 11.

CONCERNING ODOURS, 34-36

their virtues from the first while they are still fresh, some only after they are dried, as crocus and *melilotos*, these having a certain amount of moisture while they are fresh.

These examples may suffice for study of the characters and properties of spices.

Of the medicinal properties of certain perfumes.

VIII. Megaleion is believed to relieve the inflammation caused by any wound, and rose-perfume to be excellent for the ears. And this is probable enough. For the former is composed, as was said, of burnt resin cassia cinnamon and myrrh, and all these have astringent and drying properties: while the reason why rose-perfume is good for the ears is that salt is used in the manufacture of it: for it is by reason of the salt that it dries and warms (which is why 'sea-foam 2' is also good for the ears). Its use against strangury however needs explanation: for it is said to be specially helpful against this. The explanation may be that anything which is to remove the difficulty must first dissolve that which is to be removed; and this is the effect of the salt, while the fragrance supplies the necessary stimulus.

Why however, it may be asked, though irisperfume is fragrant, does it not give the stimulus? Perhaps it is because it is astringent and closes the passages, so that by shutting them it prevents free course. On the other hand this perfume acts as a laxative on the bowels because of its heating quality and because it astringes the passages leading to the bladder: for, when these are closed, the liquid collects in the bowels. In general iris-

² Said to be a zoophyte: cf. Arist. H.A. 9. 14. 2.

τὸ ἴρινον καὶ ἄλλα τῶν μύρων. ἡ δ' αἰτία πάντων ώς καθόλου εἰπεῖν ἐν ταῖς δυνάμεσι ταῖς εἰρημέναις, ὅτι στυπτικὰ καὶ θερμαντικά τὰ ἀρώματα γὰρ τὰ τοιαῦτα φαρμακώδη. ταῦτα μὲν οὖν ἔξω τῆς τέχνης.

- 1Χ. Κράσις δὲ καὶ μίξις οὐκ ἔστιν ὡρισμένη τῶν ἀρωμάτων, ὥστ' ἐκ τῶν αὐτῶν ἀεὶ χρηστὰ καὶ ὅμοια γίνεσθαι, ἀλλοῖα δὲ συμβαίνει διὰ τὴν ἀνωμαλίαν τῶν δυνάμεων τῶν ἐν τοῖς ἀρώμασι. τῆς δ' ἀνωμαλίας αἰτίαι πλείους. μία μέν, ἤπερ καὶ τοῖς ἄλλοις καρποῖς, ἡ τοῦ ἔτους κατάστασις αὕτη γὰρ πολυχουστέρας ότὲ δ' ἀσθενεστέρας τὰς δυνάμεις <ποιεῖ>. ἐτέρα δὲ ἐν τῆ συλλογῆ, τὸ προτερῆσαι τῆς ἀκμῆς ἡ ὑστερῆσαι· καὶ γὰρ τοῦτο οὐ μικρὸν διαφέρει. τρίτη δ' ἡ μετὰ τὴν συλλογήν, ὅσα χρόνου δεῖται πρὸς τὴν ἀκμήν, ὥσπερ ἐλέχθη· καὶ γὰρ ἐνταῦθά ἐστι τὸ προτερεῖν καὶ ὑστερεῖν.
- 38 Τούτων δὲ τὸ μὲν τῶν ἐτῶν οὐκ ἐφ' ἡμῖν, πλὴν εἰς τὸ εἰδέναι τὰ ποῖα σφοδροτέρας καὶ ἀσθενεστέρας ἔχει τὰς δυνάμεις τὰ δὲ κατὰ τὰς ἀκμὰς τῆς τε συλλογῆς καὶ μετὰ τὴν συλλογὴν ἐφ' ἡμῖν ἐστί, τῷ εἰδότι μᾶλλον τὸ ἐπιτυγχάνειν.

'Η μèν οὖν γένεσις καὶ σύνθεσις τῶν μύρων ἐκ τούτων.

CONCERNING ODOURS, 36-38

perfume, as well as others, has medicinal properties. And the explanation in all cases, to put it generally, lies in the above-mentioned properties of astringency and heating; for it is spices possessing these properties that are medicinal. However these matters lie outside our subject of study.

Of rules for the mixture of spices, and of the storing of various perfumes.

IX. There is no fixed rule for the combination and mixture of spices in the sense that the same components will always produce a satisfactory and a uniform result: the result varies by reason of the varying quality of the virtues found in the spices. For this there are several reasons. One, which applies also to fruits, is the character of the season; this causes the virtues to be sometimes much more than ordinarily powerful, sometimes less so. Another is to be found in the time of collection, according as it is made before or after the spices are in their prime. A third cause operates after the collection, that is, in the case of those spices which need time to come to their prime, as was said 1: for here too it is possible to be too soon or too late.

Of these causes that which depends on the seasons is not within our control, except so far as we can discover which spices in a particular season have powerful, and which weak virtues.² But we can control those which depend on collecting them when in their prime, or on keeping them after they are collected, that is, if we know pretty well how to

hit the right moment.

So much for the origin and composition of perfumes.

¹ 34. ² i.e. and we can select accordingly.

Πολυχρονιώτατον δ' έστὶ τό τ' Αἰγύπτιον καὶ τὸ ἔρινον καὶ τὸ ἀμαράκινον καὶ τὸ νάρδινον. πάντων δε μάλιστα ή στακτή, διαμένει γάρ όποσονοῦν χρόνον. μυροπώλης δέ τις έφη παρ' αὐτῷ μεμενηκέναι Αἰγύπτιον μεν όκτὼ έτη, Ιρινον δὲ εἴκοσι, καὶ ἔτι διαμένειν βέλτιον ον τῶν ἀκμαζόντων. ή μεν οθν γρονιότης εν τούτοις.

Τὰ δ' ἄνθινα πάντα ἀσθενη, συμβαίνει δὲ τοις ανθίνοις ακμάζειν μεν ώς έπι το παν μετά δίμηνον, μεταβάλλειν δ' έπὶ τὸ χείρον ένιαυτοῦ προελθόντος καὶ περικαταλαβούσης τῆς ώρας έν ή την άκμην λαμβάνει το άνθος. άνὰ λόγον δὲ τῆ ἀσθενεία και τὸ εὐπέπαντα είναι και ὅλως εὐδιἄπνευστα· τὰ δ' ἐκ τῶν ῥιζῶν καὶ τῶν λοιπῶν γρονιώτερα πλείων γάρ ή όσμη και ισχυροτέρα

καὶ σωματωδεστέρα.

40 Διαφθείρει δὲ τὰ μύρα καὶ ὥρα θερμὴ καὶ τόπος καὶ ὁ ήλιος, αν τεθώσι διὸ καὶ οἱ μυροπώλαι ζητούσι τὰς οἰκίας ὑπερώους καὶ μὴ προσηλίους άλλ' ὅτι μάλιστα παλισκίους ἀφαιρείται γὰρ τὰς ὀσμὰς ὁ ἥλιος καὶ τὸ θερμὸν καὶ όλως έξίστησι της φύσεως μαλλον του ψυγρού. τὸ δὲ ψυχρὸν καὶ ὁ πάγος, εἰ καὶ ἀοσμότερον ποιεί διὰ τὸ συστέλλειν, ἀλλ' οὐκ ἀφαιρείταί γε την δύναμιν τελέως. ή πονηρά γάρ φθορά, καθάπερ τῶν οἴνων καὶ τῶν ἄλλων χυλῶν, τῷ 41 τὸ οἰκεῖον ἀφαιρεῖσθαι θερμόν. διὸ καὶ εἰς ἀγγεῖα μολυβδα εγγέουσι και τους αλαβάστρους ζητοῦσι τοιούτου λίθου. ψυχρον γάρ και πυκνον και ό

CONCERNING ODOURS, 38-41

Those which last longest are the Egyptian, the iris, the sweet marjoram and the spikenard perfumes: but myrrh-oil has the longest life of any; for it will keep any time. A certain perfumer said that he had had Egyptian perfume in his shop for eight years, and iris-perfume for twenty, and that it was still in good case, in fact better than fresh perfume. These are instances of perfumes which will keep a long time.

On the other hand all those made from flowers have little vigour. These are usually at their best after two months, but they deteriorate when a year has past and the season has come round again at which the flowers are at their best. Also, as these perfumes lack vigour, so also do they quickly mellow, and, in most cases, quickly evaporate. Those made from roots and the other parts of the plant last longer, their odour being fuller stronger and more substantial.

Perfumes are ruined by a hot season or place or by being put in the sun. This is why perfumers seek upper rooms which do not face the sun but are shaded as much as possible. For the sun or a hot place deprives the perfumes of their odour, and in general makes them lose their character more than cold treatment: while cold and frost, even if they make them less odorous by congealing them, yet do not altogether deprive them of their virtue. For the most destructive thing that can happen to them, as to wines and other savours, is that they should be deprived of their proper heat. This is why men put them into vessels of lead and try to secure phials of alabaster—a stone which has the required effect: for lead is cold and of close texture, and stone has

μόλυβδος καὶ ὁ λίθος ὁ τοιοῦτος καὶ ἄριστος τοῖς μύροις ὁ μάλιστα τοιοῦτος. ὥστε δι' ἄμφω τηροῦσι, καὶ τῷ ψυχρῷ καὶ τῷ πυκνῷ, μήτε διιέντες ἔξω τὴν ὀσμὴν μήθ' ὅλως ἐπιδεχόμενοι μηδέν. καὶ γὰρ ἡ ἀναπνοὴ φθείρει καὶ τὸ ἔξωθεν ἐπεισιὸν καὶ ἀλλότριον ἐπεὶ καὶ τὰ πνεύματα φθείρει καὶ καταναλίσκει, καθάπερ ἐλέχθη, τὰς ὀσμάς, ἄλλως τε καὶ τὰς μὴ φυσικάς.

42 Χ. Κεφαλαλγή δὲ τῶν μὲν πολυτελῶν τὸ ἀμαράκινον καὶ τὸ νάρδινον καὶ μεγαλεῖον, τῶν δ' εὐτελῶν ὅλως μὲν τὰ πλεῖστα μάλιστα δὲ τὸ δάφνινον. ἐλαφρότατα δὲ τὸ ῥόδινον καὶ ἡ κύπρος, ἃ καὶ τοῖς ἀνδράσι μάλιστα ἁρμόττειν δοκεῖ, καὶ πρὸς τούτοις τὸ κρίνον ταῖς δὲ γυναιξὶν ἡ στακτὴ καὶ τὸ μεγαλεῖον καὶ τὸ Αἰγύπτιον καὶ τὸ ἀμαράκινον καὶ τὸ νάρδινον διὰ γὰρ τὴν ἰσχὺν καὶ τὸ πάχος οὐκ εὐαπόπνοα οὐδ' εὐαφαίρετα ζητοῦσι <γὰρ> τὰ χρόνια.

 $^{^{1}}$ e.g. alabaster, which here at least is spoken of as a kind of stone. 2 $\gamma \lambda \rho$ ins. Sch.

CONCERNING ODOURS, 41-43

the same character, that being the best for keeping perfumes which has it in the highest degree.¹ So that vessels made of these materials keep the perfume well for both reasons, their coolness and their closeness of texture: they neither let the odour pass away through them, nor do they take in anything else. For evaporation destroys the perfume, and so also does any foreign substance which finds its way in: for even draughts of air destroy odours and cause them to waste, as was said, especially those odours which do not belong to a thing's essential nature.

Of the properties of certain perfumes.

X. Headache is caused by sweet marjoram spikenard and megaleion among costly perfumes: most of the cheap ones have also this effect, notably that made from bay. The lightest are rose-perfume and kypros, which seem to be the best suited to men, as also is lily-perfume. The best for women are myrrh-oil, megaleion, the Egyptian, sweet marjoram, and spikenard: for these owing to their strength and substantial character do not easily evaporate and are not easily made to disperse, and ² a lasting perfume is what women require.

Inasmuch however as some perfumes are stronger than others, the stronger being those made from roots and the others already mentioned, for this reason those derived from flowers are more fragrant if they are not bruised, while bruising improves those made from roots and the others. For the former kind evaporate and pass off as they are warmed by the bruising, thus losing their character, while the latter owing to their strength have, as it

ὥσπερ ἀνοιγομένων τινῶν πόρων ἐκ τῆς τρίψεως 44 ἐμφανεστέραν ποιεῖ τὴν ὀσμήν. ὁ καὶ ἐπ' αὐτῶν τῶν ἡιζῶν καὶ ὅλως τῶν στερεῶν συμβαίνει, καθάπερ ἐλέχθη. κατὰ δὲ τῶν ἀνθῶν ἐναντίως, ὥστε ἡκολούθηκεν ἑκάτερα τῆ ἀρχῆ. τὰ δ' ἐκ τῆς σμύρνης εὐλογώτατα δι' ἄμφω· καὶ γὰρ μίγνυται μᾶλλον καὶ <ἡ> θερμότης ἡ τῆς τρίψεως οὐκ ἀλλοτρία, μαλακή τις οὖσα· καὶ γὰρ <ἡ> σμύρνα ζητεῖ τινα πύρωσιν. ἀπλῶς δὲ πᾶν τὸ πολύοδμον ἄντ' εὐῶδες ἄντε κακῶδες ἄντε δριμὺ ἄντ' ὀξὸ ἄντ' ὁποιονοῦν τυγχάνη, κινούμενον ἐμφανέστερον· τότε γὰρ ὥσπερ ἐνεργεία ἀναμίγνυται μᾶλλον τῷ ἀέρι.

Των δε μύρων το Αιγύπτιον και ή στακτή και εί τι άλλο πολύοδμον [και] μιγνύμενα τῷ οἴνω τῷ εὐωδει ἡδίω· παραιρεῖται γὰρ ἡ βαρύτης αὐτῶν ἐπεὶ καὶ ἡ σμύρνη αὐτὴ πρὸς τὴν ἀναθυμίασιν βρεχθεῖσα ἐν τῷ γλυκεῖ, καθάπερ ἐν τοῖς πρότερον

έλέχθη.

Προς δε τὰς δυνάμεις σκοπουμένοις δόξειεν ἂν ἄτοπον εἶναι τὸ συμβαῖνον ἐπὶ τοῦ ροδίνου κουφότατον γὰρ δν καὶ ἀσθενέστατον ἀφανίζει τὰς τῶν ἄλλων ὀσμὰς ὅταν προμυρισθῶσι. διὸ καὶ οἱ μυροπῶλαι τοὺς ἐπιδιστάζοντας καὶ μὴ ἀνουμένους παρ' αὐτῶν ἐπιμυρίζουσι τούτῷ πρὸς τὸ μὴ αἰσθάνεσθαι τὰ παρὰ τῶν ἄλλων. αἴτιον δ' ὅτι λεπτότατον ὃν καὶ προσφιλὲς τῆ αἰσθήσει διὰ τὴν κουφότητα μάλιστα διικνεῖται καὶ συμ-

¹ ἡ ins. W. ² ἡ ins. W.

³ The words ἄντε δριμὸ are omitted in both W.'s editions, but represented in his Latin version.

CONCERNING ODOURS, 43-45

were, certain passages opened by the bruising, and so their fragrance is made more obvious. This, as was said, also takes place in the case of the roots themselves and of the solid things in general; but the result in the case of flowers is just the opposite, so that both kinds behave according to their origin. That this should apply to the perfumes made from myrrh is quite natural for both reasons; they mingle more than others with the air, and the heat 1 due to the bruising is not prejudicial, since it is gentle, and myrrh 2 in fact requires a certain amount of heating. And in general any strong odour, whether it be pleasing or the reverse, whether it be pungent³ or sharp, or whatever its character, becomes more pronounced with movement; for then it becomes, as it were, active and mingles more with the air.

The Egyptian perfume, myrrh-oil, and any others that have a strong odour become sweeter if they are mixed with fragrant wine; for then their heavy quality is removed. In fact myrrh itself is made to exhale a more fragrant odour by being steeped in

sweet wine, as was said 5 in a former treatise.

If one has regard to the virtues of the perfumes in question, one may well be surprised at what happens in the case of rose-perfume:—though it is lighter and less powerful than any other, if one has first been scented with it, it destroys the odour of the others. And this is why perfumers, if a purchaser hesitates and is not inclined to buy this perfume, scent him with it so that he is not able to smell the others. The explanation is that, being very delicate and acceptable to the sense of smell, by reason of its lightness it penetrates as no

⁴ I have bracketed καλ. ⁵ C.P. 6. 17. 2.

πληροί τοὺς πόρους, ὥσθ' ἡ αἴσθησις κατειλημ
46 μένη καὶ πλήρης οὖσα κρίνειν ἀδυνατεί. δύο
γάρ εἰσι τρόποι, τάχα δὲ τρείς, οἱ κωλύοντες τὴν
κρίσιν. εἰς μὲν ὁ νῦν εἰρημένος. ἄλλος δ' ὁ ἀπὸ
τῶν ἰσχυρῶν ὥσπερ μεθύσκων τὴν αἴσθησιν καὶ
καρηβαρᾶν ποιῶν. τρίτος δ' ὅταν προκαταληφθῆ
τῷ βελτίονι. τὸ γὰρ ἐπεισάγειν τὸ χείρον οὐ
ῥάδιον. οὐ δέχεται γὰρ ἡ αἴσθησις, ὥσπερ οὐδ'
ἐπὶ τῶν χυλῶν καὶ ὅλως τῶν κατὰ τὴν τροφήν.

47 Κατισχυαίνειν δὲ δοκεῖ τὸ ῥόδον καὶ τὴν σύνθετον ὀσμήν· ὅταν γὰρ ἀκμιίζη τὸ ἄνθος, ῥοδίζουσι τὰς συνθέσεις, ἀνοιγόμεναι δ᾽ ἐξόζουσι τούτου μόνου καὶ μάλιστα. παύεται δὲ ταχὰ καὶ λήγει διὰ τὴν ἀσθένειαν καὶ λεπτότητα, δι᾽ ἢν καὶ ἐξόζει τῶν ἄλλων· λεπτὴ γὰρ οὖσα ἡ ἀναπνοὴ καὶ ἤθροισμένη τῆ κατακλείσει προτερεῖ τε τῶν λοιπῶν καὶ διαδίδοται πανταχοῦ. διὰ ταὐτὸ δὲ τοῦτο καὶ ἀπολήγει ταχὰ καὶ κατακρατεῖται πάλιν· ἀσθενεῖ γὰρ τὸ λεπτὸν καὶ μαλακόν.

48 Ποιοῦσι δέ τινες τοῦτο καὶ τῶν οἴνων, ὥστε προποθέντες ἀφανίζειν τὴν τῶν ἄλλων ἡδονήν. ἔνιοι δ' ὥστε μὴ ἐπιδέχεσθαι ῥαδίως τοὺς ἄλλους,

¹ cf. 57, 58. σύνθετος δσμή or σύνθεσις seems to mean a kind of pot-pourri, which was from time to time renewed with fresh rose-petals. Sch. understands σύνθεσις to mean 'clothes in a wardrobe' (cf. Lat. synthesis), but it must surely have the same meaning here as σύνθετος δσμή: Sch.'s citation from 57 does not seem to prove his point, and μένουσι πολὺν χρόνον αί συνθέσεις in 58 is conclusive against him. cf. also 69.

CONCERNING ODOURS, 45-48

other can and fills up the passages of the sense, so that being entirely taken up and filled with it, it is unable to judge of others. For the power of judging is inhibited in two, or possibly in three ways; one is that which has just been mentioned; another is that the sense of smell is, as it were, intoxicated with its powerful virtues and becomes stupefied: the third is that the sense may be preoccupied with the superior odour; for then it is not easy to introduce after it what is inferior, since the sense of smell refuses this—just as the sense of taste in like case refuses flavours and foods in general.

It is also thought that the rose even weakens the effect of compound perfume 1; for, when the flower is at its best, they treat compound perfumes with it; and, when these come to be opened, they smell only or chiefly of rose. However this effect is only temporary and transient because of the weakness and delicacy of the rose-scent, (the very quality which also causes it to assert 2 itself over the scent of the other ingredients). For, as it is so delicate and is compressed by confinement, it is exhaled before the others and disperses in all directions. It is also for this reason that the rose-scent only asserts itself for a short time and then is overpowered again; for anything that is delicate and subtle must be lacking in vigour.

Certain wines have also a similar effect: if they are first drunk, there is no satisfaction in others. Some again make it even difficult to take others

369

VOL. II. B B

 $^{^2}$ i.e. when the pot-pourri is first opened: the 'delicacy' of the rose-scent causes it to be given off quickly and so (1) to be the first scent perceived, (2) to be volatile. $^{\ell\xi}\delta\zeta\omega$ in this passage is used with gen. in two distinct senses.

ὥσπερ ὁ Ἐρυθραῖας άλυκός τις ὧν καὶ μαλακός. τὴν αἰτίαν <δὲ> πειρατέον ἐκ τῶν ὁμοίων λαμβάνειν· ἔχει δὲ τοῦτ' ἴδιον τὸ ῥόδινον, ὥσπερ σχεδὸν καὶ μικρῷ πρότερον εἴρηται· τὰ μὲν γὰρ ἄλλα πάντ' ἡ τὰ πλεῖστα κεφαλαλγῆ, τοῦτο δέ, ὥσπερ ἐλέχθη, λυτικὸν καὶ βάρους καὶ ἀλγηδόνος καὶ τῆς ἀπὸ τῶν μύρων.

Ή δ' αἰτία φανερὰ διὰ τῶν προειρημένων, εἴπερ ἐπικρατεῖ καὶ διαδύεται πανταχοῦ. τὰ μὲν γὰρ ἄλλ' ὅσα κεφαλαλγῆ βαρέα διὰ τὸ ἐκ τοιούτων συγκεῖσθαι τὰ μὲν ρίζῶν τὰ δ' ὀπῶν· τοῦτο δὲ καὶ τῆ ὀσμῆ ἐλαφρὸν καὶ τῆ θερμότητι σύμμετρον εἰς τὸ συμπέψαι καὶ διανοῖξαι τοὺς πόρους. οἱ γὰρ δὴ πόνοι τῆς κεφαλῆς ἡ καθυγραινομένης ἡ πνευματουμένης τῷ ἐναπολαμβάνεσθαι, ὅστε τὸ μὲν ἐκκρῦναι δεῖ τὸ δὲ πέψαι ἡ ἀφελεῦν.

Πρὸς ἄπαντα δὲ ἡ θερμότης χρήσιμον, καὶ εἰς ἀφαίρεσιν καὶ ἔτι μᾶλλον εἰς τὸ πέττειν καὶ διανοίγειν τοὺς πόρους, εἰς ἃ συμβάλλεται τὸ ἐν τῷ ἀλὶ πεποιῆσθαι· καὶ γὰρ ἀναστομοῦσι καὶ διαθερμαίνουσιν οἱ ἄλες. ἡ δ' εὐοσμία καὶ ὁρμήν τινα ποιεῖ πρὸς τὴν κίνησιν. ἀγαθὸν δὲ καὶ δοκεῖ πρὸς τοὺς κόπους εἰναι τῆ θερμότητι σύμμετρον ὂν καὶ τῆ κουφότητι καὶ τῆ διαδύσει πρὸς τοὺς ἐντὸς πόρους· ὡς δὲ τινές φασιν, οὐχ ἦττον ἡ κύπρος ἔτι τούτου· μαλακὴ γὰρ ἡ ὀσμὴ καὶ

¹ cf. 52. ² δè ins. W.

³ i.e. the case is so far analogous to that of rose-perfume; but the comparison does not hold as to what follows.

CONCERNING ODOURS, 48-50

after them; this is the effect for instance of wine of Erythrae,1 which has a taste of brine and is subtle. The 2 explanation one must endeavour to find by comparing analogous cases.3 However there is one peculiarity which as we have already more or less indicated, is possessed by rose-perfume only; while all or most of the others are heady, this, as was said, gives actual relief from heaviness and discomfort. even from that caused by other perfumes.

The reason for this is plain in view of what has been already said, seeing that this perfume overpowers others and penetrates everywhere. For the others that are heady are heavy because they are made of heavy substances, whether roots or juices; while this perfume is both light as to its scent and also by its heat well adapted to bring the passages to a suitable temperature and to open them. pains in the head are due to an excess of moisture in it, or of air which gets confined in it, so that it is necessary to get rid of the one, and to raise the temperature of the other or to remove it.

And for all such purposes heat is useful, both for removing the moisture or air, and, still more, for raising the temperature of the passages and opening them: and to these ends it is helpful that the perfume should have been prepared with salt, since the effect of salt is to open the passages and to warm them thoroughly. Again the fragrance also supplies a stimulus to movement. This perfume is also considered to be good against lassitude, because its heat and its lightness make it suitable, and also because it penetrates to the inner passages. Some however say that kypros is quite as efficacious: for this too has a delicate scent which is grateful to the

προσφιλής τῷ χρωτὶ καὶ ή ταύτης. καὶ ταῦτα μέν καὶ τὰ ὅμοια τούτοις ὥσπερ ἴδια αν είη.

- ΧΙ. Τοῦ ὁοδίνου δὲ αἰ μίξεις καὶ ἐν ταῖς ὀσμαῖς καὶ ἐν τοῖς χυμοῖς, ἐὰν ἡρμοσμέναι τυγχάνωσιν, έγουσί τινα γρείαν, αι μεν άφαιρουσαι την βαρύτητα καὶ τὴν ἰσχύν, αἱ δ' εὐοδμίαν τινὰ αἱ δὲ γλυκύτητα έμποιούσαι, καθάπερ καὶ έπὶ τῶν οίνων, και γαο ο έν Θάσω ο έν τω πρυτανείω διδόμενος, θαυμαστός τις ώς ξοικε την ήδονήν. ηρτυμένος έστίν εμβάλλουσι γάρ είς το κεράμιον σταίς μέλιτι φυράσαντες, ώστε την μέν οσμην απ' αυτού, την δε γλυκύτητα από τού σταιτός λαμβάνειν τὸν οἶνον.
- 52 Συμβαίνειν δὲ τοῦτο καὶ κατὰ τὰς τῶν οἴνων μίξεις οίον εάν τις κεράση σκληρον και εὔοσμον μαλακώ καὶ ἀόσμω, καθάπερ τὸν Ἡρακλεώτην καὶ τὸν Ἐρυθραῖον, τοῦ μὲν τὴν μαλακότητα τοῦ δὲ τὴν εὐοσμίαν παρεχομένου συμπίπτει γὰρ άμα τὰ κακὰ ἀλλήλων ἀφανίζειν τῆ μαλακότητι θατέρου <καὶ τη εὐοσμία θατέρου>. πολλάς δὲ καὶ ἄλλας οἱ ἔμπειροι λέγουσι καὶ ἴσασι μίξεις. δ καὶ ἐπὶ τῶν ὀσμῶν εὕλογον συμβαίνειν, καὶ ἐπὶ τῶν γρωμάτων ἄν τις λαμβάνη τὰς ἁρμοττούσας μίξεις. τοῦτο μεν οὖν ἴδιον τοῦ ῥοδίνου.

53 Τὸ δὲ κοινὸν ἐπὶ πάντων ἀπόρημα, τί δή ποτε

Quoted by Athen. 1. 58.
 Cf. 48.
 This sentence must be defective: as it stands, the effect of only one wine is given, though the effect is said to be

CONCERNING ODOURS, 50-53

skin. These and similar properties may be considered peculiar to these particular perfumes.

Of other properties and peculiarities of perfumes.

XI. The admixture of rose-perfume, whether in scents or in flavours, if it be well blended, is beneficial, in the one case by removing the heaviness and strength of the scent, in the other by imparting a fragrant scent or a sweet taste to the flavour, as in the case of wines. ¹Thus the wine which is served in the town-hall of Thasos, which appears to be of wonderfully delightful quality, is thus flavoured. For they put into the jar a lump of dough which has been kneaded up with honey, so that the wine gets its fragrance from itself, but its sweet taste from the honeyed dough.

This result also follows, it is said, from the mixture of different wines,—for example, if a strong fragrant wine be mixed with one that is mild and without fragrance, (for instance, if wine of Heraclea be mixed with wine of Erythrae), since the latter contributes its mildness and the former its fragrance: for the effect is that they simultaneously destroy one another's inferior qualities through the mildness of the one and the fragrance of the other. There are many other such blends mentioned by and known to experts. And it is quite to be expected that such a result should follow from blending odours, as it does from blending colours, if one discovers the suitable combinations. This then is peculiar to rose-perfume.

However there is one question which applies to all perfumes, namely, why it is that they appear to mutual. I have added καὶ τῆ εὐοσμία θατέρου after Sch.; his text however is συμπίπτει γὰρ ἄμα, καὶ τὰ κακὰ ἀλλήλων

άφανίζει, τη μαλακότητι < καὶ τη εὐοσμία> θατέρου.

ἀπὸ τοῦ καρποῦ τῆς χειρὸς ἥδιστα φαίνεται, διὸ καὶ οἱ μυροπῶλαι τοῦτο μυρίζουσι τὸ μέρος. τὴν δ' αἰτίαν ἐκ τοῦ ἐναντίου ληπτέον, ὅτι τὸ θερμὸν ἐξίστησι καὶ ἀλλοιοῦ ταχεῖα δ' ἤδη ἡ αἴσθησις τοῖς μύροις ἀναμιγνυμένοις τῷ χρωτί.

Α 'Απορείται δὲ δίοτι οἱ μὴ εἰωθότες μυρίζεσθαι μᾶλλον ἐξόζουσι τῶν συνεχῶς μυριζομένων· εἴη μὲν γὰρ ἄν λέγειν καὶ ὅτι φαντασίαι καὶ οὐκ ἀλήθειαι διὰ τὸ μὴ εἰωθός· εἰ δ' οὖν καὶ ἀληθές, ἔοικε τὸ μὲν οἷον συναναμίγνυσθαι πλείοσιν ὀσμαῖς ἐτέραις ὑφ' ὧν ἀμαυροῦται, συγκαταμιγνυμένου καὶ τοῦ χρωτός, τὸ δὲ ὥσπερ ἀκέραιον δέχεσθαι τὸ μανὸν καὶ ἐκφαίνειν τῆ αἰσθήσει χρονιζόμενον. εἴη δ' ἄν καὶ ἐναντίως λαβεῖν ὡς ἤττον δεχομένου διὰ τὸ ἀσύνηθες, βραδύτερον δ' ἀναμιγνύμενα πλείω χρόνον ἐξόζειν. καὶ τοῦτο μὲν ἔλαττον καὶ οὐ φανερῶς ὁμολογούμενον.

55 "Απτεται δε μάλιστα τοῦ χρωτὸς καὶ κεφαλῆς καὶ τῶν ἄλλων καὶ πλεῖστον χρόνον ἐμμένει τὰ

¹ Sc. a part of the body which, not being fleshy, does not spoil the scent by its warmth. So Plin. 13. 19. appears to give the point—experimentum (unguentorum) inversa manu capitur, ne carnosae partis calor vitiet, though it may be questioned whether inversa manu represents καρποῦ. Pliny's

CONCERNING ODOURS, 53-55

be sweetest when the scent comes from the wrist¹; so that perfumers ² apply the scent to this part. The explanation must be sought by observing what happens in the contrary case, inasmuch as heat changes or destroys the character of a scent, and the effect on the sense of smell is immediately perceived when perfumes are brought into close contact with the skin.

The question is also raised why those who do not habitually use perfumes smell of them more strongly. when they do so, than those who use them habitually. The suggestion might be made that this is an illusion due to the fact that the use is not habitual, and does not represent what really happens. If however it does, it would appear that in the one case the perfume becomes, as it were, confused with a number of other scents which weaken its force (the smell of the skin also becoming mixed with it), while in the other case the porous condition of the skin takes in the scent as it were uncontaminated, and so makes it perceptible by the sense of smell, because it lingers for some time. One might also make a suggestion of opposite character, that the skin takes in perfumes less readily because it is not used to them, and so, as the perfumes mingle with it more slowly, they preserve 3 their scent for a longer time. One may add that this is a small point and that all do not agree as to the fact.

Those perfumes whose scent is strongest get the best hold on the skin head and other parts of the body, and last for the longest time: such are phrase presumably means the back of the hand, and suggests

that ἀπὸ τοῦ καρποῦ may be corrupt.

2 Sc. in offering samples for choice.

³ Sc. it is not absorbed by the skin.

ἰσχυρότατα ταῖς ὀσμαῖς, οἶον μεγαλεῖον, Αἰγύπτιον, ἀμαράκινου· τὰ δ' ἀσθενῆ καὶ <οὐ> πολύοδμα, κούφην ἔχουτα τὴν ἀναπνοήν, ταχεῖαν ποιεῖ καὶ τὴν ἀπόλειψιν, ὥσπερ τό τε ῥόδινον καὶ ἡ κύπρος.

Ένια δὲ καὶ εἰς τὴν ὑστεραίαν οὐ χεῖρον ὄζει, διαπεπνευκυίας εἴ τις ἐνῆν βαρύτης. τὰ δὲ καὶ ὅλως ἔμμονα μᾶλλον, ὥσπερ ἡ νάρδος καὶ τὸ ἴρινον, πάντων δὲ μάλιστα <τὰ> ἰσχυρότατα. καὶ τὰ μὲν ἔν τε τοῖς λουτροῖς καὶ τῆ ἀνέσει διατηρεῖ πως τὴν ὀσμὴν ἡ οὐ συγκακύνει· τὰ δὲ κακυνόμενα πλείω ποιεῖ δυσωδίαν αὐτῶν τῶν ἰδρώτων, ὡς ἃν σήψεώς τινος ἡ διαφθορᾶς γινομένης.

Καὶ τὰ μὲν περὶ τῆς τῶν μύρων ποιήσεώς τε καὶ δυνάμεως ἐπὶ τοσοῦτον εἰρήσθω.

57 ΧΙΙ. Τὰ δὲ περὶ τὴν τῶν ξηρῶν μίξιν, ἐξ ὧν <τὰ> διαπάσματα καὶ αἱ συνθέσεις, οὐκ ἔτι ζητεῦ μίξιν τῶνδέ τινων ὡρισμένων, ἀλλ' ὅσφ ἄν τις πλείω καὶ ποικιλώτερα μίξη, τοσούτω καὶ ἡ ὀσμὴ λαμπροτέρα καὶ ἡδίων, ὥσπερ καὶ ἐξ αὐτῶν τῶν ἀρωμάτων τῶν προχείρων εἰς ταὐτὸ γὰρ μιγνύντες ἄπαντα χρῶνται. ζητοῦσι δ' ἐν τούτοις καὶ σπεύδουσιν ὥστε μὴ ἑνὸς ἀλλὰ πάντων

¹ τà ins. Sch.

CONCERNING ODOURS, 55-57

megaleion, Egyptian perfume and sweet marjoramperfume. Those on the other hand which are weak and have not a powerful scent, since they are volatile and evaporate, also quickly come to an end: for

instance rose-perfume and kypros.

There are some however whose scent is even better on the second day, when any heavy quality that they possessed has evaporated. Some again are altogether more permanent, as spikenard and irisperfume, and the stronger 1 a perfume is, the longer it lasts. Again some perfumes for some reason keep their scent in the bath when the body is relaxed, or at least do not help to produce a disagreeable effect; while others become disagreeable and cause an even more unpleasant odour than the sweat, as though some sort of decomposition or decay took place.

Let this suffice for an account of the manufacture

and properties of perfumes.

Of the making of perfume-powders and compound perfumes.

XII. As to the mixing of solid substances to make powders 1 and compound perfumes, we do not find it here necessary to mix certain specified ingredients: the more numerous and the more various the perfumes that are mixed, the more distinguished and the more grateful will be the scent—just as though one were mixing whatever spices themselves 2 were procurable. As a matter of fact the custom is to use a mixture made of all kinds. Again in perfumes of this class the aim and object is not to make the mixture smell of some one particular thing, but to



 $^{^2}$ Sc. the natural products from which the manufactured perfumes are made.

κοινήν τινα την όσμην είναι. διὸ καὶ ἀνοίγοντες διά τινων ήμερων το έξόζον έξαιρουσιν άει και τῶν ἰσγυρῶν ἐλάττω μιγνύουσιν, ώσπερ . . . τὰ δ' όλως ου μιγνύουσιν, ώσπερ το έρυσίσκηπτρον.

ύπερ ου και άρτίως ελέγθη.

Βρέγουσι δὲ συντιθέντες τῷ οἴνω τῷ εὐώδει. ἔοικε δ' οὖν χρήσιμος εἶναι πρὸς τὰς εὐοσμίας, εἴ γε καὶ οἱ μυρεψοὶ χρώνται. μένουσι δὲ πολύν χρόνον αι συνθέσεις. ή δε χρήσις τούτων μεν είς την των ιματίων όσμην, των δε διαπασμάτων είς την στρωμνήν, όπως πρός τον χρώτα προσπίπτη. καὶ γὰρ ἄπτεται μᾶλλον καὶ έμμονώτερον τοῦτο, καὶ ώσπερ ἀντ' ἐκείνου τοῦτο ποιοῦσιν. οἱ δὲ πρότερον ενέβαλον οίνω καταβρέχοντες εὐώδει πρὸς τὸ παραιρεῖσθαι τὴν ὀσμήν, ἔνια δὲ καὶ μελικράτω καὶ οἴνω μιγνύντες ἀνέδευον, τὰ δὲ καὶ αὐτῷ τῷ μελικράτῳ. τὸ γὰρ ὅλον ἄμφω ταῦτα συνεργεῖ πρὸς εὐοσμίαν. διαμένουσι δὲ αί συνθέσεις. Φανερον δ' έκ τούτων ὅπερ καὶ πρότερον έλέγθη, διότι τὰ ξηρὰ καὶ εὐοσμότερα $\pi \rho \delta s \ \mathring{a} \lambda \lambda \eta \lambda a < \mu \iota \chi \theta \acute{e} \nu \tau a > \tau a \iota s \delta \sigma \mu a \iota s$.

Εὐλόγως δὲ τὰ μύρα φαρμακώδη διὰ τὴν τῶν 59 (61) ἀρωμάτων δύναμιν· καὶ γὰρ τὰ ἀρώματα τοιαῦτα. δηλοί δὲ τά τε καταπλάσματα καὶ α δή τινες

The reference of excluor is obscure.

3 μιγθέντα add. Turn.

¹ The example is missing. Turn. supplies costum et amomum from Plin. 13. 16, which does not however certainly refer to this passage; see 69, where this passage seems to be repeated.

CONCERNING ODOURS, 57-59

produce a general scent derived from them all. This is why every few days they open the vessel and remove each time that perfume whose scent is overpowering the others, adding at the same time smaller quantities of the less powerful scents, such as . . .,¹ while some perfumes are never added, such as

galingale, of which we spoke just now.

When they make compound perfumes, they moisten the spices with fragrant wine: and this certainly seems to be useful for producing fragrance, seeing that perfumers also use it. These compound perfumes last a long time. They are used to impart a pleasant odour to clothes, while the powders are used for bedding, so that they may come in contact with the skin: for this kind of preparation gets a better hold and is more lasting. so that men use it thus instead of scenting their bodies directly.2 Some, before putting the powder in the bedding, soak it in fragrant wine, so that it may acquire its scent: and some powders they moisten by mixing them with mead and wine, or again simply with mead. For in general both these things help to give them fragrance. Compound perfumes also last well. From which what was said above becomes manifest, inasmuch as solid perfumes, when mixed 3 with one another, acquire a greater fragrance.

*It is to be expected that perfumes should have medicinal properties in view of the virtues of spices: for these too have such virtues. The effects of

⁴ In W.'s text, which I have followed, there is some rearrangement (after Furlanus) of the order of sentences in this chapter and the next: e.g. part of § 61 is transferred to § 59. Both figures are retained for convenience of reference.

μαλάγματα καλοῦσιν οἵας ἀποδείκνυται δυνάμεις τά τε φύματα καὶ τὰ ἀποστήματα διαχέοντα καὶ ἄλλα πλείω τῶν κατὰ τὸ σῶμα διαλλοιοῦντα, ἐπιπολῆς μὲν ἀλλὰ καὶ τὰ ἐν βάθει, οἶον, ἄν τις καταπλάση τὰ ὑποχόνδρια καὶ τὸ στῆθος, εὐθὺς σὺν τοῖς ἐρυγμοῖς ἀποδίδωσιν εὐώδεις τὰς ὀσμάς. . . .

61 ΧΙΙΙ. Αί δὲ τῶν ζώων ὀσμαὶ κατὰ τὰς ἰδίας (59) γίνονται φύσεις· ἐκάστῷ γάρ ἐστί τις οἰκεία κατὰ τὴν κρᾶσιν. αὖται δ' ἡδεῖαι μὲν καὶ καθαραὶ [καὶ] κατὰ τὰς ἀκμὰς καὶ ὅταν εὖ ἔχωσιν ἑαυτῶν, ἔτι δὲ ἡδίους ἀπαλῶν καὶ νέων ὄντων. πλεῖσται δὲ καὶ κακωδέσταται περὶ τὰς ὀχείας καὶ ὅλως συντηκομένων καὶ καμνόντων σωμάτων· διὸ καὶ οἱ τράγοι καὶ οἱ ἔλαφοι καὶ λαγοὶ καὶ τἆλλα τότε μάλιστα ὄζει.

Θαυμαστὸν δὲ καὶ ἴδιον τὸ συμπάσχειν τὰς τραγέας, ὅταν ἡ ὅρα καθήκη τῆς ὁρμῆς. αἴτιον δὲ δηλονότι τὸ ὑπολείπεσθαί τινα ἐν τῷ δέρματι δύναμιν ἡ ὑγρότητα τοιαύτην ἀφ' ἡς ἡ ὁρμὴ γίνεται καὶ ζώντων κινουμένης οὖν καὶ διαθερμαινομένης ταύτης ὑπὸ τοῦ ἀέρος εὔλογον καὶ τὰ δέρματα καθ' ὅσον ἐπιβάλλει. διὸ καὶ ὡς

380

 $^{^{1}}$ § 60 on some other medicinal effects of perfumes is omitted. 2 $\kappa\alpha l$ bracketed by W.

CONCERNING ODOURS, 59-62

plasters and of what some call 'poultices' prove what virtues they display, since they disperse tumours and abscesses and produce a distinct effect on various other parts of the body, on its surface, but also on the interior parts: for instance, if one lays a plaster on his abdomen and breast, the patient forthwith produces fragrant odours along with his eructations.¹

Of the characteristic smells of animals, and of certain curious facts as to the smell of animal and vegetable products.

XIII. The smells of animals correspond to their several characters: each has a smell of its own according to its particular composition. These smells are pleasant and pure when 2 the animal is in its prime and in good condition, and even pleasanter when they 3 are young and tender. But the smell is strongest and least pleasant at the breeding season, and generally when the body is wasting or out of condition: wherefore goats stags hares and other animals have most smell at such times.

It is a remarkable fact and peculiar to the goat that goat-skins ⁴ are sympathetically affected when the breeding season comes round. The reason plainly is that there remains somehow in the hide the sort of virtue or moisture from which arises the breeding impulse when the animal is alive. It is natural therefore that, when this is excited and warmed by the air, the skin also should be excited so far as it belongs to it to be so affected. Wherefore the original cause as it were of the

4 i.e. the skin of a dead goat.

Digitized by Google

^{*} ἐαυτῶν can hardly be sound: ? αὐτὰ (sc. τὰ ζῶα).

[.] W. adds κινείσθαι after ἐπιβάλλει.

πρώτον αἴτιον ἡ διάθεσις· τότε γὰρ καὶ οἱ μὴ ὀχεύοντες ὄζουσι καὶ οἱ ἄγονοι καὶ αἱ αἶγες ὅλως. ἡ δ' ὀχεία τότε μὲν μεγάλην μερίδα συμβάλλεται, καθ' αὐτὴν δ' αἰτία γίνεται ἡ διάθεσις.

- 63 Συμβαίνει δὲ τρόπον τινὰ καὶ ἐν ἄλλοις ἡ τοιαύτη συμπάθεια· καὶ γὰρ ὁ οἰνος ἄμα τῆ σταφυλῆ δοκεῖ συνανθεῖν καὶ τὰ σκόροδα καὶ τὰ κρόμυα τότε δριμύτατον ὅζειν, ὅταν <τὰ> ἐν τῆ γῆ βλαστάνη· πλὴν τούτοις ἄμα συμβαίνει καὶ αὐτοῖς βλαστάνειν. ὅλως δὲ πάντα κινεῖται τὰ φλοιόριζα καὶ σαρκόριζα μὴ ἀπεξηραμμένα κατὰ τὰς βλαστητικὰς ὥρας· ἡ γὰρ ἐνυπάρχουσα δύναμις ἐν αὐτοῖς κινεῖται. θαυμασιώτατον δὲ τῶν τοιούτων τὸ ἐπὶ τοῦ στέατος τῆς ἄρκτου συμβαίνον, εἴπερ ἄμα ταῖς φωλίαις ἐπαίρεται καὶ ἐκπληροῖ τὰ ἀγγεῖα.
- 84 ΧΙΥ. Τί δή ποτε Δημόκριτος τοὺς μὲν χυμοὺς πρὸς τὴν γεῦσιν ἀποδίδωσι, τὰς δ' ὀσμὰς καὶ τὰς χρόας οὐχ ὁμοίως πρὸς τὰς ὑποκειμένας αἰσθήσεις; ἔδει γὰρ ἐκ τῶν σχημάτων. ἢ τοῦτό γε πρὸς ἄπαντας κοινόν; ἄπαντες γὰρ οἱ μὲν μόνης

¹ i.e. to form a 'crust.' 2 7à ins. Sch.

 ³ cf. H.P. 1. 6. 7.
 ⁴ i.e. when the fat of the living bear becomes abnormally developed. Plin. 8. 128. expresses his surprise at T.'s credulity 382

CONCERNING ODOURS, 62-64

phenomenon is the special condition of the animal at such periods: for at these times even those males which are not breeding have the smell, and the sterile goats and the females in general. Indeed, though at that particular time the fact that animals are actually breeding is a powerful factor in producing the smell, yet their condition is in itself a cause.

Similar sympathetic behaviour is found in a manner in other things also. Thus wine appears to 'bloom' 1 at the same time as the growing grape, and stored garlic and onions appear to have the most pungent smell at the season when those 2 in the ground are sprouting: however in this case sprouting takes place in the stored vegetables also. And in general any plant whose root is in layers 3 or fleshy becomes active at the season of sprouting, unless it has been completely dried: for it is the force latent in such plants which is stirred into activity. But the most remarkable phenomenon of the kind is what occurs with bears' grease: it makes active growth at the time of the bear's winter sleep 4 and completely fills the vessels in which it is kept.

Of odours as compared with other sense-impressions.

XIV. What can be the reason why Democritus, though he assigns various flavours to the sense of taste, yet does not in like manner assign various smells and colours to the senses to which they belong? According to his system he should have done so. Perhaps the same criticism should apply to all who have dealt with the subject: for they all

in this matter: his version (coctas ursorum carnes) adds to the marvel.

οί δὲ μάλιστα ταύτης τὰ πάθη λέγουσι καὶ τὰς διαφοράς, ὡς ἐν χρώμασι λευκὸν καὶ μέλαν, καὶ ἐν χυμοῖς γλυκὺ καὶ πικρόν, οὐχ οὕτω δ' ἐν ὀσμαῖς· οὐδὲν γὰρ πλὴν τό τ' εὔοσμον καὶ τὸ κάκοσμον. οὐδ' ἐν ἀπτοῖς· πλείω γὰρ εὐθὺ τὰ ὑποκείμενα, σκληρὸν μαλακὸν τραχὺ λεῖον.

65 'Αλλὰ μᾶλλον ἐν φωναῖς, ὀξὺ καὶ βαρύ. ἔτι δὲ τὰ μὲν μικτὰ τὰ δ' ἄμικτα. ἄμικτοι χυλοὶ οἱ μὲν τῷ μὴ καταμερίζεσθαι ὅστ' ἐξ ἀμφοῖν, οἶον ὕδωρ ἔλαιον φλέγμα αἶμα, ὅλως πᾶν τὸ ἐπινέον ἡ τὸ διαιροῦν, ὥσπερ τὸ ὄξος καὶ τὸ γάλα. τὸ γὰρ τῆ πιέσει καὶ τρίψει μιγνύμενον ἔτερον εἶδος. ἄλλον δὲ τρόπον οἱ μὴ εὔμικτοι πρὸς τὴν χρείαν ἡ καὶ λυμαινόμενοι ἀλλήλους, οἶον ἡ θάλαττα καὶ τὰ νιτρώδη καὶ πικρὰ ὕδατα τοὺς οἴνους καὶ τὰ πότιμα, ἐὰν μὴ εὐθὺς χρῆταί τις.

86 'Οσμαὶ δὲ αἱ μὲν οὕτως ἄμικτοι πλείους καὶ ὥστε καθόλου λαβεῖν αἱ κακώδεις ταῖς κακώδεσι. ὡς δὲ βέλτιόν τι τὸ ἐξ ἀμφοῖν ἔργον εὐρεῖν εἰ

¹ i.e. taste.

² There seems to be some confusion here, as in the first sentence of the section T. complained that colours are not classified. The following passage is unusually elliptical, and the text is probably defective. T.'s complaint seems to be that sense-experiences in general have been inadequately classified (cf. 2); and in 66 foll. he specially mentions smells.

CONCERNING ODOURS, 64-66

either give the various qualities and distinguish the experiences of this sense 1 alone or at least comparatively neglect the others: thus with colours 2 they distinguish white and black, and with flavours sweet and bitter, yet they make no corresponding classification of smells, but merely class them as 'pleasant' or 'unpleasant.' So too they fail to distinguish different experiences of the sense of touch, whereas several belong immediately to this sense, as hardness, softness, roughness, smoothness.

In sounds still more are there differences, as that between shrill and deep. Again some sense-experiences are simple, some compound. Flavours are simple first in the sense that they cannot be resolved into two components 3: instances are water oil phlegm blood,4 and in general anything which floats, like milk,5 or which causes separation, like vinegar. (Where mixture can be produced by pressure or crushing, it is quite a different matter). Secondly there are flavours which do not readily combine in another sense, namely for human use, or which even spoil one another if they are mixed, as sea-water, or water with soda in it or which has a bitter taste: these spoil wines or other things that are good to drink, unless they are taken at once.

Now the odours which in this sense do not combine are numerous, and, speaking generally, it is the pleasant odours which do not combine with the unpleasant ones. It would indeed be difficult, if not impossible, to find a case in which mixture is an

385

Sch. and W. after Turn. add < ἐν γίνεσθαι> after ἀμφοῖν, which seems unnecessary.

⁴ i.e. a liquid which, in one way or another, refuses to mix with another liquid.

⁵ ? cream.

μὴ ἀδύνατον, εἰς τὴν τοιαύτην δὲ δύναμιν <οὐχ>
ἄπαν ὡς εἰπεῖν πρὸς πᾶν εὔοσμον. ἀλλ' ἔνθα
μὲν ἴσως χείρω ποιεῖ ἔνθα δὲ βελτίω, καθάπερ
ἐπὶ τῶν μύρων τὰ μὲν γὰρ ἀφαιρεῖται τὸ ἄκρατον καὶ σκληρόν, τὰ δ' ἀποθηλύνει καὶ ὥσπερ
ἔξυδατοῖ τὰς ὀσμάς. ἐν δὲ τοῖς ξηροῖς ἄπασαι
πάσαις μικταί.

Τὰ γὰρ διαπάσματα ὅσφ ἃν ἢ πλειόνων ἀμείνω. ποιεῖ δὲ καὶ ἡ τοῦ οἴνου κατάμιξις καὶ μύρα ἔνια καὶ θυμιάματα εὐοσμότερα, καθάπερ τὴν σμύρναν. δοκεῖ δὲ καὶ τὸ μύρον ἡδύνειν τοὺς οἴνους, διὸ καὶ οἱ μὲν ἐν τἢ οἰνοποιτα μιγνύουσιν οἱ δὲ οὕτως ἐπιχεόμενον πίνουσιν. οὐκ ἄλογον δὲ συνέγγυς τὰς αἰσθήσεις οὕσας καὶ ἐν τοῖς αὐτοῖς ὑποκειμένοις ἔχειν τινὰ ἐπικοινωνίαν· ὡς γὰρ ἐπὶ τὸ πᾶν οὐδεὶς οὕτε χυλὸς ἄοσμος οὕτε ὀσμὴ ἄχυλος· τοῦτο δὲ ὅτι οὐδεμία ἐκ μὴ ἔχοντος χυλόν.

68 Συμβαίνει δὲ καὶ μεταβάλλειν τὰς ὀσμὰς ἄμα τοῖς χυλοῖς, ὥσπερ ἐπί τε τοῦ οἴνου καὶ ἐπὶ καρπῶν τινῶν· ἐνίων δὲ καὶ ἐν τῷ ἄνθει πρότερον, ὥσπερ τῶν βοτρύων· ἡ δὲ τῶν μύρων εἰς ἀκμὴν μόνον καὶ οἷον φθίσιν. μετακινοῦνται δ' ἐν ταῖς ἐτείαις ὥραις πάνθ' ὡς εἰπεῖν, μάλιστα δὲ τὰ

¹ I have inserted οὐχ, suggested by Sch.

² Like Sch. I fail to see the relevance of this remark. The sense required is 'while the fruit is still on the tree': 386

CONCERNING ODOURS, 66-68

improvement to the odour: in fact one might say that not¹ every combination of one fragrant thing with another will produce such a quality, but though sometimes the effect of such mixture is an improvement, sometimes it may be the reverse, as in the case of perfumes: for while the effect of some admixtures is to remove excessive strength or harshness, in other cases the odour is enfeebled and made, as it were, insipid. With solids however all combinations are possible.

In fact powders are the better, the more ingredients they have. Also the admixture of wine makes some perfumes and things used for incense more fragrant, for instance myrrh. It appears also that perfume sweetens wines, wherefore some add it in the manufacture, some put it in at the time of drinking. Nor is it unnatural that between these senses, since they are akin and are affected by the same objects, there should be a sort of reciprocity: for, to speak generally, no taste is unaccompanied by smell and no smell is unaccompanied by taste, the reason being that a thing which has no taste produces no smell.

It is also the case that smells actually change along with tastes, for instance in wine and certain fruits. And in some cases, as with grapes, the change takes place earlier, during 2 the flowering period: while in perfumes it occurs only when they have reached their best and are about, as it were, to go off. Almost all perfumes undergo alteration at certain seasons of the year, and this applies specially to the weakest kinds: in the case of those made

possibly $\delta \nu \theta \epsilon \hat{\imath}$ has got in from below and we should read $\kappa a \rho \pi \hat{\psi}$.

387

ἀσθενέστατα, τὰ δ' ἄνθινα καθ' ἡν ώραν ἀνθεῖ τὸ ăνθos.

[Τὰς συνθέσεις ποιοῦσιν ἐκ τῶν ἀρωμάτων: θραύσαντες πολλά και μίξαντες είς ταὐτὸ κλείουσιν είς κιβώτιον, είτ' ανοίγοντες διά τινων ήμερων ὅτι ἀν μάλιστα ὄζειν δοκή τοῦτ' αἴρουσι, καὶ πάλιν δὲ καὶ πάλιν διαλείποντες χρόνον, όπως αν μηδενός εξόζη. θαυμαστήν δ΄ οσμήν λαμβάνει τὰ ἱμάτια εἰς ταῦτα τιθέμενα.

Τὸ δὲ τῆς βαλάνου τῆς Αἰγυπτίας μύρον αὐτὸ μεν οὐκ ἄγαν ἀναπνεῖ, μιγνύμενον δε ποιεῖ τάλλα βελτίω μάλιστα δὲ τὴν ἰριν]. . . .

 $^{^{2}}$ cf. 57, of which this section seems to be a repetition. 3 cf. 15.

CONCERNING ODOURS, 68-69

from flowers this period is that at which the plants from which they are made are in bloom.1

[Compound 2 perfumes are made from spices: they bruise and mix a variety of these and shut them up together in a box. Then after a few days they open the box and take out the spice which seems to have the strongest smell: this treatment is repeated at intervals, so that the smell of no one ingredient may overpower the others. And clothes put away with such perfumes acquire a marvellous fragrance.

The perfume 3 made of the Egyptian balanos, though it has not much scent of its own, when mixed with others, especially iris-perfume, improves them]. . . . 4

389

⁴ The remaining sentences (§§ 70, 71) seem to be disconnected scraps, which perhaps do not belong to this treatise at all. The text of them being defective, it seems not worth while to attempt translation.

ΠΕΡΙ ΣΗΜΕΙΩΝ ΤΔΑΤΩΝ ΚΑΙ ΠΝΕΥΜΑ-ΤΩΝ ΚΑΙ ΧΕΙΜΩΝΩΝ ΚΑΙ ΕΤΔΙΩΝ

Ι. Σημεῖα ὑδάτων καὶ πνευμάτων καὶ χειμώνων καὶ εὐδιῶν ὧδε ἐγράψαμεν καθ' ὅσον ἢν ἐφικτόν,
 ὰ μὲν αὐτοὶ προσκοπήσαντες ὰ δὲ παρ' ἑτέρων

ούκ άδοκίμων λαβόντες.

Τὰ μεν οὖν ἐπὶ τοῖς ἄστροις δυομένοις καὶ ἀνατέλλουσιν ἐκ τῶν ἀστρονομικῶν δεῖ λαμβά2 νειν. εἰσὶ δὲ δύσεις διτταί· οἴ τε γὰρ ἀφανισμοὶ δύσεις εἰσί· τοῦτο δὲ ἐστιν ὅταν ἄμα συνδύνη τῷ ἡλίῳ τὸ ἄστρον, καὶ ὅταν ἀνατέλλοντος δύνη. ὁμοίως δὲ καὶ ἀνατολαὶ διτταί, αἱ μὲν ἑῷοι ὅταν προανατέλλη τοῦ ἡλίου τὸ ἄστρον, αἱ δὶ ἀκρόνυχοι ὅταν ἄμα δυομένῳ ἀνατέλλη.

Αι μεν οὖν τοῦ ἀρκτούρου λεγόμεναι ἀνατολαὶ ἀμφοτέρως συμβαίνουσιν· ἡ μεν γὰρ τοῦ χειμῶνος ἀκρόνυχός ἐστιν, ἡ δε μετωπορινὴ εωία. τῶν δ' ἄλλων αι πλεισται τῶν ὀνομαζομένων εῷαι, οἶον

Πλειάδος καὶ 'Ωρίωνος καὶ Κυνός.

3 Τῶν δὲ λοιπῶν σημείων ἔνια μὲν ἴδια κατὰ πάσας χώρας ἐστὶν ἐν ὅσαις ὅρη ὑψηλὰ καὶ αὐλῶνές εἰσι, μάλιστα δὲ ὅσα πρὸς θάλασσαν καθήκει τῶν ὑψηλῶν· τῶν τὲ γὰρ πνευμάτων ἀρχομένων τὰ νέφη προσπίπτει πρὸς τοὺς τοιούτους τόπους, καὶ μεθισταμένων εἰς τοὐναντίον

CONCERNING WEATHER SIGNS

Introductory: general principles.

I. The signs of rain wind storm and fair weather we have described so far as was attainable, partly from our own observation, partly from the informa-

tion of persons of credit.

Now those signs which belong to the setting or rising of the heavenly bodies must be learnt from astronomy.\(^1\) Their settings are twofold, since they may be said to have set when they become invisible. And this occurs when the star sets along with the sun, and also when it sets at sunrise. In like manner their risings are twofold: there is the morning rising, when the star rises before the sun, and there is the rising at nightfall, when it rises at sunset.

Now what are called the risings of Arcturus occur at both times, his winter rising being at nightfall and his autumn rising at dawn. But the rising of most of the familiar constellations is at dawn, for

instance, the Pleiad Orion and the Dog.

Of the remaining signs some belong specially to all such lands as contain high mountains and valleys, specially where such mountains extend down to the sea: for, when the winds begin to blow, the clouds are thrown against such places, and, when the winds

¹ Or, perhaps, 'from my astronomical works.'

ἀντιμεθίστανται καὶ ὑγρότερα γινόμενα διὰ βάρος εἰς τὰ κοῖλα συγκαθίζει. διὸ δεῖ προσέχειν οὖ ἄν τις ἱδρυμένος ἢ. ἔστι γὰρ ἀεί τινα λαβεῖν τοιοῦτον γνώμονα καὶ ἔστι σαφέστατα σημεῖα τὰ ἀπὸ τούτων.

Διὸ καὶ ἀγαθοὶ γεγένηνται κατὰ τόπους τινὰς ἀστρονόμοι ἔνιοι, οἶον Ματρικέτας ἐν Μηθύμνη ἀπὸ τοῦ Λεπετύμνου, καὶ Κλεόστρατος ἐν Τενέδφ ἀπὸ τῆς Ἰδης, καὶ Φαεινὸς ᾿Αθήνησιν ἀπὸ τοῦ Λυκαβηττοῦ τὰ περὶ τὰς τροπὰς συνείδε, παρ' οῦ Μέτων ἀκούσας τὸν τοῦ ἐνὸς δέοντα εἴκοσιν ἐνιαυτῶν <κύκλον> συνέταξεν. ἢν δὲ ὁ μὲν Φαεινὸς μέτοικος ᾿Αθήνησιν ὁ δὲ Μέτων ᾿Αθηναῖος. καὶ ἄλλοι δὲ τὸν τρόπον τοῦτον ἠστρολόγησαν.

Κλλα δέ ἐστι σημεῖα ἃ λαμβάνεται ἀπό τε ζώων τῶν κατ' οἰκίαν καὶ ἑτέρων τινῶν τόπων καὶ παθημάτων, μάλιστα δὲ κυριώτατα <τὰ> ἀπὸ τοῦ ἡλίου καὶ τῆς σελήνης ἡ γὰρ σελήνη νυκτὸς οἶον ἥλιός ἐστι· διὸ καὶ αἱ σύνοδοι τῶν μηνῶν χειμέριοί εἰσιν, ὅτι ἀπολείπει τὸ φῶς τῆς σελήνης ἀπὸ τετράδος φθίνοντος μέχρι τετράδος ἱσταμένου. ὥσπερ οὖν ἡλίου ἀπόλειψις γίνεται κατὰ τὸν ὅμοιον τρόπον καὶ τῆς σελήνης ἔκλειψις.
 δεῖ οὖν προσέχειν μάλιστα ταῖς ἀνατολαῖς ταῖς

6 δεῖ οὖν προσέχειν μάλιστα ταῖς ἀνατολαῖς ταῖς τούτων καὶ ταῖς δύσεσιν ὁποίας ἃν ποιῶνται τὸν βουλόμενον προγινώσκειν.

² Plin. 5. 140. Of Matriketas nothing is known.

¹ αντιμεθίστανται. ? αντιμεθίσταται.

³ Said (Plin. 2. 31) to have first recognised the Ram and the Archer. Athen. (7. 278 b) connects him with Tenedos. 39²

CONCERNING WEATHER SIGNS, 3-6

change, the clouds also change 1 and take a contrary direction, and, as they become laden with moisture, they settle down in the hollows because of their weight. Wherefore good heed must be taken to the local conditions of the region in which one is placed. It is indeed always possible to find such an observer, and the signs learnt from such persons are the most trustworthy.

Thus in some parts have been found good astronomers: for instance, ² Matriketas at Methymna observed the solstices from Mount Lepetymnos, Cleostratus ⁸ in Tenedos from Mount Ida, Phaeinos at Athens from Mount Lycabettus: Meton, who made the cycle ⁴ of nineteen years, was the pupil of the last-named. Phaeinos was a resident alien at Athens, while Meton was an Athenian. Others also have made astronomical observations in like manner.

Again there are other signs which are taken from domestic animals or from certain other quarters and happenings. Most important of all are the 5 signs taken from the sun and moon: for the moon is as it were a nocturnal sun. Wherefore also the meetings of the months are stormy, because the moon's light fails from the fourth day from the end of one month to the fourth day from the beginning of the next: there is therefore a failure of the moon corresponding to the failure of the sun. Wherefore anyone who desires to forecast the weather must pay especial heed to the character of the risings and settings of these luminaries.

⁵ τὰ seems necessary. ? κύρια τὰ.

Digitized by Google

⁴ Called 'the great year': cf. Aelian. V.H. 10. 7. τδν τοῦ ἐνὸς δέοντα εἴκοσιν ἐνιαυτῶν <κύκλον> conj. Sch. ἐνιαυτὸν conj. W.

Πρῶτον μὲν οὖν ληπτέον ὅτι αἱ διχοτομίαι διορίζουσι τὰς ὥρας, ὥστε ἐπὶ τούτων δεῖ ἀθρεῖν καὶ ἐνιαυτὸν καὶ μῆνα καὶ ἡμέραν. διχοτομεῖ δὲ τὸν μὲν ἐνιαυτὸν Πλειάς τε δυομένη καὶ ἀνατέλλουσα· ἀπὸ γὰρ δύσεως μέχρι ἀνατολῆς τὸ ἤμισυ τοῦ ἐνιαυτοῦ ἐστίν. ὥστε δίχα τέμνεται ὁ τ πᾶς χρόνος. ὁμοίως δὲ καὶ αἱ τροπαὶ καὶ ἰσημερίαι ποιοῦσιν. οἵα τις ἂν οὖν ἢ κατάστασις τοῦ ἀέρος Πλειάδος δυομένης, οὕτω ἔχει ὡς ἐπὶ τὸ πολὺ μέχρι τροπῶν, κᾶν μεταβάλλη, μετὰ τροπάς· ἐὰν δὲ μὴ μεταβάλλη, διέχει ἔως ἰσημερίας, κἀκείθεν ὡσαύτως μέχρι Πλειάδος, καὶ ἀπὸ ταύτης μέχρι τροπῶν θερινῶν, καὶ ἐντεῦθεν μέχρι ἰσημερίας, καὶ ἀπὸ ἰσημερίας μέχρι Πλειάδος δύσεως.

'Ως δ' αὕτως ἔχει καὶ περὶ τὸν μῆνα ἕκαστον διχοτομοῦσι γὰρ αἵ τε πανσέληνοι καὶ αἱ ὀγδόαι καὶ αἱ τετράδες, ὥστε ἀπὸ νουμηνίας ὡς ἀπὰ ἀρχῆς δεῖ σκοπεῖν. μεταβάλλει γὰρ ὡς ἐπὶ τὸ πολὺ ἐν τῆ τετράδι, ἐὰν δὲ μή, ἐν τῆ ὀγδόη, εἰ δὲ μή, πανσελήνω ἀπὸ δὲ πανσελήνου εἰς ὀγδόην φθίνοντος, καὶ ἀπὸ ταύτης εἰς τετράδα, ἀπὸ δὲ τετράδος εἰς τὴν νουμηνίαν.

Ως δ' αὕτως καὶ ἐπὶ τῆς ἡμέρας ἔχουσιν αὶ μεταβολαὶ ὡς ἐπὶ τὸ πολύ. ἀνατολὴ γὰρ καὶ πρωὶ καὶ ρέσημβρία καὶ δείλη καὶ δύσις, καὶ τὰ τῆς νυκτὸς μέρη τὰ ἀνάλογα ταὐτὸ ποιεῖ τοῖς εἰρημένοις περὶ πνευμάτων καὶ χειμῶνος καὶ εὐδίας. μάλιστα γὰρ ἐὰν μέλλη μεταβάλλειν, ἐν

CONCERNING WEATHER SIGNS, 6-9

Now the first point to be seized is that the various periods are all divided in half, so that one's study of the year the month or the day should take account of these divisions. The year is divided in half by the setting and rising of the Pleiad 1: for from the setting to the rising is a half year. So that to begin with the whole period is divided into halves: and a like division is effected by the solstices and equinoxes. From which it follows that, whatever is the condition of the atmosphere when the Pleiad sets, that it continues in general to be till the winter solstice, and, if it does change, the change only takes place after the solstice: while, if it does not change, it continues the same till the spring equinox: the same principle holds good from that time to the rising of the Pleiad. from that again to the summer solstice, from that again to the autumnal equinox, and from that to the setting of the Pleiad.

So too is it with each month; the full moon and the eighth 2 and the fourth days make divisions into halves, so that one should make the new moon the starting-point of one's survey. A change most often takes place on the fourth day, or, failing that, on the eighth, or, failing that, at the full moon; after that the periods are from the full moon to the eighth day from the end of the month, from that to the fourth day from the end, and from that to the new moon.

The divisions of the day follow in general the same principle: there is the sunrise, the mid-morning, noon, mid-afternoon, and sunset; and the corresponding divisions of the night have like effects in the matter of winds storms and fair weather; that is to say, if there is to be a change, it will generally

¹ Plin. 18. 280. ² cf. Arat. 73 f.

ταῖς διχοτομίαις μεταβάλλει. καθόλου μὲν οὖν τὰς ὥρας οὕτω δεῖ παρατηρεῖν, καθ΄ ἔκαστα δὲ τῶν σημείων κατὰ τὸν ὑπογεγραμμένον τρόπον.

- 10 "Υδάτος μὲν οὖν σημεῖα τὰ τοιαῦτα δοκεῖ εἶναι.
 ἐναργέστατον μὲν οὖν τὸ ἑωθινόν, ὅταν πρὸ ἡλίου
 ἀνατολῆς φαίνηται ἐπιφοινίσσον σημεῖον ἡ γὰρ
 αὐθημερινὸν ἐπισημαίνει ἡ τριῶν ἡμερῶν ὡς ἐπὶ
 τὸ πολύ. δηλοῖ δὲ καὶ τὰ ἄλλα σημεῖα· ἐὰν
 γὰρ μὴ πρότερον, τριταῖα μάλιστα σημαίνει
 τὸ ἐπιφοινίσσον καὶ δύνοντος, ἡττον δὲ ἡ τὸ
 ἑωθινοῦ.
- Καὶ ἐὰν δύνη χειμῶνος ἡ ἔαρος εἰς νεφέλιον, τριῶν ἡμερῶν ὡς τὰ πολλὰ ἐπισημαίνει. καὶ ἐὰν ῥάβδοι νοτόθεν, ταὐτὰ δὲ ταῦτα βορρᾶθεν γινόμενα ἀσθενέστερα. καὶ ἐὰν ἀνίσχων μέλαν σημεῖον ἴσχη, καὶ ἐὰν ἐκ νεφελῶν <ἀν>έχη, ὑδατικόν, καὶ ἐὰν ἀκτῖνες ἀνίσχοντος ἀνατείνωσι πρὶν ἀνατεῖλαι, κοινὸν ὕδατος σημεῖον καὶ ἀνέμου. καὶ ἐὰν καταφερομένου τοῦ ἡλίου ὑφίστηται νέφος, ὑφ' οῦ ἐὰν σχίζωνται αὶ ἀκτῖνες, χειμερινὸν τὸ σημεῖον. καὶ ὅταν καυματίας δύηται καὶ ἀνατέλλη, ἐὰν μὴ ἄνεμος γένηται, ὕδατος τὸ σημεῖον.

Τὰ αὐτὰ δὲ σημαίνει καὶ σελήνη πανσελήνω ἀνίσχουσα, ἀσθενέστερα δὲ ὁ μείς. ἐὰν μὲν ἢ πυρώδης, πνευματώδη σημαίνει τὸν μῆνα, ἐὰν δὲ ζοφώδης, ὑδατώδη· σημαίνει δὲ ὅτι ἂν σημαίνη

τριταίος ών ὁ μείς.

¹ τον ύπογ. τρόπον seems to mean the same as the Aristotelian τον υφηγημένον τρόπον, e.g. Eth. Nic. 2. 7. 9. The rendering 'the following method' would however suit the context.

CONCERNING WEATHER SIGNS, 9-12

occur at one of these divisions. In general therefore one should observe the periods in the way indicated, though as to particular signs we must follow the accepted method.¹

The signs of rain.

Now the signs of rain appear to be as follows: most unmistakable is that which occurs at dawn, when the sky has a reddish appearance before sunrise; for this usually indicates rain within three days, if not on that very day. Other signs point the same way: thus a red sky at sunset indicates rain within three days, if not before, though less

certainly than a red sky at dawn.

Again, if the sun sets in a cloud in winter or spring, this generally indicates rain within three days. So too, if there are streaks of light from the south, while, if these are seen in the north, it is a less certain sign. Again, if the sun when it rises has a black mark, or if it rises 2 out of clouds, it is a sign of rain; while, if at sunrise there are rays 3 shooting out before the actual rising, it is a sign of rain and also of wind. Again if, as the sun sinks, a cloud forms below it and this breaks up its rays, it is a sign of stormy weather. Again, if it sets or rises with a burning heat, and there is no wind, it is a sign of rain.

Moonrise gives similar indications, at the time of full moon: they are less certain when the moon is not full. If the moon looks fiery, it indicates breezy weather for that month, if dusky, wet weather; and, whatever indications the crescent moon gives, are given when it is three days old.

² ἀνέχη conj. Sch. ⁸ Plin. 18. 344.

13 'Αστέρες πολλοὶ διάττοντες ὕδατος ἢ πνεύματος, καὶ ὅθεν ἀν διάττωσιν ἐντεῦθεν τὸ πνεῦμα ἢ τὸ ὕδωρ. καὶ ἐὰν ἀκτῖνες ἀθρόαι ἀνίσχωσιν ἀνιόντος ἢ δύνοντος, σημεῖον <ὕδατος>. καὶ ὅταν ἀνίσχοντος τοῦ ἡλίου αἱ αὐγαὶ οἶον ἐκλείποντος χρῶμα ἴσχωσιν, ὕδατος σημεῖον. καὶ ὅταν νεφέλαι πόκοις ἐρίων ὅμοιαι ὧσιν, ὕδωρ σημαίνει. [ὑετοῦ δὲ σημεῖα] πομφόλυγες ἀνιστάμεναι πλείους ἐπὶ τῶν ποταμῶν ὕδωρ σημαίνουσι πολύ. ὡς δ' ἐπὶ τὸ πολὺ ἰρις περὶ λύχνον ἢ διὰ λύχνου διαφαινομένη νότια σημαίνει ὕδατα.

Καὶ οἱ μύκητες ἐὰν νότια ἢ, ὕδωρ σημαίνουσι, σημαίνουσι δὲ καὶ ἄνεμον κατὰ λόγον ὡς ᾶν ἔχωσι πλήθους καὶ μεγέθους, σμικροὶ δὲ καὶ κεγχρώδεις καὶ λαμπροὶ ὕδωρ καὶ ἄνεμον. καὶ ὅταν χειμῶνος τὴν φλόγα <ὁ λύχνος> ἀπωθῆ διαλιπὼν οἶον πομφόλυγας, ὕδατος σημεῖον, καὶ ἐὰν πηδῶσιν αἱ ἀκτῖνες ἐπ' αὐτόν, καὶ ἐὰν σπιν-

θηρες έπιγένωνται.

15 Θρυιθές λουόμενοι μὴ ἐν ὕδατι βιοῦντες ὕδωρ ἡ χειμῶνας σημαίνουσι. καὶ φρύνη λουομένη καὶ βάτραχοι μᾶλλον ἄδοντες σημαίνουσιν ὕδωρ. καὶ ἡ σαύρα φαινομένη, ἡν καλοῦσι σαλαμάνδραν, ἔτι δὲ καὶ χλωρὸς βάτραχος ἐπὶ δένδρου ἄδων ὕδωρ σημαίνει. χελιδόνες τῆ γαστρὶ τύπτουσαι τὰς λίμνας ὕδωρ σημαίνουσι. βοὺς τὴν προσθίαν ὁπλὴν λείξας χειμῶνα ἡ ὕδωρ σημαίνει. ἐὰν

cf. 37.
 δατος ins. Furl.
 Plin. 18. 356.
 δετοῦ δὲ σημεῖα bracketed by Sch.

^a Plin. 18, 356.
^a νετού δε σημεία bracketed by Sch cf. Arist. Meteor. 3. 4; Plut. Quaest. Nat. 1. 2.

cf. 42.

⁸ i.e. breaks up into small 'grains' (?). cf. 25, 42, 54.

CONCERNING WEATHER SIGNS, 13-15

Many shooting ¹ stars are a sign of rain or wind, and the wind or rain will come from that quarter from which they appear. Again, if at sunrise or sunset the sun's rays appear massed together, it is a sign of rain.² Also it is a sign of rain when at sunrise the rays ³ are coloured as in an eclipse; and also when there are clouds ⁴ like a fleece of wool. The rising of bubbles ⁵ in large numbers on the surface of rivers is a sign of abundant rain. And in general, when a rainbow ⁶ is seen round or through a lamp, it signifies rain from the south.

Again, if the wind is from the south, the snuff? of the lamp-wick indicates rain; it also indicates wind in proportion to its bulk and size: while if the snuff is small, like millet-seed, and of bright colour, it indicates rain as well as wind. Again, when in winter the lamp rejects the flame but catches, as it were, here and there in spurts, it is a sign of rain: so also is it, if the rays of light leap

up on the lamp, or if there are sparks.

It is a sign of rain or storm when birds which are not aquatic take a bath. It is a sign of rain when a toad takes a bath, and still more so when frogs are vocal. So too is the appearance of the lizard known as 'salamander,' 10 and still more the chirruping of the green frog in a tree. It is a sign of rain when swallows 11 hit the water of the lakes with their belly. It is a sign of storm or rain when the ox

10 cf. de igne 60, where it is explained why the salamander

puts fire out.

¹¹ Plin. 18. 363; Verg. Georg. 1. 377.

⁹ i.e. refuses to light properly. The appearance seems to be that described Verg. Georg. 1. 391 (scintillare oleum). In the same passage putres concrescere fungos perhaps illustrates the comparison of the snuff to millet-seed above.

δὲ εἰς τὸν οὐρανὸν ἀνακύπτων ὀσφραίνηται, ὕδωρ σημαίνει.

Κορώνη ἐπὶ πέτρας κορυσσομένη ἢν κῦμα κατακλύζει ὕδωρ σημαίνει καὶ κολυμβῶσα πολλάκις καὶ περιπετομένη ὕδωρ σημαίνει. κόραξ πολλὰς μεταβάλλειν εἰωθὼς φωνάς, τούτων ἐὰν ταχὺ δὶς φθέγξηται καὶ ἐπιρροιζήση καὶ τινάξη τὰ πτέρα, ὕδωρ σημαίνει. καὶ ἐὰν ὑετῶν ὄντων πολλὰς μεταβάλλη φωνὰς καὶ ἐὰν φθειρίζηται ἐπ' ἐλαίας. καὶ ἐἀν τε εὐδίας ἐάν τε ὕδατος ὄντος μιμῆται τῆ φωνῆ οἶον σταλαγμούς, ὕδωρ σημαίνει. ἐάν τε κόρακες ἐάν τε κολοιοὶ ἄνω πέτωνται καὶ ἰερακίζωσιν, ὕδωρ σημαίνουσι. καὶ ἐὰν κόραξ εὐδίας μὴ τὴν εἰωθυῖαν φωνὴν ἵη καὶ ἐπιρροιβδῆ, ὕδωρ σημαίνει.

Τ΄ Ἐὰν ἱέραξ ἐπὶ δένδρου καθεζόμενος καὶ εἴσω εἰσπετόμενος φθειρίζηται, ὕδωρ σημαίνει. καὶ θέρους ὅταν πολλοὶ ἀθρόοι φανῶσιν ὅρνιθες οἱ βιοτεύουσιν ἐν νήσῳ, ὕδωρ σημαίνουσιν ἐὰν δὲ μέτριοι, ἀγαθὸν αἰξὶ καὶ βοτοῖς, ἐὰν δὲ πολλοὶ ὑπερβολῆ, αὐχμὸν ἰσχυρόν. ὅλως δὲ ὅρνιθες καὶ ἀλεκτρυόνες φθειριζόμενοι ὑδατικὸν σημεῖον, καὶ

όταν μιμώνται ύδωρ ώς δον.

Καὶ ἡ νῆττα ἥμερος <ἐὰν> ὑπιοῦσα ὑπὸ τὰ γεῖσα ἀποπτερυγίζηται, ὕδωρ σημαίνει, ὁμοίως δὲ καὶ κολοιοὶ καὶ ἀλεκτρυόνες, ἐάν τε ἐπὶ λίμνη ἡ θαλάττη ἀποπτερυγίζωνται, ὡς νῆττα ὕδωρ ση-

¹ Plin. 18. 364; Verg. Georg. 1. 375.

δετῶν ὅντων can hardly mean 'while it is raining.'

² ἐπιρροιζήση. Sc. with his wings probably; not, as LS. 'croaks.' Plin. (18. 362) seems to have had a fuller text, or to have drawn also on some other authority.

CONCERNING WEATHER SIGNS, 15-18

licks his fore-hoof; if he puts 1 his head up towards the sky and snuffs the air, it is a sign of rain.

It is a sign of rain when a crow puts back its head on a rock which is washed by waves, or when it often dives or hovers over the water. It is a sign of rain if the raven, who is accustomed to make many different sounds, repeats one of these twice quickly and makes a whirring 2 sound and shakes his wings. So too if, during a rainy season, he utters many different sounds, or if he searches for lice perched on an olive-tree. And if, whether in fair or wet weather, he imitates, as it were, with his voice falling drops, it is a sign of rain. So too is it if ravens or jackdaws fly high and scream 4 like hawks. And, if a raven in fair weather does not utter his accustomed note and makes a whirring with his wings, it is a sign of rain.

It is a sign of rain if a hawk perches on a tree, flies right into it and proceeds to search for lice: also, when in summer a number of birds living on an island pack together: if a moderate number collect, it is a good sign for goats and flocks, while if the number is exceedingly large, it portends a severe drought. And in general it is a sign of rain when cocks and hens search for lice; as also when they

make a noise like that of falling rain.

Again it is a sign of rain when a tame 6 duck gets under the eaves and flaps its wings. Also it is a sign of rain when jackdaws and fowls flap their wings whether on a lake or on the sea—like the duck. It

⁸ ήμερος. ? ή ήμερος.

401

VOL. II.

D D

⁴ ieρaκίζωσιν. ? 'hover like hawks.' However, Arat. 231 understood it to refer to the voice: so LS.

⁵ ἐπιρροιβδη̂. Exact sense uncertain. cf. Soph. Ant. 1004.

μαίνει. καὶ ἐρωδιὸς ὄρθριον φθεγγόμενος ὕδωρ ἡ πνεῦμα σημαίνει καὶ ἐὰν ἐπὶ θάλατταν πετόμενος βοα, μαλλον ύδατος σημείον ή πνεύματος, καὶ ὅλως Βοῶν ἀνεμῶδες.

19 Καὶ ὁ σπίνος ἐν οἰκία οἰκουμένη ἐὰν φθέγξηται εωθεν, ύδωρ σημαίνει ή χειμώνα. και χύτρα σπινθηρίζουσα πάσα περίπλεως ύδατος σημείον. καὶ ἴουλοι πολλοὶ πρὸς τοῖχον ἔρποντες ὑδατικόν. δελφίς παρά γην κολυμβών και αναδυόμενος

πυκνὰ ὕδωρ ἡ χειμῶνα σημαίνει.

"Υμηττος ελάττων, ἄνυδρος καλούμενος, εαν τώ κοίλφ νεφέλιον έχη, ύδατος σημείον καὶ έὰν ὁ μέγας "Υμηττος τοῦ θέρους ἔχη νεφέλας ἄνωθεν καὶ ἐκ πλαγίου, ὕδατος σημείου. καὶ ἐὰν ὁ ἄνυδρος "Υμηττος λευκάς έχη ἄνωθεν καὶ έκ πλαγίου. καὶ ἐὰν περὶ ἰσημερίαν λὶψ πνεύση, ὕδωρ ση-

μαίνει.

Αί δὲ βρονταὶ αί μὲν χειμεριναὶ καὶ έωθιναὶ μαλλον <άνεμον ή> ὕδωρ σημαίνουσιν αί δὲ θεριναί μεσημβρίας καὶ έσπεριναί βρονταὶ ύδατικὸν σημείον. ἀστραπαὶ δὲ ἐάν γε πανταχόθεν γένωνται, ΰδατος αν η ανέμου σημειον. και έσπεριναί ώσαύτως, και έαν ακρωρίας νότου πνέοντος νοτόθεν ἀστράψη, ὕδωρ σημαίνει ἡ ἄνεμον. καὶ ζέφυρος ἀστράπτων προς βορείου η χειμώνα ή ύδωρ σημαίνει. καὶ θέρους αί έσπέριαι άστραπαὶ ὕδωρ αὐτίκα σημαίνουσιν ἡ

cf. Plin. 18. 361; Cic. Div. 2, 70. 8 ἐὰν τφ. ? ἐὰν ἐν τφ.

¹ Sch. cites Plin. 18. 364, vermes terreni erumpentes, as representing this, which seems doubtful.

CONCERNING WEATHER SIGNS, 18-21

is a sign of wind or rain when a heron utters his note at early morning: if, as he flies towards the sea, he utters his cry, it is a sign of rain rather than of wind, and in general, if he makes a loud cry, it portends wind.

It is a sign of rain or storm if a chaffinch kept in the house utters its note at dawn. It is also a sign if any pot filled with water causes sparks to fly when it is put on the fire. It is also a sign of rain when a number of millepedes 1 are seen crawling up a wall. A dolphin 2 diving near land and frequently reappearing indicates rain or storm.

If the lesser Mount Hymettus, which is called the Dry Hill, has cloud in ³ its hollows, it is a sign of rain: so also is it, if the greater Hymettus has clouds in summer on the top and on the sides: or if the Dry Hymettus has white clouds on the top and on the sides; also if the south-west wind ⁴ blows at

the equinox.

Thunder in winter and at dawn indicates wind be rather than rain; thunder in summer at midday or in the evening is a sign of rain. If lightning is seen from all sides, it will be a sign of rain or wind, and also if it occurs in the evening. Again, if when the south wind be is blowing at early dawn, there is lightning from the same quarter, it indicates rain or wind. When the west wind is accompanied by lightning from the north, it indicates either storm or rain. Lightning in the evening in summer time indicates rain within three days, if not immediately.

403

⁴ cf. Arist. Probl. 26. 26.

ἄνεμον ἡ add. Furl. from Plin. 18. 354.
 cf. Soph. Aj. 257; Arist. Probl. 26. 20.

⁷ akowolas. cf. 42. So Arat. 216 renders.

τριών ήμερών. καὶ ὀπώρας βορράθεν ἀστραπαὶ ύδατικὸν σημεῖον.

- 2 'Η Εὔβοια ὅταν διαζωσθῆ μέση, ὕδωρ διὰ ταχέων. καὶ ἐὰν ἐπὶ τὸ Πήλιον νεφέλη προσίζη, ὅθεν ἃν προσίζη, ἐντεῦθεν ὕδωρ ἡ ἄνεμον σημαίνει. ὅταν ἰρις γένηται, ἐπισημαίνει ἐάν τε πολλαὶ ἔριδες γένωνται, σημαίνει ὕδωρ ἐπὶ πολύ. ἀλλὰ πολλάκις καὶ οἱ ὀξεῖς ἥλιοι, ὅταν ἐκ νεφέλης. μύρμηκες ἐν κοίλῳ χωρίφ ἐὰν τὰ ῷὰ ἐκφέρωσιν ἐκ τῆς μυρμηκιᾶς ἐπὶ τὸ ὑψηλὸν χωρίον, ὕδωρ σημαίνουσιν, ἐὰν δὲ καταφέρωσιν, εὐδίαν. ἐὰν παρήλιοι δύο γένωνται καὶ ὁ μὲν νοτόθεν ὁ δὲ βορρᾶθεν, καὶ ἄλως ἄμα ὕδωρ διὰ ταχέων σημαίνουσι. καὶ ἄλως αι μέλαιναι ὑδατικὸν καὶ μᾶλλον αι δείλης.
- 23 'Εν τῷ Καρκίνω δύο ἀστέρες εἰσίν, οἱ καλούμενοι 'Όνοι, ὧν τὸ μεταξὺ τὸ νεφέλιον ἡ Φάτνη
 καλουμένη. τοῦτο ἐὰν ζοφῶδες γένηται, ὑδατικόν.
 ἐὰν μὴ ἐπὶ Κυνὶ ὕση ἡ ἐπὶ 'Αρκτούρω, ὡς ἐπὶ
 τὸ πολὺ πρὸς ἰσημερίαν ὕδωρ ἡ ἄνεμος. καὶ τὸ
 δημόσιον τὸ περὶ τὰς μυίας λεγόμενον ἀληθές:
 ὅταν γὰρ δάκνωσι σφόδρα, ὕδατος σημεῖον.
 σπίνος φθεγγόμενος ἔωθεν μὲν ὕδωρ σημαίνει ἡ χειμῶνα, δείλης δὲ ὕδωρ.

Της δὲ νυκτὸς ὅταν τὸν "Υμηττον κάτωθεν τῶν ἄκρων νεφέλη διαζώση λευκὴ καὶ μακρά, ὕδωρ γίνεται ὡς τὰ πολλὰ μετρίων ἡμερῶν. καὶ ἐὰν

¹ Evidently an Attic saying, of days when only the upper part of the Eubocan mountains was visible.

CONCERNING WEATHER SIGNS, 21-24

Lightning from the north in late summer is a sign of rain.

1 When Euboea has a girdle about it up to the waist, there will be rain in a short space. If cloud clings about Mount Pelion, it is an indication of rain or wind from the quarter to which it clings. When a rainbow appears, it is an indication of rain: if many rainbows appear, it is an indication of longcontinued rain. So too is it often when the sun appears 2 suddenly out of cloud. It is a sign of rain if ants 3 in a hollow place carry their eggs up from the ant-hill to the high ground, a sign of fair weather if they carry them down. If two mocksuns 4 appear, one to the south, the other to the north, and there is at the same time a halo, these indicate that it will shortly rain. A dark halo round the sun indicates rain, especially if it occurs in the afternoon.

In the Crab are two stars called the Asses, and the nebulous space between them is called the Manger⁵; if this appears dark, it is a sign of rain. If there is no rain at the rising of the Dog or of Arcturus, there will generally be rain or wind towards the equinox. Also the popular saying about flies is true; when they bite excessively, it is a sign of rain. If a chaffinch 6 utters its note at dawn, it is a sign of rain or storm, if in the afternoon, of rain.

When at night a long stretch of white cloud encompasses Hymettus below the peaks, there will generally be rain in a few days. If cloud settles on

² cf. H.P. 8. 10. 3.

³ Plin. 18. 364; Verg. Georg. 1. 379. ⁴ cf. 29. ⁵ cf. 43, 51.

⁶ cf. 19, of which this seems to be in part a repetition.

έν Λίγίνη [καὶ] ἐπὶ τοῦ Διὸς τοῦ Ἑλλανίου νεφέλη καθίζηται, ὡς τὰ πολλὰ ὕδωρ γίνεται. ἐὰν ὕδατα πολλὰ γίνηται χειμερινά, τὸ ἔαρ ὡς τὰ πολλὰ γίνεται αὐχμηρόν ἐὰν δ' αὐχμηρὸς ὁ χειμών, τὸ ἔαρ ὑδατῶδες. ὅταν χιόνες πολλαὶ γίνωνται, ὡς τὰ πολλὰ εὐετηρία γίνεται.

25 Φασὶ δέ τινες καὶ εἰ ἐν ἄνθραξι λαμπρὰ χάλαζα ἐπιφαίνηται, χάλαζαν προσημαίνειν ὡς τὰ πολλά· ἐὰν δὲ ὥσπερ κέγχροι μικροὶ λαμπροὶ πολλοί, ἀνέμου μὲν ὄντος εὐδίαν, μὴ ἀνέμου δὲ ὕδωρ ἡ ἄνεμον. ἔστι δ' ἄμεινον πρῶτον γίνεσθαι βόρειον ὕδωρ νοτίου καὶ τοῖς φυομένοις καὶ τοῖς ζώοις· δεῖ δὲ γλυκὸ εἶναι καὶ μὴ ἀλμυρὸν τοῖς γευομένοις. καὶ ὅλως ἔτος βέλτιον νοτίου βόρειον καὶ ὑγιεινότερον. καὶ ὅταν <πάλιν> ὀχεύωνται πρόβατα ἡ αἶγες, χειμῶνος μακροῦ σημεῖον.

28 ΙΙ. "Υδατος μέν οὖν ταῦτα λέγεται σημεῖα· ἀνέμου δὲ καὶ πνευμάτων τάδε. ἀνατέλλων ὁ ἥλιος καυματίας, κᾶν μὴ ἀποστίλβῃ, ἀνεμῶδες τὸ σημεῖον· καὶ ἐὰν κοῦλος φαίνηται ὁ ἥλιος, ἀνέμου ἡ ὕδατος τὸ σημεῖον. καὶ ἐὰν ἐπὶ πολλὰς ἡμέρας καυματίας, αὐχμοὺς καὶ ἀνέμους πολυχρονίους σημαίνει. ἐὰν αὶ ἀκτῖνες αὶ μὲν πρὸς βορρᾶν αἱ δὲ πρὸς νότον σχίζωνται τούτου μέσου

406

¹ So called also by Pind. Nem. 5. 19. Paus. 2. 30. 3 calls it the temple of Zevs Πανελλήνιος. και bracketed by Sch. ² cf. C.P. 2. 2.

CONCERNING WEATHER SIGNS, 24-26

the temple of Zeus Hellanios¹ in Aegina, usually rain follows. If a great deal of rain falls in winter, the spring is usually dry; if the winter has been dry, the spring is usually wet. When there is much² snow in winter, a good season generally follows.

Some say that, if in the embers ⁸ there is an appearance as of shining hail-stones, it generally prognosticates hail; while, if the appearance is like a number of small shining millet-seeds, ⁴ it portends fair weather, if there is wind at the time, but, if there is no wind, rain or wind. It is better both for plants and for animals that rain should come from the north before it comes from the south; it must however be fresh and not briny to the taste. And in general a season ⁵ in which a north wind prevails is better and healthier than one in which southerly winds prevail. It is a sign of a long winter when sheep or goats have a second ⁶ breeding season.

The signs of wind.

II. Such then are said to be the signs of rain. The following are signs of wind and breezes. ⁷ If the sun rises with a burning heat but does not shine brilliantly, it is a sign of wind. If the sun has a hollow appearance, it is a sign of wind or rain. If it blazes with a burning heat for several days, it portends long-continued drought or wind. If at dawn its rays are parted, some pointing to the north and some

⁷ Plin. 18. 342.

³ ἄνθραξι conj. Sch., supported by Plin. 18. 358; Arat. 309. ἀστράσι MSS.

⁴ cf. 14, 42, 54. ⁵ cf. C.P. 2. 2.

⁶ πάλιν ins. Sch.; text probably defective.

όντος κατ' όρθρον, κοινὸν ὕδατος καὶ ἀνέμου

σημειόν έστιν.

Τ΄ Εστι δὲ σημεῖα ἐν ἡλίῳ καὶ σελήνη, τὰ μὲν μέλανα ὕδατος τὰ δ' ἐρυθρὰ πνεύματος. ἐὰν δὲ καὶ ὁ μεὶς βορείου ὄντος ὀρθὸς εἰστήκη, ζέφυροι εἰώθασιν ἐπιπνεῖν καὶ ὁ μὴν χειμερινὸς διατελεῖ. ὅταν μὲν ἡ κεραία <ἡ ἄνω> τοῦ μηνὸς ἐπικύπτη, βόρειος ὁ μείς・ ὅταν δ' ἡ κάτωθεν, νότιος・ ἐὰν δ' ὀρθὸς καὶ μὴ καλῶς ἐγκεκλιμένος μέχρι τετράδος καὶ εὔκυκλος, εἴωθε χειμάζειν μέχρι διχομηνίας. σημαίνει ζοφώδης μὲν ὧν ὕδωρ πυρώδης δὲ πνεῦμα.

Αἴθυιαι καὶ νῆτται [πτερυγίζουσαι] καὶ ἄγριαι καὶ τιθασσαὶ ὕδωρ μὲν σημαίνουσι δυόμεναι, πτερυγίζουσαι δὲ ἄνεμον. οἱ κέπφοι εὐδίας οὕσης ὅποι ἀν πέτωνται ἄνεμον προσημαίνουσι. στρουθοὶ χειμῶνος ἀφ' ἐσπέρας θορυβοῦντες ἡ ἀνέμου μεταβολὴν σημαίνουσιν ἡ ὕδωρ ὑέτιον. ἐρωδιὸς ἀπὸ θαλάσσης πετόμενος καὶ βοῶν πνεύματος σημεῖόν ἐστι· καὶ ὅλως βοῶν μέγα

ἀνεμώδης.

Κύων κυλινδούμενος χαμαὶ μέγεθος ἀνέμου σημαίνει. ἀράχνια πολλὰ φερόμενα πνεῦμα ἡ χειμῶνα σημαίνει. ἡ ἄμπωτις βόρειον πνεῦμα σημαίνει, πλημμύρα δὲ νότιον. ἐὰν μὲν γὰρ ἐκ βορείων πλημμύρα ἡκŋ, εἰς νότιον μεταβάλλει, ἐὰν δ' ἐκ νοτίων ἄμπωτις γίνηται, εἰς βόρειον

¹ Plin. 18. 343 suggests that this is the meaning: text perhaps defective. cf. Verg. Georg. 1, 445.

³ Lit. 'the crescent moon has a northerly character.' ἡ ἄνω add. Furl.

CONCERNING WEATHER SIGNS, 26-29

to the south, while the orb itself is 1 clearly seen

between, it is a sign of rain and wind.

Also black spots on the sun or moon indicate rain, red spots wind. Again, if, while a north wind blows, the horns 2 of the crescent moon stand out straight, westerly winds will generally succeed, and the rest of the month will be stormy. When the upper horn of the crescent moon is bent, northerly winds 3 will prevail for that part of the month: when the lower horn is bent, southerly winds will prevail. 4 If however the horns up to the fourth day point straight and have not a graceful bend inwards but round to a circle, it will generally be stormy till the middle of the month. If the moon is dusky, it indicates rain, if fiery, it indicates wind.

It is a sign of rain when gulls and ducks, whether wild or tame, plunge under water, a sign of wind when they flap their wings. Wherever the bird called kepphos flies during a calm, it is a sign of coming wind. If sparrows in winter begin to be clamorous at evening, it is a sign of a coming change or of a fall of rain. A heron flying from the sea and screaming is a sign that a breeze is coming: so is it in general a sign of wind when he screams loudly.

A dog rolling on the ground is a sign of violent wind. A number of cobwebs 5 in motion portends wind or storm. The ebb-tide indicates a north wind, the flowing tide a wind from the south. For, if the flowing tide sets from the north, there is a change to the south, and if an ebb-tide comes from the south, there is a change to the north. It is

Digitized by Google

⁴ cf. 38; Plin. 18. 347; Verg. Georg. i. 428; the English sign, 'the young moon with the old moon in her arm.'

⁵ Plin, 11, 84; Arist, Probl. 26, 61.

μεταβάλλει. θάλασσα οἰδοῦσα καὶ ἀκταὶ βοῶσαι καὶ αἰγιαλὸς ἡχῶν ἀνεμώδης. καὶ ὁ μὲν βορέας λήγων ἐλάττων ὁ δὲ νότος ἀρχόμενος. παρήλιος

οπόθεν αν ή ύδωρ ή άνεμον σημαίνει.

30 Ἡ πέμπτη καὶ δεκάτη ἀπὸ τροπῶν τῶν χειμερινῶν ὡς τὰ πολλὰ νότιος. βορείων δὲ γινομένων ξηραίνει πάντα, νοτίων δὲ ὑγραίνει. ἐὰν δὲ νοτίων ὄντων ψοφῆ <τι> τῶν κεκολλημένων, εἰς τὰ νότια σημαίνει τὴν μεταβολήν ἐὰν δὲ πόδες οἰδῶσι, νοτία ἡ μεταβολή. τὸ δὲ αὐτὸ σημεῖον καὶ ἐκνεφίου. καὶ ὀδαξῶν τὸν δεξιόν. ἐχῦνος ὁ χερσαῖος σημαντικόν ποιεῖται δὲ δύο ὀπὰς ὅπου ᾶν οἰκῆ, τὴν μὲν πρὸς βορρᾶν τὴν δὲ νοτόθεν ὁποτέραν δ' ᾶν ἀποφράττη, ἐντεῦθεν πνεῦμα σημαίνει, ἐὰν δ' ἀμφοτέρας, ἀνέμου μέγεθος.

81 Έλν όρος ..., πρὸς βορρὰ ἄνεμον προσημαίνει. ἐὰν ἐν θαλάττη ἐξαίφνης πνεύματος γαλήνη γίνηται, μεταβολὴν πνεύματος ἡ ἐπίδοσιν. ἐὰν ἄκραι μετέωροι φαίνωνται ἡ καὶ νῆσοι ἐκ μιᾶς πλείους, νοτίαν μεταβολὴν σημαίνει· γῆ τε μέλαινα ὑποφαινομένη <βόρειον>, λευκὴ δὲ νότιον. αἱ ἄλωνες περὶ τὴν σελήνην πνευματώδεις μᾶλλον ἡ περὶ ἥλιον· σημαίνουσι δὲ πνεῦμα ῥαγεῖσαι περὶ ἄμφω, καὶ ἡ ἄν ῥαγῆ ταύτη πνεῦμα. ἐπι-

² cf. Arist. Probl. 26. 12 ad fin.

⁴ νότια MSS.; βόρεια conj. Furl., surely with good reason. cf. Arist. Probl. 1. 24.

¹ cf. 40; Plin. 18. 359; Verg. Georg. 1. 356.

³ ξηραίνει, ὑγραίνει seem to be used quasi-imporsonally; but the text is perhaps defective.

After δεξιδν Sch. and W. mark a lacuna, which does not seem necessary.
 cf. Arist. H.A. 9. 6 ad fin.

CONCERNING WEATHER SIGNS, 20-31

a sign of wind when the sea 1 has a swell or promontories moan or there is loud noise on the beach. Now the north wind has less force as it ceases to blow, the south wind as it begins. A mock sun, in whatever quarter it appears, indicates rain or wind.

The fifteenth 2 day after the winter solstice is generally marked by southerly winds. If there is a northerly wind, everything gets dried 3 up, if a southerly, there is abundant moisture. If, while a south wind is blowing, glued articles make a cracking sound, it indicates a change to a south 4 wind. the feet swell, there will be a change to a south wind. This also sometimes indicates a hurricane. So too does it, if a man has a shooting pain in the right foot. The behaviour of the hedgehog is also significant: this animal makes two holes wherever he lives, one towards the north, the other towards the south: now whichever hole he blocks up, it indicates wind from that quarter, and, if he closes both, it indicates violent wind.

If a mountain 7 it indicates wind from the north. If at sea during a wind there is a sudden calm, it indicates a change or an increase of wind. If promontories 8 seem to stand high out of the sea, or a single island looks like several, it indicates a change to south wind. If the land looks black from the sea, it indicates a north wind,9 if white, a south wind. A halo 10 about the moon signifies wind more certainly than a halo about the sun: but in either case, if there is a break in the halo, it indicates wind, which will come from the quarter in which the break is. If the sky is overcast in whatever quarter

⁷ I have marked a lacuna after δροs. Furl. renders si mons versus aquilonem extenditur, venti signum est, with what meaning I cannot see.

8 cf. Arist. Meteor. 3. 4 ad init.

9 βόρειον add. Furl.

10 cf. 51.

νεφέλων ὅθεν αν ἀνατέλληται, ἐντεῦθεν ἄνεμος. αὶ κηλάδες νεφέλαι θέρους ἄνεμον σημαίνουσι.

2 'Εὰν ἀστραπὴ πανταχόθεν γίνηται, ὕδωρ σημαίνει, καὶ ὅθεν ὰν αἱ ἀστραπαὶ πυκναὶ γίνωνται, ἐντεῦθεν πνεύματα γίνεται. θέρους ὅθεν ὰν ἀστραπαὶ καὶ βρονταὶ γίνωνται, ἐντεῦθεν πνεύματα γίνεται ἰσχυρά· ἐὰν μὲν σφόδρα καὶ ἰσχυρὸν ἀστράπτη, θᾶττον καὶ σφοδρότερον πνεύσουσιν, ἐὰν δ' ἡρέμα καὶ μανῶς, κατ' ὀλίγον. τοῦ δὲ χειμῶνος καὶ φθινοπώρου τοὐναντίον παύουσι γὰρ τὰ πνεύματα αἱ ἀστραπαί· καὶ ὅσω ὰν ἰσχυρότεραι γίνωνται ἀστραπαὶ καὶ βρονταί, τοσούτω μᾶλλον παύονται· τοῦ δ' ἔαρος ἡττον ἄν ταὐτὰ σημεῖα λέγω, ὥσπερ καὶ χειμῶνος.

Έλν νότου πνέουτος βορράθεν ἀστράπτη, παύεται· ἐὰν ἔωθεν ἀστράπτη εἴωθε παύεσθαι τριταῖος, οἱ δὲ ἄλλοι πεμπταῖοι ἑβδομαῖοι ἐνναταῖοι, οἱ δὲ δειλινοὶ ταχὺ παύονται. οἱ βορέαι παύονται ώς ἐπὶ τὸ πολὺ ἐν περιτταῖς οἱ δὲ νότοι ἐν ἀρτίαις. ἄνεμοι αἴρονται ἀμ' ἡλίφ ἀνατέλλοντι καὶ σελήνη. ἐὰν ἀνατέλλων ὁ ἥλιος καὶ σελήνη παύσωσιν, ἐπιτείνει τὰ πνεύματα· χρονιώτερα δὲ καὶ ἰσχυρότερα τὰ πνεύματα γίνεται τὰ ἡμέρας ἡ νύκτωρ

ἀρχόμενα.

34 Εάν ἐτησίαι πολὺν χρόνον πνεύσωσι καὶ μετόπωρον γένηται ἀνεμῶδες, ὁ χειμῶν νήνεμος γίνεται, ἐὰν δ' ἐναντίως, καὶ ὁ χειμῶν ἐναντίος.

⁸ αν. Sc. εΙναι, which perhaps should be added.

 ¹ κηλάδες, i.e. a 'mackerel sky' (?) The word seems to occur nowhere else except in Hesych, who renders ἄνυδρος: derivation obscure. It should probably be read in § 51 for κοιλάδες.
 2 Plin. 18. 354.

CONCERNING WEATHER SIGNS, 31-34

the sun is first seen, there will be wind from that quarter. Light¹ clouds in summer-time indicate wind.

If lightning comes from all sides, it indicates rain, and from any quarter from which the flashes come in quick succession there will be wind. In summer² from whatever quarter lightning and thunder come, there will be violent winds: if the flashes are brilliant and startling, the wind will come sooner and be more violent; if they are of gentler character and come at longer intervals, the wind will get up gradually. In winter and autumn however the reverse happens, for the lightning causes the wind to cease: and, the more violent the lightning and thunder are, the more will the wind be reduced. In spring I consider that the indications would ³ not so invariably have the same meaning,—and this is also true of winter.

If, while a south wind is blowing, there comes lightning from the north, the wind ceases. If there is lightning at dawn, the wind generally ceases on the third day: other winds than a south wind however do not cease till the fifth seventh or ninth day, though a wind which got up in the afternoon will cease sooner. A north 4 wind generally ceases in an odd, a south wind in an even number of days. Winds get up at sunrise or moonrise. If the rising sun or moon have caused the wind to cease, presently 5 it gets up again with more force, and winds which begin to blow in the day-time last longer and are stronger than those which begin at night.

If periodic winds have been blowing for a long time, and a windy autumn follows, the winter is windless: if however the contrary happens, the character


⁴ Plin, 2, 129.

⁵ So Furl. renders: W. inserts μη after σελήνη.

πρὸς κορυφης όρους ὁπόθεν ἃν νεφέλη μηκύνηται, ταύτη ἄνεμος πνευσείται. αὶ νεφέλαι ἐκ τῶν ὅπισθεν πνευσοίζουσαι καὶ ὅπισθεν πνευσοῦνται. ᾿Αθως μέσος διεζευγμένος νότιος, καὶ ὅλως τὰ ὅρη διεζωσμένα νότια ὡς τὰ πολλά. οἱ κομῆται ἀστέρες ὡς τὰ πολλὰ πνεύματα σημαίνουσιν, ἐὰν δὲ πολλοί, καὶ αὐχμόν. μετὰ χιόνα νότος, μετὰ πάχνην βορέας εἴωθε πνεῖν. μύκητες ἐπὶ λύχνου νότιον πνεῦμα ἡ ὕδωρ σημαίνουσιν.

Αί δὲ στάσεις τῶν πνευμάτων οὕτως ἔχουσιν ώς ἐν τῷ γράμματι διώρισται. τῶν δ' ἀνέμων ἔτι πνέουσι τοῖς ἄλλοις ἐπιπίπτουσι μάλιστα ἀπαρκτίας θρακίας ἀργέστης. ὅταν δὲ μὴ ὑπ' ἀλλήλων διαλύωνται τὰ πνεύματα, ἀλλ' αὐτὰ καταμαρανθῶσι, μεταβάλλουσιν εἰς τοὺς ἐγομέ-

g. de Ventis 50; Arist. Probl. 26.3.
 g. de Ventis 50; Arist. Probl. 26.3.
 g. de Ventis 50; Arist. Probl. 26.3.
 g. de Ventis 50; Arist. Meteor. 2. 6. describes such a figure (ὑπογραφή), which may be reconstructed thus:—



¹ cf. 22. ² cf. 57.

CONCERNING WEATHER SIGNS, 34-35

of winter is also reversed. From whatever quarter cloud streams out from a mountain peak, wind will blow in the direction thus indicated. Clouds which cling to the back of the mountain will also produce wind from the back of it. If there is a girdle 1 of cloud half way up Mount Athos, and if mountains in general wear such a girdle, there will generally follow a southerly wind. Comets 2 usually indicate wind, and, if there are many of them, drought is also indicated. After snow 3 a south wind, after hoar-frost a north wind generally blows. Snuff 4 in a lamp indicates wind or rain from the south.

The points from which the winds come are as they are given in the figure.⁵ The winds which most often come on the top of other winds while these are still blowing are the north wind (aparktias),⁶ the north-north-east and the north-west. When however the winds are not dispersed by one another but die down of their own accord, they change ⁷ to the next winds on the figure, reckoning from left

Arist. does not seem to distinguish $\beta o \rho \ell as$ and $\delta \pi a \rho \kappa \tau \ell as$: his $\delta \rho a \kappa \ell as$ is T.'s $\delta \rho a \kappa \ell as$: his eight principal winds (underlined in diagram) correspond to those represented on the famous Tower of the Winds at Athens, built about two hundred years later.

⁶ cf. Arist. l.c. ⁷ Plin. 2. 128.

νους ἐπὶ δεξιά, ὥσπερ ἡ τοῦ ἡλίου ἔχει φορά. ὁ νότος ἀρχόμενος ξηρὸς τελευτῶν δὲ ὑγρός. καὶ ὁ εὖρος. ὁ δ' ἀπηλιώτης ἀπὸ ἀνατολῆς ἰσημερινῆς ὑδατώδης. διὰ λεπτῶν δὲ ἄγει τὰ ὕδατα.

'Υγροὶ δὲ μάλιστα ὅ τε καικίας καὶ λίψ· χαλαζώδης δ' ἀπαρκτίας καὶ θρακίας καὶ ἀρ-γέστης· νιφετώδης δὲ ὅ τε μέσης [ἢ βορέας] καὶ ἀπαρκτίας· καυματώδης δὲ νότος καὶ ζέφυρος καὶ εὖρος· οἱ μὲν οἰς ᾶν ἐκ πελάγους προσπίπτωσυν, οἱ δὲ οἰς ᾶν διὰ γῆς. δασύνουσι δ' οὐράνὸν νέφεσι καὶ καλύπτουσι καικίας μάλιστα εἰτα λίψ. καὶ οἱ μὲν ἄλλοι ἄνεμοι ἀφ' ἑαυτῶν τὰ νέφη ἀθοῦσι, καικίας δὲ μόνος πνέων εἰς ἑαυτόν. αἴθριοι δὲ μάλιστα θρακίας καὶ ἀργέστης καὶ τῶν λοιπῶν ἀπαρκτίας· ἐκνεφίαι δὲ μάλιστα ὅ τε ἀπαρκτίας καὶ ὁ θρακίας καὶ ὁ ἀργέστης.

87 Γίνονται δὲ ἐκνεφίαι ὅταν εἰς ἀλλήλους ἐμπίπτωσι πνέοντες μάλιστα μὲν μετοπώρου τῶν δὲ λοιπῶν ἔαρος. ἀστραπαῖος δὲ θρακίας καὶ ἀργέστης καὶ ἀπαρκτίας καὶ μέσης. ἐὰν ἐν τῆ θαλάττη πάπποι φέρωνται πολλοὶ οἱ γινόμενοι ἀπὸ τῶν ἀκανθῶν, ἄνεμον σημαίνουσιν ἔσεσθαι μέγαν. ὅθεν ἃν ἀστέρες διάττωσι πολλοί, ἄνεμον

I have bracketed ħ βορέαs as probably a gloss on ἀπαρκτίαs;
 ἡ is difficult to account for otherwise. See diagram.
 Plin. 18. 360.

CONCERNING WEATHER SIGNS, 35-37

to right according to the course of the sun. When the south wind begins to blow, it is dry, but it becomes wet before it ceases: so too does the southeast wind. The east wind, coming from the quarter where the sun rises at the equinox, is wet: but it

brings the rain in light showers.

The north-east and south-west are the wettest winds: the north the north-north-east and the northeast bring hail; snow comes with the north-northeast 1 and north. The south, the west, and the south-east winds bring heat. Some of these have their effect on places which they strike as they come from the sea, others on places which they visit as they come over land. The winds which more than any others make the sky thick with cloud and completely cover it are the north-east and the southwest, especially the former. While the other winds repel the clouds from themselves, the north-east alone attracts them as it blows. Those winds which chiefly bring a clear sky are the north-north-west and the north-west, and next after them the north. Those which most have the character of a hurricane are the north the north-north-west and the northwest.

They acquire this character when they fall upon one another as they blow, especially in autumn, but to some extent in spring. Those which are accompanied by lightning are the north-north-west the northwest the north and the north-north-east. If at sea 2 a quantity of down is seen blown along, which has come from thistles, it indicates that there will be a great wind. Wind 3 may be expected from any quarter in which a number of shooting stars are

³ cf. 13; Plin. 18. 352; Verg. Georg. 1. 365.

έντεθθεν έὰν δὲ πανταχόθεν όμοίως, πολλά πνεύματα σημαίνουσι.

Πνευμάτων μέν οὖν σημεῖα ταῦτα.

- 38 III. Χειμῶνος δὲ τάδε. ἥλιος δυόμενος εἰς μὴ καθαρόν. καὶ ὡς ἂν μερισθῆ δυόμενος, οὕτως αἰ ἡμέραι ἐπιτελοῦνται. οἶον εἰ τὸ τρίτον μέρος ἀπολειφθείη ἡ τὸ ἥμισυ. τὸ σελήνιον ἐὰν ὀρθὸν ἡ μέχρι τετράδος καὶ εἰ εὔκυκλον, χειμάσει μέχρι διχοτόμου· γέρανοι ἐὰν πρωὶ πέτωνται καὶ ἀθρόοι, πρωὶ χειμάσει, ἐὰν δὲ ὀψὲ καὶ πολὺν χρόνον, ὀψὲ χειμάσει. καὶ ἐὰν ὑποστραφῶσι πετόμενοι, χειμῶνα σημαίνουσι.
- 39 Χηνες βοώντες μάλλον η περί σίτον μαχόμενοι χειμέριον. σπίνος στρουθός σπίζων εωθεν χειμέριον. όρχιλος [ώς] εἰσιὼν καὶ εἰσδυόμενος εἰς όπὰς χειμῶνα σημαίνουσι καὶ ἐριθεὺς ὡσαύτως. κορώνη ἐὰν ταχὺ δὶς κρώζη καὶ τρίτον, χειμερία. καὶ κορώνη καὶ κόραξ καὶ κολοιὸς ὀψὲ ἄδοντες χειμέριοι. στρουθὸς ἐὰν λευκὸς ἡ χελιδὼν ἡ ἄλλο τι τῶν μὴ εἰωθότων λευκῶν, χειμῶνα μέγαν σημαίνουσιν, ὅσπερ καί, μέλανες ἐὰν πολλοὶ φανῶσιν, ὕδωρ.
- 40 Καὶ ἐὰν ἐκ πελάγους ὅρνιθες φεύγωσι, χειμῶνα σημαίνουσι. καὶ σπίνος ἐν οἰκίᾳ οἰκουμένη φθεγγόμενος χειμέριον. ὅσα ὕδωρ σημαίνει, χειμῶνα ἄγει, ἐὰν μὴ ΰδωρ, χιόνα καὶ χειμῶνα.

¹ i.e. and the succeeding day will be more or less stormy in proportion. $\frac{\lambda\pi\omega\lambda\epsilon\iota\mu\theta\epsilon(\eta)}{2}$? $\frac{\lambda\pi\omega\lambda\eta\mu\theta\epsilon(\eta)}{2}$ i.e. it is possible, more or less, to see the whole circle.

⁴¹⁸

CONCERNING WEATHER SIGNS, 37-40

seen. If these appear in every quarter alike, it indicates many winds.

Such then are the signs of winds.

III. The following are signs of storm. The sun becoming obscured as it sinks indicates storm. And, according as its orb is divided as it sets, so the succeeding days turn out; for instance, a third or a half of the orb may remain visible. If the horns 2 of the moon point straight up till the fourth day, and if it rounds to a circle, it will be stormy till the middle of the month. If cranes fly early and in flocks, it will be an early winter 3; if they fly late and for a long time, it will be a late winter; and, if they wheel as they fly, it indicates stormy weather.

⁴ It is a sign of storm when geese make more clamour than usual or fight for their food; so too is it when a sparrow or chaffinch twitters at dawn. It indicates a storm when the goldcrest⁵ goes into holes and hides itself; so also when the redbreast does the same. It is a sign of storm when the crow caws twice in quick succession and then a third time; also when the crow or raven or jackdaw makes its call late. It is a sign of a great storm when a white sparrow or swallow is seen, or a white specimen of any other bird which is not usually white, even as the appearance of a large number of such birds of a dark colour signifies rain.

It is also an indication of storm when birds flee from the sea. A chaffinch uttering its note in an inhabited house is a sign of storm. All the signs which indicate rain bring stormy weather, that is to say, snow and storm, if not rain. If the raven utters

³ So Arat. 343 f. interprets. ⁴ Plin. 18. 363. ⁵ ‰ bracketed by Sch.

κόραξ φωνὰς πολλὰς μεταβάλλων χειμώνος χειμέριον. κολοιοὶ ἐκ τοῦ νότου πετόμενοι καὶ τευθίδες χειμέριαι. φωνὴ ἐν λιμένι ἀποψοφοῦσα καὶ πολύπλοκον ἤχοῦσα χειμέριον. καὶ οἱ πνεύμονες οἱ θαλάττιοι ἐὰν πολλοὶ φαίνωνται ἐν τῷ πελάγει, χειμερινοῦ ἔτους σημεῖον. πρόβατα ἐὰν πρωὶ ὀχεύηται, πρώϊον χειμῶνα σημαίνουσι.

Μετοπώρω ἐὰν πρόβατα ἡ βόες ὀρύττωσι καὶ κοιμῶνται ἀθρόοι πρὸς ἀλλήλους ἔχοντες τὰς κεφαλάς, τὸν χειμῶνα χειμέριον σημαίνει. ἐν δὲ τῷ Πόντω φασίν, ὅταν ᾿Αρκτοῦρος ἀνατείλη θᾶττον, ἐναντίους τῷ βορρῷ νέμεσθαι. βόες μᾶλλον ἐσθίοντες τοῦ εἰωθότος καὶ ἐπὶ τὸ δεξιὸν κατακλινόμενοι χειμέριον. καὶ ὧτα κρούων ὄνος χειμέριον καὶ μαχόμενα πρόβατα καὶ ὅρνιθες περὶ σίτου παρὰ τὸ ἔθος προπαρασκευάζονται γάρ καὶ μῦες τρίζοντες καὶ ὀρχούμενοι χειμέριον.

42 Καὶ κύων τοῖς ποσὶν ὀρύττουσα καὶ ὁλολυγὼν ἄδουσα μόνη ἀκρωρίας χειμέριον. γῆς ἔντερα πολλὰ φαινόμενα χειμῶνα σημαίνει. καὶ ἐὰν πῦρ μὴ θέλη ἄπτεσθαι, χειμέριον καὶ ἐὰν λύχνος ἄπτεσθαι μὴ ἐθέλῃ, χειμῶνα σημαίνει καὶ τέφρα πηγνυμένη νιφετόν. λύχνος εὐδίας ἡσυχαῖος καιόμενος χειμῶνα σημαίνει καὶ ἐὰν χειμῶνος ὄντος μύκαι μέλαιναι ἐπιγίνωνται, χειμῶνα σημαίνει καὶ ἐὰν ὥσπερ κέγχροις πολλοῖς κατά-

² cf. 21, 29. πολύπλοκον is Furlanus' conj. for Vulg. πολύποδον.

³ πνεύμονες. Plin. 18. 359. pulmones: cf. 9. 154.

 $^{^{1}}$ $\tau \epsilon \nu \theta \ell \delta \epsilon s$. The word is perhaps corrupt and conceals the name of a bird.

CONCERNING WEATHER SIGNS, 40-42

a great variety of sounds in winter, it is a sign of storm. Jackdaws flying from the south are a sign of storm, and so are cuttle-fish.1 It is a sign of storm when a loud 2 voice is heard in harbour, which is reechoed many times. It is a sign of a stormy season when a number of jelly-fish 3 appear in the sea. indicates an early winter when the breeding season

of sheep begins early.

If in autumn sheep or oxen dig holes and lie keeping their heads close to one another, it indicates a severe winter. They say that in Pontus when Arcturus rises, (the cattle 4) face northwards as they graze. It is a sign of storm when cattle eat more than usual and lie down on their right sides.⁵ So is it when the ass shakes 6 his ears, or when sheep or birds fight for their food more than usual, since they are then trying to secure a store against bad weather: also when mice squeak and dance.

A bitch digging holes with her paws and a treefrog croaking alone at early dawn 7 are signs of storm: it indicates storm when a number of the worms 8 called 'the earth's entrails' appear. It is a sign of storm if the fire refuses to catch, or if a lamp refuses to light: while, if much ash is formed, it is a sign of snow. If a lamp burns steadily in fine weather, it is a sign of storm: so is it if in wintertime dark snuff of forms: if it is, as it were, full of numerous millet-seeds, there will be stormy weather;

⁴ θᾶττον is clearly corrupt, and words indicating what the sign portends are missing. ⁵ cf. 54. ⁶ δτα κρούων doubtful. Sch. suggests οδδαs for δτα.

⁷ ἀκρωρίας. cf. 21.

⁸ γης έντερα. So Arat. 225 explains. One might guess 'worm-casts.'

⁹ cf. 14, 25, 34, 54.

πλεως ή, χειμερίσει καὶ ἐὰν κύκλφ περὶ τὸ

λαμπρου ώσιν ευδίας ούσης, χιονικόν.

Υ΄Η τοῦ ὄνου Φάτνη εἰ συνίσταται καὶ ζοφερὰ γίνεται, χειμῶνα σημαίνει. καὶ ἐὰν ἀστραπὴ λαμπρὰ μὴ ἐν τῷ αὐτῷ μένη, χειμέριον. ἐπὶ Πλειάδι δυομένη ἐὰν λάμψη κατὰ Πάρνηθα καὶ Βρίληττον καὶ "Υμηττον, ἐὰν μὲν ἄπαντα καταλάμψη, μέγαν χειμῶνα σημαίνει, ἐὰν δὲ τὰ δύο, ἐλάττω, ἐὰν δὲ Πάρνηθα μόνον, εὐδιεινόν· καὶ ἐὰν χειμῶνος ὄντος νεφέλη μακρὰ ἐπὶ τὸν "Υμηττον ἢ, χειμῶνος ἐπίτασιν σημαίνει. *Αθως καὶ 'Ολυμπος καὶ ὅλως ὀρέων κορυφαὶ κατεχόμεναι ὑπὸ νεφελῶν χειμέριον. ἐὰν εὐδίας γινομένης νεφέλιον φαίνηται ἐν τῷ ἀέρι παρατεταμένον καὶ τιλλόμενον, οὕπω παύεται ὁ γειμών.

Έαν το μετόπωρον εὐδιείνου παρά το εἰκος γένηται, το ἔαρ γίνεται ψυχρον ὡς τὰ πολλά. ἐὰν πρωὶ χειμάζειν ἄρξηται, πρωὶ παύεται καὶ ἔαρ καλόν, ἐὰν δὲ τοὐναντίον, καὶ ἔαρ ὅψιον ἔσται. ἐὰν χειμὼν ὑέτιος, τὸ ἔαρ αὐχμηρόν, ἐὰν δ΄ αὐχμηρὸς ὁ χειμών, τὸ ἔαρ καλόν. ἐὰν ἡ ὅπωρα γίνηται ἐπιεικής, τὰ πολλὰ γίνεται τοῖς προβάτοις λιμός. ἐὰν τὸ ἔαρ καὶ τὸ θέρος ψυχρὰ γίνηται, ἤ τε ὅπωρα γίνεται καὶ <τὸ> μετόπωρον

πνιγηρον και ούκ άνεμωδες.

46 Οἱ πρῖνοι ἐὰν εὐκαρπῶσι, χειμῶνες πολλοὶ σφόδρα γίνονται. ἐὰν ἐπὶ κορυφῆς ὅρους νέφος ὀρθὸν στῆ, χειμῶνα σημαίνει, ὅθεν καὶ ᾿Αρχίλοχος ἐποίησε " Γλαῦχ' ὅρα· βαθὺς γὰρ ἤδη κύμασιν

δνου φάτνη. cf. 23, 51. See LS, s.v. δνος; Theocr. 22.
 Plin. 18. 353, sunt in signo Cancri duae stellae parvae

CONCERNING WEATHER SIGNS, 42-45

and if these in fine weather appear in a circle round

the flame, it is a sign of snow.

If the 'Ass's Manger 1' shrinks in size and becomes dark, it is a sign of storm; also if there is vivid lightning which does not remain in the same quarter. If at the setting of the Pleiad there is lightning over Parnes Brilessus and Hymettus—when it appears over all three mountains, it indicates a great storm; when over the two lesser heights, a less violent storm; when over Parnes alone, fine weather. Again, if during a storm a long cloud stretches over Hymettus, it signifies that the storm will increase in force. It is a sign of storm when Athos Olympus and mountain-peaks in general are covered with clouds. If during fine weather a cloud appears in the sky stretching a long way and torn to shreds, stormy weather will continue.

If the autumn is unusually fine, the succeeding spring is generally cold. If winter begins early, it ends early and there is a fair spring; if the reverse, spring will also be late. If the winter is wet, the spring will be dry, if the winter is dry, the spring will be fair. If the late summer is satisfactory, the sheep will generally suffer from hunger. If the spring and summer are cold, the late summer and

autumn 2 will be stifling hot and windless.

If the kermes-oak struits well, there follows a long succession of storms. If a cloud stands upright on a mountain-peak, it indicates storm; whence Archilochus' lines "Mark you, Glaucus; deep ocean

aselli appellatae, exiguum inter illas spatium obtinente nubecula, quam praesepia appellant. 2 75 add. Sch. 3 cf. 49.

A comparison of war to stormy weather. Quoted also by Plut. de Superstitione, 72, and by Heraclides, Allegoriae

Homericae, 4. In both citations the Greek is corrupt.

ταράσσεται Πόντος ἀμφὶ δ' ἄκρα <Γυρῶν> ὀρθὸν ίσταται νέφος Σήμα χειμώνος." ἐὰν δ' ὁμόχρων ή υμένι λευκφ, χειμέριον. ὅταν ἐστώτων νεφῶν ἔτερα ἐπιφέρηται τὰ δ΄ ἠρεμῆ, χειμέρια.

< Ο ήλιος> εαν χειμώνος διαλάμψας πάλιν ἀποκρυφθη καὶ τοῦτο ποιήση δὶς ἡ τρίς, ἡμέρα χειμέριος δίεισιν. ὁ τοῦ Ἑρμοῦ ἀστηρ χειμῶνος μὲν φαινόμενος ψύχη σημαίνει θέρους δὲ καῦμα. όταν μέλιτται μη ἀποπέτωνται μακράν άλλ αὐτοῦ ἐν τῆ εὐδία πέτωνται, χειμῶνα ἐσόμενον σημαίνει. λύκος ὧρυόμενος χειμῶνα σημαίνει διὰ τριῶν ἡμερῶν. λύκος ὅταν πρὸς τὰ ἔργα όρμα ή είσω χειμώνος ώρα, χειμώνα σημαίνει *คั*บθีบร.

*Εστι δὲ σημεῖον χειμώνων μεγάλων καὶ ὅμβρων και δταν γένωνται έν τῷ μετοπώρφ πολλοί σφήκες, και όταν όρνιθες λευκοί προς τα έργάσιμα πλησιάζωσι, καὶ όλως τὰ ἄγρια θηρία ἐὰν πρὸς τὰ ἐργάσιμα, βόρειον καὶ χειμῶνος μέγεθος ση-μαίνει. τῆς Πάρνηθος ἐὰν τὰ πρὸς ζέφυρον άνεμον και τὰ πρὸς Φύλης φράττηται νέφεσι βορείων όντων, χειμέριον τὸ σημείον.

48 "Όταν πνίγη γίνηται ἰσχυρά, ὡς τὰ πολλὰ ἀνταποδίδωσι καὶ γίνεται χειμὼν ἰσχυρός. ἐὰν ύδατα ἐαρινὰ πολλὰ γένηται, καύματα ἰσχυρὰ έν τοις πεδινοις και κοίλοις γίνεται. δει οθν την άρχην όραν. ἐὰν τὸ μετόπωρον εὐδιεινον γίνηται

¹ Γυρών. γυροῦν W. Heraclides gives γυρεδν, Plut. γυρεῦον; but one MS. of Plut. gives γύρων with a marginal gloss 'sc. $\pi \epsilon \tau \rho \hat{\omega} \nu$, which suggests that the word is a proper name. Od. 4. 500 mentions the Γύραι (i.e. the 'round-backed rocks')

CONCERNING WEATHER SIGNS, 45-48

is now stirred up with waves, and about the heights of the Gyrae ¹ there rises a cloud erect, the sign of storm." If the clouds are of uniform colour, like ² a white membrane, it is a sign of storm. When, as some clouds are motionless, others move towards them while they remain at rest, it is a sign of storm.

If the sun in winter after gleaming out is again obscured, and this is repeated two or three times, it will be stormy all day. If the star Hermes appears in winter, it indicates cold, if in summer, heat. When in fine weather bees do not fly ⁸ long distances, but fly about where they are, it indicates that there will be a storm. The howling of a wolf indicates a storm within three days. When a wolf approaches or enters cultivated ground in the season of winter, it indicates that a storm will come immediately.

It is also a sign of great storms and heavy rain when many wasps appear in autumn, or when white birds approach cultivated lands; and in general when wild creatures approach such lands, it indicates a north wind and a severe storm. If the western side of Parnes and the side towards Phyle are blocked with clouds during a north wind, it is a sign of storm.

When there is severe heat, generally there is compensation and a severe winter follows. If there is much rain in spring, it is followed by severe heat in low-lying districts and valleys; so that one should mark how the season begins. If the autumn is where Aias Oileus perished. The word is missing in the MSS.

2 δμοῖον has perhaps dropped out after δμόχρων ¾; the adjective seems to agree with νέφον.

s cf. Arist. H.A. 9. 40 ad fin.

⁴ Plin. 18. 363: presumably gulls, etc.

σφόδρα, τὸ ἔαρ ὡς τὰ πολλὰ γίνεται ψυχρόν ἐὰν δὲ τὸ ἔαρ ὄψιον γένηται καὶ ψυχρόν, ἡ ὅπωρα ὀψία γίνεται καὶ <τὸ> μετόπωρον ὡς τὰ πολλὰ πνιγηρόν.

- 49 Οἱ πρίνοι ὅταν εὐκαρπῶσι σφόδρα, ὡς μὲν τὰ πολλὰ χειμῶνα ἰσχυρὸν σημαίνουσιν, ἐνίστε δὲ καὶ αὐχμούς φασι γίνεσθαι. καὶ ἐάν τις σπάλακα λαβῶν ὑποπάσας ἄργιλον εἰς πιθάκνην θῆ, σημαίνει ταῖς φωναῖς αἶς ἀφίησιν ἄνεμον καὶ εὐδίαν. καὶ τὸ πανταχοῦ δὲ λεγόμενον σημεῖον δημόσιον χειμέριον, ὅταν μῦες περὶ φορυτοῦ μάχωνται καὶ φέρωσιν.
- 50 IV. Εὐδίας δὲ σημεῖα τάδε. ἥλιος μὲν ἀνιὼν λαμπρὸς καὶ μὴ καυματίας καὶ μὴ ἔχων σημεῖον μηδὲν ἐν ἑαυτῷ εὐδίαν σημαίνει. ὡς δ' αὕτως σελήνη πανσελήνῳ. καὶ δυόμενος ἥλιος χειμῶνος εἰς καθαρὸν εὐδιεινός, ἐὰν μὴ ταῖς προτέραις ἡμέραις εἰς μὴ καθαρὸν δεδυκὼς ἢ ἐξ εὐδιῶν οὕτω δὲ ἄδηλον. καὶ ἐὰν χειμάζοντος ἡ δύσις γένηται εἰς καθαρόν, εὐδιεινόν καὶ ἐὰν δύνων χειμῶνος ὡχρὸς ἢ, εὐδίαν σημαίνει.
- 61 Καὶ ὁ μεὶς ἐὰν τριταῖος ὢν λαμπρὸς ἢ, εὐδιεινόν. καὶ ἡ τοῦ ὄνου Φάτνη ὅτε ἂν καθαρὰ καὶ λαμπρὰ φαίνηται, εὐδιεινόν. ἄλως δὲ ἐὰν ὁμαλῶς

³ σπάλακα Vulg.; σπάκα Bas. Ald.; σκολόπακα (woodcock?) conj. Furl.

¹ τδ add. Sch. ² cf. 45.

⁴ i.e. (reading σκολόπακα) for the bird to find worms in with its long beak (Sch.). It is hard to say, without illus-

CONCERNING WEATHER SIGNS, 48-51

exceedingly fine, generally the spring is cold: if the spring is late and cold, the summer goes on late and

the 1 autumn is usually scorching hot.

When the kermes-oak ² fruits exceedingly well, it generally indicates a severe winter, and sometimes they say that this sign is followed by droughts. If one takes a mole³ and puts it in a tub, the bottom⁴ of which has been covered with clay, it indicates by the sounds which it utters wind or fine weather. There is also the sign of storm which is popularly recognized everywhere, namely when mice fight for the possession of chaff and carry it about.

The signs of fair weather.

IV. The following are signs of fair weather. ⁵ If the sun rises brilliant but without scorching heat and without showing any special sign in his orb, it indicates fair weather. The same may be said of the moon when it is full. If in winter that part of the sky into which the sun goes down is clear, it is a sign of fair weather, unless on the preceding days that part has not been clear, though it was clear above the horizon: in that case the prospect is uncertain. It is also a sign of fair weather, if during stormy conditions that part of the sky into which the sun sets is clear; and also if, in winter at the time of setting, the sun has a pale colour.

Again, it indicates fair weather if the outline of the moon on the third day is bright; also if the 'Ass's 6 Manger' is clear and bright. If the halo 7 forms and disappears evenly, it is a sign of fair

tration, which is the more convincing of the creatures suggested.

⁵ Plin. 18. 342.

⁶ cf. 23, 43.

⁷ cf. 22, 31; Plin. 18. 345; Arist. Meteor. 3. 3.

παγή καὶ μαρανθή, εὐδίαν σημαίνει. αἱ κηλάδες νεφέλαι χειμῶνος εὐδιειναί. "Ολυμπος δὲ καὶ "Αθως καὶ ὅλως τὰ ὅρη τὰ σημαντικὰ ὅταν τὰς κορυφὰς καθαρὰς ἔχωσιν, εὐδίαν σημαίνει. καὶ ὅταν τὰ νέφη πρὸς τὴν θάλασσαν αὐτὴν παραζωννύη, εὐδιεινόν. καὶ ὅταν ὕσαντος πρὸς δυσμὰς χαλκῶδες τὰ νέφη χρῶμα ἔχη εὐδία γὰρ ὡς τὰ

πολλά τῆ ὑστεραία.

Οταν δε όμίχλη γένηται, ὕδωρ οὐ γίνεται ἡ ἔλαττον. ὅταν γέρανοι πέτωνται καὶ μὴ ἀνακάμπτωσιν, εὐδίαν σημαίνει οὐ γὰρ πέτονται πρὶν ἡ ἀν πετόμενοι καθαρὰ ἴδωσι. γλαῦξ ἡσυχαῖον φθεγγομένη ἐν χειμῶνι εὐδίαν προσημαίνει καὶ νύκτωρ χειμῶνος ἡσυχαῖον ἄδουσα. θαλαττία δε γλαῦξ ἄδουσα χειμῶνος μὲν εὐδίαν σημαίνει, εὐδίας δὲ χειμῶνα. καὶ κόραξ δὲ μόνος μὲν ἡσυχαῖον κράζων, καὶ ἐὰν τρὶς κράξη μετὰ τοῦτο πολλάκις κράξη, εὐδιεινός. . . .

53 Και κορώνη εωθεν εὐθὺς ἐὰν κράξη τρίς, εὐδίαν σημαίνει, και ἐσπέρας χειμῶνος ἡσυχαίον ἄδουσα. και ὀρχιλος ἐξ ὀπῆς ἐκπετόμενος και ἐξ ἑρκίων και ἐξ οἰκίας ἔξωθεν εὐδίαν σημαίνει. και ἐὰν χειμῶνος βορεύοντος βορρᾶθεν ὑπόλαμψις γένηται λευκή, νοτόθεν δὲ ἐναντία τεταγμένη ἢ νεφέλη ὀγκώδης, ὡς ἐπὶ τὸ πολὺ εἰς εὐδίαν σημαίνει μεταβολήν. και ὅταν βορέας νεφέλας πολλὰς κινῆ

έκπνέων μέγας, εὐδίαν σημαίνει.

428

Digitized by GOOGLE

¹ κηλάδες I conj. cf. 31, to which this statement answers. κοιλάδες MSS.

² Plin. 18. 356. ³ Plin. 18. 357. cf. Verg. Georg. 1. 401.

⁴ Plin. 18. 362.

⁵ ἐν χειμῶνι. ? 'in winter.' The same ambiguity occurs in many places: the sense seems fixed here by the next sentence.

CONCERNING WEATHER SIGNS, 51-53

weather. Light 1 clouds in winter are a sign of fine weather. It is a sign of fine weather when Olympus Athos and in general the mountains which give signs have their tops 2 clear: so too is it, when clouds encompass them at the sea-level. 3 Also when after rain the clouds have a bronze colour towards sundown: in that case there will generally be fine

weather the next day.

When there is mist, little or no rain follows. When cranes 4 take flight and do not come back, it is a sign of fair weather: for they do not do so till they see a clear sky before them as they fly. It is a sign of fair weather when during a storm 5 an owl makes a low hoot, or at night during a storm it utters a low sound. If the sea-owl utters its note during a storm, it indicates fair weather, if during fair weather, it indicates a storm. It is a sign of fair weather if a solitary raven makes a low croak, and, after croaking three times, repeats the sound again and again. . . .6

If the crow caws thrice directly the dawn appears, it indicates fair weather, as also if it makes a low note in the evening during a storm. It is a sign of fair weather if a goldcrest flies out abroad from a hole or from a hedge or from its nest. Again, if during a storm from the north there is a white gleam from that quarter, while in the south a solid mass of cloud has formed, it generally signifies a change to fair weather. Again when the north wind (Boreas) as it begins to blow violently stirs up a number of clouds, it indicates fair weather.

⁶ I have marked a lacuna: the answer to $\mu \hat{\epsilon} \nu$ is missing, presumably a statement about the significance of more than one raven. cf. Verg. Georg. 1. 410.

54 Πρόβατα όψε όχευόμενα εὐδιεινον ἀποτελοῦσι τὸ σημείον. καὶ βοῦς ἐπὶ τὸ ἀρίστερον ἰσχίον κατακλινόμενος εύδίαν σημαίνει καὶ κύων ώσαύτως έπι δεξιον δε χειμώνα. τέττιγες πολλοί γινόμενοι νοσώδες τὸ έτος σημαίνουσι. χειμώνος καιόμενος ήσυχαίος εὐδίαν σημαίνει. καὶ ἐὰν ἐπ' ἄκρω οίον κέγχρους ἔχη λαμπράς. καὶ ἐὰν ἐν κύκλω τὴν μύξαν περιγράφη λαμπρὰ γραμμή.

Ο της σχίνου καρπός σημαίνει τούς άρότους. έχει δὲ τρία μέρη καὶ ἔστιν ὁ πρῶτος τοῦ πρώτου αρότου σημείου, ο δεύτερος τοῦ δευτέρου, ο τρίτος τοῦ τρίτου καὶ ὡς αν τούτων ἐκβαίνη κάλλιστα καὶ γένηται άδρότατος, οῦτως ἔξει καὶ ὁ κατά

τοῦτον ἄροτος.

Λέγεται δὲ καὶ τοιάδε σημεῖα ὅλων τε τῶν ένιαυτών γίνεσθαι καλ τών μορίων. έαν άρχομένου τοῦ χειμῶνος ζόφος ή καὶ καύματα γίνηται καὶ ταῦτα ἄνευ ὕδατος ὑπ' ἀνέμων διαλυθῆ, πρὸς τὸ ἔαρ σημαίνει χάλαζαν ἐσομένην. μετά την εαρινην ισημερίαν ομίχλαι πίπτωσι, πνεύματα καὶ ἀνέμους σημαίνουσιν εἰς εβδομον μηνα άμφοτέρων άριθμουμένων. δσαι μεν άμα μηνοειδεί τη σελήνη πίπτουσιν, αύται μέν πνεύματα σημαίνουσιν είς εκείνον τὸν χρόνον, ὅσαι δ' άμφικύρτου ούσης της σελήνης ύδατα, καὶ ὅσω

¹ cf. 41. ² cf. 14, 25, 42. ³ H.P. 7. 13. 6 the same is said of σκίλλα.

CONCERNING WEATHER SIGNS, 54-56

When sheep begin to breed late, it is a sign which fulfils itself in fair weather. So is it when an ox lies ¹ on his left side, and also when a dog does the same: if they lie on the right side, it indicates storm. The appearance of a number of cicadas indicates that the season will be unhealthy. If a lamp burns quietly during a storm, it indicates fair weather. So also if it has on the surface an appearance like shining millet-seeds: ² also if a bright line surrounds the lamp-nozzle.

The fruiting of the mastich ⁸ gives signs as to the seasons of sowing: ⁴ it takes place at three several periods, which indicate respectively the time for the first the second and the third sowing: and according as one or other of these fruiting-times turns out ⁵ best and produces the most abundant fruit, so too will be the success of the corresponding time of

sowing.

Miscellaneous signs.

The following signs are said to affect either the whole year or whole periods 6 of it. If at the beginning of winter there is dull weather followed by heat, and these conditions are dispersed by wind without rain, it indicates that towards the spring there will be hail. Again, if after the spring equinox mists come down, it is an indication of breezes and winds by the seventh month, reckoning inclusively. Those mists which come down when the moon is in its first quarter indicate breezes for that period, those which come down when the moon is in its third quarter indicate rain. And the more mists

4 cf. H.P. 7. 1. 1 foll.

6 cf. 6.

⁵ ἐκβαίνη I conj.: cf. H.P. 7. 13. 6; κλίνη MSS.

ầν μᾶλλον ἐφ' ἐκατέρφ τῷ σχήματι ὁμίχλαι πίπτωσι, μᾶλλον τὰ εἰρημένα σημαίνει.

Σημαίνει δὲ καὶ τὰ πνεύματα ἄμα ταῖς ὁμίχλαις έπιπιπτούσαις γινόμενα καλ έαν μεν άπ' γους και μεσημβρίας γίνηται τὰ πνεύματα, ὕδατα σημαίνει· ἐὰν δ' ἀφ' ἐσπέρας και ἀπὸ τῆς ἄρκτου πνεύματα καὶ ψύχη. οθς δὲ κομήτας Αἰγύπτιοι λέγουσιν οὐ μόνον τὰ προειρημένα σημαίνουσιν όταν φαίνωνται άλλὰ καὶ ψύχη ἐπὶ δὲ τοῖς άστροις είωθεν ώς έπι τὸ πολύ σημαίνειν και ταις ισημερίαις και τροπαίς, οὐκ ἐπ' αὐταίς ἀλλ' ἡ πρὸ αὐτῶν ἡ ὕστερον μικρῶ.

 ¹ cf. 34; Arist. Meteor. 1. 6.
 ² Text seems doubtful, as cold weather was included above.

CONCERNING WEATHER SIGNS, 56-57

come down when the moon is assuming either shape, the more certainly is the result just mentioned indicated.

Also the winds which accompany the falling of the mists are significant: if the breezes come from the east or south, rain is indicated; if from the west or north, breezes and cold weather. And the stars which the Egyptians ¹ call 'comets' indicate not only the conditions just mentioned but also cold ² weather. ³ In the case of the rising of the stars the indication, as in the case also of the equinoxes and solstices, is given not at the actual time but a little earlier or later.

3 The text of this sentence can hardly be sound. σημαίνειν has no subject and ταῖε ἰσημερίαιε καὶ τροπαῖε no construction.

433

FF

VOL. II.

NOTE TO THE INDEX OF PLANTS

SPRENGEL made the first comprehensive attempt to determine in modern nomenclature the plants mentioned by Theophrastus: Wimmer gives the result in the Introduction to his 1842 edition. Sprengel adopted the most probable identifications of earlier botanists, supplemented by his own conjectures and Sibthorp's exploration of the Greek flora. The ambitious but uncritical Conspectus Florae Classicae of Frass did not add much to our knowledge, which throughout had been vitiated by failure to recognise the fact that the Mediterranean flora differed from that of Western and Central Europe. Halacsy's Conspectus Florae Graecae now gives us a scientific enumeration of the native plants of Greece; a Greek plant-name can be wedded to a plant which at any rate is Greek. Incidentally much has been cleared up by special research at the hands of De Candolle, Hanbury, Yule, Schweinfurth, Bretzl, and others.

The identifications in the following Index are drawn from various sources; for their selection in view of the botanical data available I am indebted to Sir William Thiselton-Dyer. A considerable number may be accepted as certain, many are

probable, some no more than possible.

comp.=compared.
= denotes a synonym. Where a reference is added (see e.g. ἀτρακτυλίς), it indicates that Theophrastus himself states that the names are synonymous.

άβρότονον, southernwood, Artemisia arborescens

1. 9. 4. evergreen; 6. 1. 1. in list of under-shrubs; 6. 3. 6. an unnamed plant comp.: see App. (23): 6. 7. 3. propagation; 6. 7. 4. much seed: roots described.

άγνος (= olσος), chaste-tree, Vitex Agnus-castus

- 1. 3. 2. a shrub which becomes tree-like; 1. 14. 2. bears fruit at the top; 3. 12. 1. growth of κράνεια comp.; 3. 12. 2. roots of θηλυκράνεια comp.; 4. 10. 2. ἐλαίαγνος comp.; 9. 5. 1. size of κινάμωνον and κασία comp.
- αγριέλαιος (? = κότινος), wild olive, Olea Oleaster
- 2. 2. 5. comes from seed of ἐλάα.
 ἄγρωστις, dog's tooth grass, Cynodon Dactylon
 - 6. 7. root jointed; 1. 6. 10. roots large and numerous;
 2. 2. 1. propagation; 4. 6. 6. φῦκος (6) comp.; 4. 10. 5-6 root described; 4.11.13. an unnamed form of κάλαμος comp.; root of κ. ὁ Ἰνδικός comp.; 9. 13. 6. habit of ἐρευβέδανον comp.

of ἐρευθέδανον comp. ἄγχουσα, alkanet, Anchusa tinctoria 7. 8. 3. leaves 'on the ground': 7. 9. 3. roots red.

άδίαντον, maiden-hair, Adiantum Capillus-Veneris, etc.

7. 10. 5. evergreen; 7. 14. 1. leaf cannot be wetted: two kinds

(see below): medicinal use: grows in damp places.

άδίαντον τὸ λευκόν (= τριχομανές 7. 14. 1), English maiden-hair, Asplenium Trichomanes

14. 1. described by comparison with ά. το μέλαν: medicinal use: likes shady places.

αδίαντον το μέλαν, maiden-hair, Adiantum Capillus Veneris

7. 14. 1. comp. with a. το λευκόν.

aδράφαξυς, orach, Atriplex rosea

1. 14. 2. bears fruit both on top
and at sides; 3. 10. 5. seeds of
φίλυρα comp.; 7. 1. 2-3. time
of sowing and of germination;
7. 2. 6. root described; 7. 2. 7-8.
root of βλίτον comp.; 7. 2. 8.
root: 7. 3. 2. seeds; 7. 3. 4. seed
borne both at top and at side;
7. 4. 1. only one kind; 7. 5. 5.
seed does not keep well.

άείζωον, house-leek, Sempervivum tectorum

1. 10. 4. leaves fleshy; 7. 15. 2. always moist and green: habitat. άθραγένη, traveller's joy, Clematis Vitalba

5. 9. 6. wood makes good firesticks: described; 5. 9. 7. the stationary piece should be made of this or κιττός.

aiγειροs, black poplar, Populus nigra
1. 2. 7. bark; 1. 5. 2. bark fleshy;
2. 2. 10. Cretan form bears fruit;
3. 1. 1. propagation; 3. 3. 1. tree
of mountain and plain; 3. 3. 4 a
question if it bears fruit; etc.;

3. 4. 2. time of budding; 3. 6. 1. quick growing; 3.14.2, described; 4. 1. 1. likes wet ground; 4. 7. 4. size of unnamed Arabian tree, see App. (12a), comp.; 4. 13. 2. shorter-lived by water; 5. 9. 4. wood makes an evil smoke when burnt for charcoal.

αἰγίλωψ (1) (= ἄσπρις), Turkey oak,

Quercus Cerris

3. 8. 2. one of the five (Idaean) kinds of oak: fruit; 3 8. 4. habit and timber; 3. 8. 6.

galls: φάσκος (q.v.).

aiγίλωψ (2) (grass), Aegilops ovata 7. 13. 5. seed sometimes takes two years to germinate: 8. 7. 1. comp. with aipa; 8. 8. 3. grows specially among κριθαί; 8. 9. 2. like a wild plant; 8.9.3. greatly exhausts the soil; 8. 11. 8-9. peculiarities about seed.

αίμόδωρον, broom-rape, Orobanche

criienta

8.8.5. parasitic on Bounépas (only):

described.

aipa, darnel, Lolium temulentum 1.5.2. 'bark 'in one layer; 2.4.1. πυρός turns into à.; 4. 4. 10. πρός turns into à.; 4. 4. 10. πρύζον comp.; 8. 4. 6. does not infest certain kinds of πυρός: contrasted with μελάμπυρον; 8. 7. 1. κριθή and especially πυρός said to change into à. under certain conditions: described: λίνον also said to change into \dot{a} : comp. with $ai\gamma(\lambda\omega\psi$ (2); 8. 8. 3. produced possibly by degeneration of κριθή and πυρός, or else specially affects such crops; 8. 9. 3. alto-

gether a wild plant.
ἀκαλύφη, nettle, Urtica urens
7. 7. 2. a λάχανον; needs cooking.
ἄκανθα (1) ἡ Λίγυπτία, acacia, Acacia arabica (and albida)

4. 2. 1. peculiar to Egypt; 4. 2. 8. described: two kinds (ή λευκή and ἡ μέλαινα) distinguished (see below); 9. 1. 2. sap 9. 1. 2. sap gummy.

ακανθα (ή Αίγυπτία) ή λευκή, acacla, Acacia albida

4. 2. 3. distinguished from a. ή μέλαινα.

ακανθα (ἡ Αἰγυπτία) ἡ μέλαινα, acacia. Acacia arabica

4. 2. 8. distinguished from a. n λευκή.

ἄκανθα (2) ή ἀκανώδης (see 4.10.6.n.). corn-thistle, Carduus arvensis 4. 10. 6. root etc. described

ἄκανθα (3) ἡ διψάς, Acacia tortilis

4. 7. 1. the only tree which grows on part of the 'Red Sea' coast.

ακανθα (4) ή Ἰνδική (see App. (9)), Balsamodendron Mukul

9. 1. 2. sap gummy: gum like σμύρνα,

ακανθα (5) ή λευκή 'Ηρακλέους (= aκaνθa (6)), Euphorbia antiquorum

4. 4. 12. described: uses of wood. äκανθα (6) (peculiar to Gedrosia). = aκavθa (5), Euphorbia anti-

auorum 4. 4. 13. described: has a blinding

juice. ακανθα (7) τις, gum arabic. Acantha arabica

9. 18. 1. said to have the property of thickening water.

ἄκανθα (?) (8) (= ἄκανος = ἰξία (2) iξίνη = χαμαιλέων ὁ λευκός
 12. 1.), pine-thistle, Atractylis gummifera.

 \tilde{a} κανος $(=\tilde{a}$ κανθα $(8)=\tilde{i}$ ξία $(2)=\tilde{i}$ ξίνη= χαμαιλέων ὁ λευκός), pine-thistle.

Atractylis gummifera

1. 10. 6. spinous-leaved; 1. 13. 3. flower attached above each seed; 6. 1. 3. has spines on the leaves: a wild under-shrub; 6. 4. 4. many stalks and side-growths; 6. 4. 5. one form only; 4. 8. root of σόγκος contrasted: χαιαιλέων comp.; 6. 4. 11. fruit-case of κάκτος (1) comp.; 6. 6. 6. seed of ρόδον comp.; 9. 12. 1. 'head' of χαμαιλέων ὁ λευκός comp.: another name for χαμαι-λέων (?); 9. 12. 2. leaf of χαμαι-λέων (λέως comp.) λέων ο μέλας comp.

ακόνιτον (= θηλύφονον = μυόφονον = σκορπιός (3)), wolf's bane, Aconi-

tum Anthora

16. 4. localities: described: habitat: eaten by no animal; 9. 16. 5. difficulty of compounding drug: effects: has no anti-

dote: 9.16.7. use requires expert knowledge: legal restrictions: proportion between times of gathering and of administering.

акорva, Cnicus Acarna

1. 10. 6. spinous-leaved; 7. 4. 3. a 'thistle-like' plant; 6. 4. 6. described.

άκτέος (?) (= ἀκτῆ), elder, Sambucus nigra

3. 4. 2. time of budding.

ἀκτή (= ἀκτέος), elder, Sambucus nigra 1. 5. 4. wood without knots; 1. 6. 4. core fleshy: has no core, according to some: 1. 8. 1. few knots; 4. 13. 2. shorter-lived by water; 5. 3. 3. character of wood.

άλθαία (= μαλάχη ή ἀγρία 9. 15. 5.), marsh-mallow, Althaea officin-

9. 15. 5. a drug, called in Arcadia μαλάχη ἡ ἀγρία; 9.18.1. root said to thicken water: described: medicinal use

äλιμον, Atriplex Halimus
4. 16. 5. Very dangerous to trees. άλίφλοιος (δρῦς), εςς δρῦς (3).

άλσίνη, Parietaria cretica 9. 13. 3. leaf of ἀριστολοχία comp. άλωπέκουρος, Polypogon monspelien-

7. 11. 2. flowers in a spike: described.

ἀμάρακον (ἀμάρακος), sweet marjoram, Origanum Majorana

1. 9. 4, evergreen; 6. 1. 1. in list of under-shrubs; 6. 7. 4. propagation: roots described; 6. 8. 3. flowering time; 9. 7. 3. in list οί ἀρώματα.

αμπελος (1) (leaf οἴναρον 9, 13. 5.), vine, Vitis vinifera

 1. 2. 1. has tendrils;
 1. 2. 7. bark;
 1. 3. 1. a typical 'tree';
 1. 3. 5. evergreen at Elephantine; 1.5.2. bark cracked and fibrous: bark in layers; 1.6.1. core fleshy; 1.6.3. roots thin; 1.6.5. roots branching upwards; 1.8.5. highest shoots 'roughest': 'eye' analogous to knot in other trees; 1.9.1. effect of pruning; 1.0.4. house breed; 1.10.5. 1. 10. 4. leaves broad; 1. 10. 5.

leaf divided; 1. 10. 7. long leafstalk: attachment of leaf-stalk; 1. 10. 8. leaves made of 'bark' and flesh; 1. 11. 4. seeds all together in a single case; 1.11.5. each grape separately attached; 1.12.1. taste of fruit; 1.12.2 taste of sap; 1.13.1. flower 'downy'; 1.13.3. flower surrounds fruit; 1. 13. 4. some kinds sterile; 1. 14. 1. bears on new shoots; 1. 14. 4. many cultivated forms; 2. 1. 3. propagation; 2.2.4. degenerates from seed; 2.3.1. sometimes spontaneously changes character; 2. 3. 2. a. ο κάπνειος varies in colour of grapes on same bunch: 2. 3. 3. sometimes bears fruit on the stem; etc.; 2.5.3. propagation; cannot be grafted; 2.5.4. propagation; 2.5.7. low ground suitable; great variety of kinds according to soil; 2. 6. 12. cuttings set upside down; 2.7.1. water-loving; 2. 7. 2. needs much pruning; 2. 7. 5. use of dust; 2. 7. 6. rootpruning; 3. 5. 4. autumn budding; 3. 17. 3. bark of κολου τία (2) comp.; 3.18.5. flower and fruit of ρους comp.; 3. 18. 12. cluster of berries of σμίλαξ (2) cluster of berries of σμίλαξ (2) comp.; 4.4.8. unnamed Indian tree (cotton-plant) planted in rows like ά.; 4.4.11. India confined to hill-country; 4.5.4. grows on Mount Tmolus and Mysian Olympus; 4.7.7. leaf of δεύδρον το ἐριφόρορν comp.; 4.7.8. occurs on island of Τylos; 4.13.2. some kinds short-lived; 4.13.4-6. said to be longest-lived of trees: reason: method of prolonging life son: method of prolonging life artificially; 4. 14. 2. young plants liable to 'sun-scorch'; 4. 14. 6. other diseases; 4. 14. 7. effects of injury to roots; 4. 14. 8. effect of rain on fruiting; 4. 14. 9. a special pest at Miletus; 4. 14. 10. effect of hot winds; 4.14.13. effects of frost; 4. 15. 1. outer bark can be stripped; 4. 16. 1. survives

splitting of stem; 4. 16. 6. natural antipathy of à. to ράφανος; 5. 3. 4. character of wood; 5. 4. 1. the less fruitful trees produce more solid wood; 5.9.4. wood, if damp, makes an evil smell when burnt for charcoal; 5. 9. 6. ἀθραγένη comp.; 8. 2. 8. à. in Melos; 9. 1. 6. time of tapping; 9. 13. 5. leaf and time of growth of πενταπετές comp.; 9. 18. 11. peculiar properties of certain local kinds.

αμπελος (2) (Mt. Ida), currant grape, Vitis vinifera, var. corinthiaca 17. 4. a local Idaean kind; 3. 17. 6. do. described.

ἄμπελος (3) ἡ ποντία, Fucus spiralis
 4. 6. 2. peculiar to certain waters;
 4. 6. 9. described.

αμπελος (4), ή άγρία (= μήλωθρον), bryony, Bryonia cretica 3. 18. 12. fruit of σμίλαξ (2) comp.

9. 14, 1. how long drug prepared from it will keep; 9, 20, 3, properties of root; medicinal use. άμυγδαλή, almond, Prunus Amyg-

dalus

 6. 3. large central root; 1. 9. 6. leaves produced early, but not shed early; 1.11.1. seed imme-diately within envelope; 1.11.3. seed in a woody shell; 1. 12. 1. taste of fruit; 1. 13. 1. flower 'leafy': flower of some kinds reddish; 1.14.1. bears on last year's wood; 2.2.5. degenerates from seed; etc.; 2.2.9. effects of cultivation; 2.2.11. effects of cultivation; z. z. 11.
do.: effect of tapping gum;
2. 5. 6. trees should be planted
far apart; 2. 7. 6. 'punishing'
the tree; 2. 7. 7. tapping the
gum; 2. 8. 1. apt to shed immature fruit; 3. 11. 4. fruit of
κράνεια comp; 3. 12. 1. leaf of
κράνεια comp; 4. 4. 7. fruit of
τέρμινθος ἡ Ἰνδική comp; 4.7.5.
rruit of unnamed Persian tree fruit of unnamed Persian tree (see App. (13)), comp.; 4.14.12. uninjured by special winds; 5.9.5. wood-ashes make pungent smoke; 7. 13. 6. flower appears before leaves and (new growth of) stem; 8. 2. 2. germination described; 9. 1. 2. sap gummy; 9. 1. 3. gum scentless; 9. 1. 5. gum useless; 9. 19. 1. leaf of δνοθήρας comp.

αμωμον, Nepaul cardamom, Amo-

mum subulatum 9. 7. 2. an ἄρωμα, Median or In-

dian. ἀνδράχλη, andrachne, Arbutus An-

drachne

drachne
1. 5. 2. bark readily drops off;
1. 9. 3. evergreen; 3. 3. 1. a
mountain tree; 3. 3. 3. evergreen; 3. 4. 2. time of budding;
3. 4. 4. time of fruiting; 3. 4. 6.
do; 3. 6. 1. slow growing (?);
3. 16. 5. described; 3. 16. 6. leaf
of κοκκυγέα comp.; 4. 4. 2. leaf
of μηλέα ἡ Περσική comp.; 4. 7.5.
an unnamed Persian tree (see
App. (14)) comp. 4. 15.1 does App. (14)), comp.; 4. 15. 1. does not perish if bark is stripped; 4. 15. 2, bark cracks; 5. 7. 6. wood used for parts of loom; 9. 4. 3. bark of σμύρνα comp.

άνδράχνη, purslane, Portulaca ole-

racea 7. 1. 2-3. time of sowing and germination; 7.2.9. root described.

ἀνεμώνη, anemone, Anemone spp. 7.8.3. leaves 'on the ground. ανεμώνη, anemone, Anemone coron-

aria 7. 7. 3. puts forth flower soon

after season of growth begins; 7. 10. 2. flowers in winter.

ανεμώνη ή λειμωνία, Anemone pavonina

6. 8. 1. flowering-time. ἀνεμώνη ἡ ὀρεία, Anemone blanda

6. 8. 1. flowering-time.

äνηθον (= ăννητος), dill, Anethum graveolens

 1. 11. 2. seeds naked; 1. 12. 2. taste of sap; 6. 2. 8. fruit of vaρης and vaρηκία comp.; also setting of flowers and fruit; 7. 1. 2-3. time of sowing and germination; 7. 2. 8. root described; 7. 3. 2. seeds described; 7. 4. 1. only one kind; 7. 6. 4. fruit of δρειοσέλινον comp.

ανθεμον, Anthemis chia, etc. (see below)

13. 3. flower attached above

each seed; 7. 14. 2. flowering begins at top: flower and fruit: several kinds (see below).

äνθεμον το ἀφύλλανθες, wild chamo-mile, Matricaria Chamomilla 7.8. 3. leaves 'on the ground.' ἀνθεμον το φυλλώδες, Anthemis chia 7.8. 3. leaves on the stem.

ανθέρικος, see ασφόδελος.

αννησον, anise, Pimpinella Anisum
1. 12. 1. scent.

ἄννητος (= ἄνηθον.) dill, Anethum graveolens 9. 7. 3. in list of ἀρώματα.

ἀντίρρινον, snapdragon, Antirrhinum Orontium

9. 19. 2. alleged magic properties: described.

ἀπάπη, dandelion, Taraxacum offici-

nale

 4. 8. (?) flower of χαμαιλέων comp.; 7. 7. 1. a λάχανον: classed as 'chicory-like' from classed as 'cnicory-inke non-its leaves; 7. 7. 3. season of growing; 7. 7. 4. prolonged flowering-time; 7. 8. 3. leaves 'on the ground'; 7. 10. 2. (?) flowers in winter, earliest of all; 7. 10. 3. flowers borne in succession; 7. 11. 3. flowering-time; 7. 11. 4. inedible: growth described.

άπαργία, hawk's beard, Crepis Columnae

7. 8. 3. leaves 'on the ground.' άπαρίνη, bedstraw, Galium Aparine 7. 8. 1. stem 'clasping,' but, for want of support, 'on the want of support, 'on the ground'; 7. 14. 3. clings to clothes: peculiar setting of

flower described; 8. 8. 4. grows specially among φακοί: growth described; 9. 19. 2. avtíppivov

comp. απιος (1), pear, Pyrus communis, var. sativa

var. saturat. 1. 2. 7. bark; 1. 3. 3. a tree whose stem is not single; 1. 8. 2. has less knots than $\Delta\chi\rho\dot{\alpha}_{5}$; 1. 10. 5. leaves round; 1. 11. 4. seeds all together in a single case; 1. 11. 5. seeds in a membrane; 1. 12. 2. taste of sap; 1. 13. 1. flower 'leafy'; 1. 13. 3. flower above fruit-case; 1. 14. 1. bears on last year's wood; 1. 14. 4. a cultivated form of άχράς; many cultivated forms; 2. 1. 2. propagation; 2. 2. 4. de-generates from seed; 2. 2. 5. seed produces wild form; 2.2.12. cannot be made out of $\frac{1}{4}\chi\rho_{0}$'s by cultivation; 2. 5. 3. grafting; 2. 5. 6. trees should be planted rather far apart; 2. 7. 7. 'punishing' the tree; 2. 8. 1. apt to shed immature fruit; 3. 2. 1. shed immature fruit; 3.2.1. produces less fruit than ἀχράς, but ripens more; 3.3.2. has better fruit and timber in low-lands; 3.4.2. time of budding; 3.6.2. formation of buds; 3.11.5. mountain and lowland forms comp.; 3.12.8. fruit of ὅη comp. as to keeping; 3.14.1. leaf of πτελέα comp.; 3.14.3. leaf of κληθρα comp.; 3.18.7. does not differ in kind from ἀχράς: 4.2.5. πεστέα comp. άχράς; 4. 2. 5. περσέα comp.; 4. 3. 1. size of λωτός (4) comp.; 4.4.2. thorns of μηλέα ή Περσική comp.; 4.5.3. abundant in Pontus; 4. 13. 1. shorter-lived than άχράς; 4.14.2. apt to get worm-eaten; 4.14.10. fruit gets wormeaten; 4. 14. 12. uninjured by special winds; 5, 3, 2, leaf of an unnamed tree comp. (see App. (20)); 9. 4. 2. leaf of λιβανωτός

comp. ἄπιος (2) (= i σχάς 9. 9. 5. = ράφαvos ή δρεία), spurge, Euphorbia Apios

9. 9. 5. medicinal use; 9. 9. 6. described.

άρακος, Vicia Sibthorpii
1. 6. 12. an unnamed plant (see App. (1)) comp.; 8.8.3. ('the rough hard kind') grows specially among φακοί.

άράχιδνα, Lathyrus amphicarpus
1.1.7. fruit underground; 1.6.12.

root like a second fruit.

άρία (= ίψος = φελλόδρυς 3. 16. 3.), holm - oak, Quercus Ilex var. agrifolia

3. 3. 8. doubt whether it has a flower; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 3. 16. 3. Dorian name for φελλόδρυς;

 17. 1. acorn of φελλός comp.;
 7. 2. (?) leaf of δάφνη (6) comp.; 5. 1. 1. time of cutting timber; 5. 3. 3. character of wood; 5.4.2. wood proof against decay; 5. 5. 1. wood hard to work; 5. 9. 1. wood makes good character. charcoal.

άριστολοχία, birthwort, Aristolochia rotunda

9. 13. 2. described: medicinal use; 9. 14. 1. how long drug will

keep; 9.15.5. grows in Arcadia; 9.20.4. cf. 9.13.2. ἄρκευθος (= κέδρας (3)), Phoenician cedar, Juniperus phoenicea

1. 9. 3. evergreen; 3. 3. 1. a mountain tree; 3. 3. 3. evergreen; 3. 3. 8. doubt whether it has a flower; 3. 4. 1. takes a year to ripen fruit; 3. 4. 5. time of fruiting; etc.; 3. 4. 6. do.; 3. 6. 1. slow-growing(?); 3. 6. 5. shallow-rooting according to Arcadians; 3.12.3-4. described: distinguished from κέδρος (1): 4. 1. 3. grows high on mountains, but not tall; 5. 7. 4. use of wood in house-building; 5. 7. 6. other uses of wood: does not decay; 9, 1, 2, sap gummy.

αρνόγλωσσον (= στελέφουρος 7.11.2., according to some, = opruk 7. 11. 2., according to some), plantain, Plantago maior

7. 8. 3. leaves 'on the ground'; 7. 10. 3. flowers borne in succession; 7. 11. 2. flowers in a spike: described by comparison with άλωπέκουρος.

 ἄρου, cuckoo-pint, Arum italicum
 1. 6. 7. root fleshy; 1. 6. 8. has a stout root and also fibrous roots: roots not tapering;
 1. 6. 10. cultivation;
 1. 18. 10. (?) flower made of flesh; 7. 2. 1. propagation; 7. 9. 4. root described; 7. 12. 2. root and leaves edible: use in surgery: special treatment to promote growth of root; one kind inedible (see δρακόντιον); 7.13.1. leaves described; 7.13, 2, no stem or flower.

άρρενόγονον (= θηλύγονον), dog mercury, Mercurialis perennis

9. 19. 5. properties: described. ἀσπάλαθος, Calycotome villosa

 7. 3. in list of ἀρώματα.
 ἄσπρις (=αἰγίλωψ (1)), Turkey oak, Quercus Cerris

3. 8. 2. one of the four Macedonian kinds of oak: acorns and

timber. ἀστέρισκος, Michaelmas daisy, Aster Amellus

4. 12. 2. seed of μελαγκρανίς comp

άσταφίς, Delphinium Staphisagria 9. 12. 1. medicinal use.

άσφάραγος, asparagus, Asparagus acutifolius 1. 10. 6. spines for leaves; 6. 1. 3.

do.; a wild under-shrub; 6, 4, 1. one of very few plants which are altogether spinous; 6. 4. 2. described.

 $\dot{a}\sigma\phi\delta\delta\epsilon\lambda\sigma$ (stem $\dot{a}\nu\theta\epsilon\rho\iota\kappa\sigma$), (= $\pi\delta\theta\sigma$ (2)), asphodel, Asphodelus ramo-

1. 4. 3. belongs to 'ferula-like' plants; 1. 10. 7. attachment of leaves; 6. 6. 9. leaves of vápκισσος (1) comp.; 7. 9. 4. root acorn-shaped; 7 12. 1. root acorn-shaped; 7 12. 1. root edible; 7.13.1. leaves described; 7.13.2-3. stem of loss comp.: largest stem of herbaceous plants: fruit inflorescence etc. described; worm which infests it: uses for food of stem and roots; 7.13.4. grown from seed; 9.9.6. leaf of ioxas comp.; 9.10.1. stem of exhégopos comp. by some.

ἄσχιον, puff-ball, Lycoperdon gigan-

teum 1. 6. 9. not a root, though underground.

άτρακτυλίς (= φόνος 6. 4. 6.), distaffthistle, Carthamus lanatus

6. 4. 3. a 'thistle-like' plant; 6. 4. 6. described: also called φόνος: reason; 9. 1. 1. juice blood-coloured.

άφάκη, tare, Vicia sativa var. angustifolia

8. 1. 4. (a pulse) sown late: 8. 5. 3. shape of pod; 8. 8. 3. mederivos

grows specially among a.; 8.11.1. seed does not keep.

άφάρκη (a natural hybrid between ανδράχλη and κόμαρος), hybrid arbutus, Arbutus hybrida

1.9.3. evergreen; 3.3.1. a mountain tree; 3.3.3. evergreen; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 5. 7. 7. uses of wood.

άφία, lesser celandine, Ranunculus Ficaria

7. 7. 3. puts forth flowers at season of growth.

άχράς, wild pear, Pyrus amygdali-

1. 4. 1. more fruitful than cultivated kind; 1. 8. 2. has more knots than aπιος; 1. 9. 7. time of shedding leaves; 1. 14. 4. a wild form of aπιος; 2. 2. 5. produced from seed of amios; 2.2.12. cannot be made into aπιος by cultivation; 3.2 1. produces more fruit than $\tilde{a}_{\pi ios}$, but ripens less; 3.3.1. a tree of mountain and plain; 3. 3. 2. has better fruit and timber in lowlands; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 3. 6. 1. slow growing (?); 3. 11. 5. mountain and lowland forms comp.; 3. 12. 8. fruit of δη comp. as to keeping; 3. 14. 2. bark of λεύκη comp.; 3. 18. 7. does not differ in kind from ἀπιος; 4. 13. 1. longer lived than ἀπιος; 5. 5. 1. cobblers' strops made of the wood.

ἀψίνθιον, wormwood, Artemisia Ab-sinthium

 1. 12. 1. taste of fruit; 4. 5. 1. seeks cold regions; 7. 9. 5. leaves and stem bitter, yet wholesome; 9. 17. 4. said to become by use non-poisonous to sheep.

βάλανος, Balanites aegyptiaca 4. 2. 1. peculiar to Egypt; 4. 2. 6. described.

βάλσαμον (gum ὁποβάλσαμον), balsam of Mecca, Balsamodendron Opo-

9. 1. 2. sap gummy; 9. 1. 7. time of tapping; 9. 4. 1. collection of gum; 9.6.1-4. described; habitat; method of collection; nowhere found wild: 9, 7, 3, in list of ἀρώματα.

βάτος, bramble, Rubus ulmifolius
1. 3. 1. a typical 'shrub'; 1. 5. 3.
thorns on wood; 1. 9. 4. evergreen; 1. 10. 6. leaf with spinous projections; 1. 10. 7. stem presently spinous; 3. 18. 3. grows in wet and dry places alike; 3. 18. 4. kinds distinguished; 3. 18. 12. cluster of berries of σμίλαξ (2) comp; 4. 8. 1. to some extent grows in marshes; 4. 12. 4. to some extent aquatic; 6. 1. 3. has spines on the shoots.

βληχώ, pennyroyal, Mentha Pulegium 9. 16. 1. leaf etc. of δίκταμνον

comp βλίτον, blite, Amaranthus Blitum

1. 14. 2. bears fruit both on top and at sides; 7.1.2-3. time of sowing and of germination; 7.2.7-8. root described; 7.3.2. seeds described; 7. 3. 4. seed borne both on top and at side;
7. 4. 1. only one kind.

βολβίνη, star-flower, Ornithogalum umbellatum

 13. 9. belongs to τὰ βολβώδη.
 βολβός, purse-tassels, Muscari comosum etc. (see below).

sum etc. (see below).

1. 6. 7. root in scales; 1. 6. 8. root not tapering; 1. 6. 9. no side roots; (part of) stem underground; 1. 10. 7. no leaf-stalk; attachment of leaves; 6. 8. 1. flowering time: used as a coronary plant; 7. 2. 1. propagation; 7. 2. 2. root makes offsets; 7. 2. 3. offsets specially numer-7. 2. 3. onsets specially fuller-ous; 7. 4. 12. formation of roots of κρόμυον comp.; 7. 9. 4. cf. 1. 6. 7.; 7. 12. 1. example of an edible root; 7. 12. 2. special treatment to promote growth of root; 7. 13. 1. leaves de-scribed; 7. 13. 2. flower-stem not the only stem; 7.13.4-5. grown from seed: seed sometimes takes two years to germinate; 7. 13. 7. root of νάρκισσος (1) comp.; 7. 13. 8.

several kinds: 7. 13. 9. roots of various plants comp.; 8. 8. 3. grows specially among mupos.

Βολβός δ ἐριόφορος, Pancratium maritimum

7. 13. 8. grows on beach: described: uses for food and cloth-

βουκέρας, $(=\tau \hat{\eta} \lambda \iota_s)$, fenugreek. Tri-

gonella Foenum-Graecum 4. 4. 10. an Indian plant (see App. (8)), comp. 8.8.5. aiµóδωρον parasitic on B.

βουμέλιος, ash, Fraxinus excelsior 3. 11. 4-5. described; 4. 8. 2. common in Egypt.

βούπρηστις,? 7. 7. 3. season of growing.

βούτομος, sedge, Carex riparia 1.5.3. stem very smooth; 1.10.5. leaves end in a point; further described; 4.8. 1. in list of τλ λοχμάδη; 4.10.4. described; 4.10.6. grows both on land and in water: grows on the floating islands of Lake Copais; 4. 10. 7. part used for food; 4.11.12. foliage of some κάλαμοι

βρόμος, oats, Avena sativa

8. 4. 1. seed has more coats than other cereals; 8. 9. 2. exhausts the soil: reason: like a wild plant.

βρύον, oyster-green, Ulva Lactuca 4. 6. 2. occurs generally in Greek waters; 4, 6, 6, described.

γήθυον, long onion, Allium Cepa var. 1.6.9. part of stem underground; 7. 1. 2-3. time of sowing and of 7.1.2-3. time of sowing and of germination; 7.1. 6. germination; 7.1. 7. bears fruit in second year: has single stem; 7.2.2-3. root makes offsets; 7.5.1. likes water; 7.5.3. bears transplanting; 7.5.5 seed does not keep well; 7.9.4. root in scales; 7.12.3. root of φάσγανον comp; 9.11.6. 'head' of TOUYNES transplanting; 7.5. Toot of φάσγανον comp; 9.11.6. 'head' of

στρύχνος ὁ μανικός comp. γήτειον (Attic for γήθνον), horn-onion, Allium Cepa var. 1. 10. 8. leaves hollow; 7, 4, 10,

described: cultivation (classed as a form of κρόμυον).

γλείνος, Acer creticum

3. 3. 1. name for lowland form of σφένδαμνος; 3. 11. 2. timber.
γλυκεία (εc. ῥίζα) (= ῥίζα Σκυθική q.v.),
9, 13. 2., liquorice, Glycyrrhiza

glabra.

γλυκυσίδη (= παιωνία q.v.), 9, 8, 6., peony, Paeonia officinalis.

γογγυλίς, truent operatus.
γογγυλίς, truent Rassica Rapa
1. 6. 6. root fleshy; 1. 6. 7. root of
bark and flesh; 7. 1. 2. time of
sowling; 7, 1. 7. germination;
7. 2. 5. survives and increases in size under a heap of soil: root described; 7. 2. 8. do.; 7. 3. 2. seeds described; 7. 3. 4. seed borne at side; 7. 4. 3. doubtful if more than one kind: seed, method of sowing, effect of weather; 7.5.3. bears trans-planting; 7.6.2. wild form distinguished; 7.9.4. root has bark.

δαῦκον (1), carrot, Daucus Carota 9. 15. 5. Arcadian drug: described

(see note).

δαῦκον (2), Malabaila aurea
9. 15. 8. grows about Patrai:
properties: root black; 9. 20. 2.
cf. 9, 15. 8.

δάφνη (1) (ἡ ημερος), (berry δαφνίς, 1. 11. 3.), sweet bay, Laurus nobilis

1. 5. 2. bark thin; 1. 6. 2. roots both stout and fine; 1.6.4. roots crooked; etc.; 1.8.1. few knots; 1. 9. 3. evergreen (cultivated and wild forms, see below); 1. 11. 3. fleshy seed in a shell (δαφνίς); 1. 12. 1. taste of fruit; 1.14.4. many cultivated forms 2. 1. 3. propagation; 2. 2. 6. sometimes improves from seed: 2. 5. 6. trees should be planted comp. to that of δ. ή λεπτόφυλλος; 3. 13. 5. leaflet of arry comp. to

leaf of δ. ή πλατύφυλλος; 3.14.3. flower (?) of κλήθρα comp.; 3. 15. 4. leaf of τέρμινθος comp; 3. 16. 4. leaf of κόμαρος comp.; 3.17.3. leaf of κολοίτια (2) comp. to δ. ή πλατύφυλλος; 4. 4. 12. leaf of an unnamed Arian shrub comp. (see App. (10)); 4. 4. 13. leaf of an unnamed Gedrosian tree comp. (see App. (11)); 4.5.3. does not thrive in cold regions; 4. 5. 4. grows in Propontis; 4.7.1. a class of marine Atlantic plants comp.; 4. 7. 4. leaf of an unnamed Arabian tree (see an unnamed Arabian tree (see App. (12b)) comp.; 4.13.3. after decaying shoots again from same stock; 4. 16. 6. spolls flavour of grape; 5. 3. 3-4. character of wood; 5.7.7. wood used for walking-sticks; 5. 8. 3. grows in lowland parts of Latium: abundant on Circeian promontory; 5. 9. 7. fire-drills made of the wood, because it does not wear away; 9. 4. 2. bark of λιβανωτός comp.; 9.4.3. leaf of λιβανωτός comp. (by some); 9. 4. 9. do.; 9. 10. 1. leaf of ελλέβορος ὁ μέλας comp. (by some); 9.15.5. δαῦκον (1) comp.: 9.20.1. one kind of πέπερι (fruit) comp.

δάφνη (2) η άγρία (= ονοθήρας), ole-ander, Nerium Oleander

9. 3. distinguished from δ. η

ημερος. δάφνη (3) η 'Αλεξανδρεία, Alexandrian laurel, Ruscus Hypophyl-

10. 8. bears fruit on leaves; 3. 17. 4. do.

δάφνη (4) ή λεπτόφυλλος, sweet bay. Laurus nobilis

12. 7. (see under δάφνη).

δάφνη (5) η πλατύφυλλος, sweet bay, Laurus nobilis 3. 11. 3., 3. 13. 5., 3. 17. 3. (see

under δάφνη). δάφνη (6) (= ϵ λάα (3) = App. (14)),

mangrove, Avicennia white officinalis

4.7.1. grows in 'Red Sea': 4.7.2. described: produces a drug for stanching blood.

δίκταμνον, dittany, Origanum Dictamnus

9. 16. 1-2. described: medicinal use: popular belief about its use to goats: comp. with ψευδο-δίκταμνον; 9. 16. 3. habitat.

δίκταμνον (ετερον), Ballota Pseudodictamnus

9. 16. 3. Cretan; has nothing in common with true & except the described: properties different.

διόσανθος, carnation, Dianthus inodorus

6. 1. 1. in list of under-shrubs; 6.6. 2. a cultivated under-shrub; a coronary plant: scentless; 6. 6. 11. grown from seed: woody; 6. 8. 3. flowering time.

διοσβάλανος (fruit κάρυον κασταναϊκόν), 4.8.11., chestnut, Castanea pesca

 12. 1. taste of fruit; 3. 2. 3. evidence that it is really wild: 3. 3. 1. a mountain tree; 3. 3. 8. doubt whether it has a flower: 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 3. 5. 5. winterbuds; 4. 5. 1. in list of northern trees; 4. 5. 4. abundant on Mount Tmolus and Mysian Olympus; 4. 8. 11. bark of root of λωτός (2) comp. to shell of chestnut.

διόσπυρος, Diospyros Lotus
3. 13. 3. fruit of κέρασος comp.
δόλιχος, calavance, Vigna sinensis
8. 3. 2. stem; 8. 11. 1. seed does not keep.

δόναξ (= κάλαμος ὁ Λακωνικός = κ. ὁ αὐλητικός = κ. ό συριγγιάς = κ. δ τοξικός), pole-reed, Arundo Donax 4.11.11. a kind of κάλαμος: habit

and habitat. δρακόντιον, edderwort, Dracunculus vulgaris

7.12.2. an inedible and poisonous kind of apov; 9.20.3. medicinal use: described.

δρυπίς, Drypis spinosa 10. 6. spinous-leaved.

δρῦς (1), oak, Quercus Robur
1. 2. 1. has galls (κηκίς); 1. 2. 7.
bark; 1. 5. 2. bark thick; bark
fleshy; 1.5.3. wood fleshy; 1.5.5.

wood heavy because it contains mineral matter; 1.6.1. core hard and close; 1. 6. 2. core called 'oak-black': core large and conspicuous; 1. 6. 3. roots many and long; 1. 6. 4. roots fleshy: deep-rooting; 1. 8. 5. diseased formation (κραδή); 1. 9. 5. an evergreen specimen; 1. 10. 6. leaves notched: leaves with spinous projections; 1. 10. 7. attachment of leaves; 1. 11. 3. seed in a leathery shell; 2. 2. 3. propagation; 2. 2. 6. deteriorates from seed; 3. 3. 1. tree of mountain and plain; 3. 3. 3. evergreen in some places; 3.3.8. doubt whether it has a flower (βρύον); 3.4.2. time of budding; 3.4.4. time of fruiting; 3.5. 1. periods of budding; etc.; 3.5. 2. galls; 3.5. 5. winter-buds; 3.6. 1. quick growing; 3.6. 5. instance of a deep-rooting tree; 3. 7. 4-6. various galls; etc.; 3. 8. 2. four or five kinds, viz. ημερίς ΟΓ έτυμόδρυς, αἰγίλοψ, πλατύφυλλος, φηγός, ἀλίφλοιος ΟΓ εὐθύφλοιος (Ναν recognised by inhabitants of Mt. ida); 3.16.1. leaf growth and bark of πρίνος comp.; 3. 16. 3. φελλόδρυς δρύς and πρίνος comp.; 4. 2. 8. common in Thebaid; 4. 5. 1. in list of northern trees; 4.5.3 grows in Porture; 4.14.10. infested by knips; 4.15. 2. survives stripping of bark for some time; 4.15. 3. effect of stripping bark in winter; 5.1.2. time of cutting timber: reasons; 5. 1. 4. do.; 5. 3. 1. core very close and heavy; 5. 3. 3. character of wood; 5. 4. 1. wood hard and heavy; 5. 4. 2. wood proof against decay; 5. 4. 3. wood does not decay if buried or soaked in water: rots in seawater: 5. 4. 8. effect of salt water on different parts; 5.5.1. wood hard to work; 5.6.1. wood contains mineral matter and so gives under weight: apt to split; 5.7.2. used for keel of triremes and for merchantmen to make

extra keel for hauling: does not glue well on to ἐλάτη οι πεύκη; 5. 7. 4. use of wood in housebuilding; 5.8.3. grows in Latium on Circeian promontory; 5. 9. 1. wood makes good charcoal, but inferior to apía and κόμαρος; 5. 9. 2. charcoal of this wood less esteemed by smiths than that of πεύκη; 8. 2. 2. germination from acorn described; 9. 9. 5. leaf of xauaiδρυς comp.

δρῦς (2) ἡ ἀγρία (= φηγός 3. 8. 2.), Valonia oak, Quercus Aegilops 1. 5. 2. rough bark; 3. 8. 2. see

under δρῦς.

δρῦς (3) ἡ ἀλίφλοιος (= δ. ἡ εὐθύ-φλοιος 3. 8. 2.), sea-bark oak, Quercus Pseudo-Robur

 8. 2. one of the five kinds of oak (Mt. Ida): = δ. η εὐθύφλοιος; (as. m. 10a); = 0. η ευθυφλοιος;
 3. 8. 3-4. acorns;
 3. 8. 6. φάσκος
 (q.v.) grows on it;
 3. 8. 7. timber;
 5. 1. 2. time of cutting timber.
 δρῦς (4) ἡ εὐθύφλοιος (= δ. ἡ ἀλόφλοιος

3. 8. 2.), sea-bark oak, Quercus Pseudo-Robur

3. 8. 2. one of the five kinds of oak (Mt. Ida).

δρῦς (5) ἡ ἡμερος (= ἐτυμόδρυς q.v. = ἡμερός (2)), true oak, Quercus Robur

8. 2. one of the five kinds of oak (Mt. Ida).

δρῦς (6) ἡ πλατύφυλλος, broad-leaved oak (scrub oak), Quercus lanuginosa

3. 8. 2. one of the five kinds of oak (Mt. Ida): fruit; 3. 8. 5. habit and timber; 3. 8. 6. galls: one of the four Macedonian oaks: has bitter acorns.

δρῦς (7) (φῦκος), Cystoseira ericoides 4. 6. 2. peculiar to certain waters; 4. 6. 7-8. described.

δρῦς (8) (ποντία), Sargassum vulgare 4. 6. 9. distinguished from δρύς (7): has a useful Bálavos.

έβένη (ἔβενος) (1), ebony, Diospyros Ebenum

1. 5. 4. wood heavy; 1. 5. 5. do. because of close grain: 1.6.1.

core hard and close; 4. 4. 6. described: two kinds distinguished (see below); 5. 3. 1. wood very close and heavy, especially the core; 5. 3. 2. colour of wood of répμιθος comp.: wood of an unnamed tree (see App. (20)) comp. to a variegated ε΄, 5. 4. 2. wood proof against decay; 9. 20. 4. colour and medicinal use of wood.

έβένη (2), Diospyros melanoxylon
4. 4. 6. a kind with inferior wood. είλετίας, 800 κάλαμος ὁ είλετίας έκτομον το μελαμπόδιον 800 έλλέβορος ο

μέλας

 cháa, olive, Olea Europea
 1. 3. 1. a typical 'tree'; 1. 5. 4.
 wood easily broken, not split: wood has many knots; 1.5.5. wood easily broken, because tough and not of straight grain; 6. 2. core not conspicuous; 1. 6. 3. roots both stout and thin; 1. 6. 4. roots branching; etc.; shallow rooting; roots crooked; etc.; 1. 8. 2. has less knotsthan κότινος; 1. 8. 6. liable to excrescences; etc.; 1.9. 3. evergreen; 1.10. 1. leaves inverted in summer; 1.10.2. colour of leaves; 1.10. 4. leaves narrow; 1.10.7. leaf-stalk short; tr. 1.11. seed enveloped in etc.; 1. 11. 1. seed enveloped in flesh and stone; 1. 11. 3. fleshy seed in a stone; 1. 11. 4. effect on fruit of rich feeding; 1. 12. 1. taste of fruit; 1. 13. 2. flower consists of one 'leaf' only partly divided; 1. 13. 3. flower surrounds fruit; etc; 1. 14. 1. bears on last year's wood; 1. 14. 2. bears fruit both on top and at side; 1. 14. 4. a cultivated form of κότινος; 2. 1. 2. propagation; 2. 14. 4. do.; 2. 2. 5. seed produces wild form; 2. 2. 12. cannot be etc.; 1. 11. 1. seed enveloped in wild form; 2. 2. 12. cannot be made out of κότινος by cultivation; 2. 3. 1. sometimes changes ton (2.3.1. sometimes changes to κότινος spontaneously; etc.; 2.5.8. grafting; 2.5.4. propagation; 2.5.6. do.: trees should be planted far apart; 2.5.7. low ground suitable; 2.7.2. needs much pruning; 2.7.3.

requires pungent manure and much water; 3. 2. 1. produces less fruit than κότινος but ripens more; 3. 12. 2. flower and fruit more; 3. 12. 2. flower and fruit of θηλυκράνεια comp.; 3. 17. 5. slze of fruit of συκή η Ιδαία comp.; 4. 2. 8. common in Thebaid; 4. 2. 9. character in Thebaid; 4. 3. 1. grows and bears well in Gyrenaics; 4. 4. 1. (?) distribution in Asia; 4. 7. 2. leaf and fruit of ἐλάα (3) comp.; 4. 7. 4. slze of fruit of unnamed Arabian tree comp. (see App. (12b)); 4. 13. 1. shorter-lived than κότινος; 4. 13. 2. story of a very old tree at Athens; 4. 13. 5. explanation of longevity; 4. 14. 2. diseases; 4. 14. 8. effect of rain on fruiting; 4. 14. 9. specially on fruiting; 4. 14. 9. specially apt to shed fruit at Taras; 4. 14. 10. other diseases: effect of hot winds; 4. 4. 11. suffers much from special winds; 4.16.1. improved by lopping branches; 5.3.3. character of wood; 5.3.7. 5.3.3. character of wood; 5.3.7. images made from the root; 5.4.2. wood proof against decay; 5.4.4. wood not eaten by teredon; 5.5.2. core not obvious: wherefore wood not apt to 'draw'; 5.5.3. core not obvious but exists; 5.6.1. wood out to split under pressure. obvious but exists; 5. 8. 1. wood apt to split under pressure; 5. 9. 6. wood good for kindling furnaces; 5. 9. 7. wood not suitable for fire-sticks: reason; 5. 9. 8. articles made of the wood have been known to produce shoots: instances; 6. 2. 1. leaf of κνέωρος ὁ λενκός comp.; 6. 2. 4. will not grow more than a short distance from the sea; 8. 2. 8. abundant in Melos; 9.18. 5. fruit of θηλύγουν comp. to βρύον of ἐ.: fruit of ἀρρενόγον comp. to undeveloped olive. rov comp. to undeveloped olive.

kh'a (2), Olea cuspidata
4. 4. 11. Indian (in hill-country

 $^{4}\lambda\dot{a}a'(3) (= \delta\dot{a}\phi\nu\eta(6) = App. (14)),$ white mangrove, Avicennia offi-

4.7.1. grows in 'Red Sea'; 4.7.2. described.

ἐλαίαγνος (properly ἐλέαγνος), goat willow, Salix Caprea
 4. 10. 1-2. in list of plants of L.

Copais: described.

έλάτη (1), silver-fir, Abies cephalonica 1. 1. 8. branches opposite; 1. 3. 6. refuses cultivation; 1. 5. 1. erect and tall; 1. 5. 2. bark in layers; 1. 5. 3. wood fibrous; 1. 5. 4. wood easily split; 1.5.5. do. because of straight grain; 1.6.3. root single; 1.6.4. roots fibrous; 1.6.5. roots not branching; 1.8.1. many knots; 1.8.2. 'male' has more knots than 'female'; 1.8.3. branches at 'remale'; 1.8. 3. branches at right angles; 1.9. 1. growth chiefly upwards; 1.9. 2. growth affected by position; 1.9. 3. evergreen; 1.10. 5. leaves described; 1.12. 1. taste of fruit; 1.12. 2. taste of sap; 1.13.1 flower yellow; 2. 2. 2. propagated only by seed; 2.7. 3. requires purport, respured. requires pungent manure; 3. 1. 2. grows only from seed; 3. 3. 1. a mountain tree; 3. 3. 3. evergreen; 3. 4. 5. time of flowering and fruiting; 3.5.1. period of budding; 3.5.3. do.; 3.5.5. winter-buds; 3.6.1. quick growing: even young tree fruits; 3. 6. 2. formation of buds; 3. 6. 4. not deep-rooting; 3. 6. 5. deep-rooting according to Arcadians; etc.; 3. 7. 1-2. dies if topped: formation of callus; 3. 9. 5. timber compared with πεύκη: etc.; 3. 9. 6. differences between 'male' and ences between 'female': described; 3. 9. 7. further comparison with meven: produces λοῦσσον; 3.9.8.do.: core and callus; 4. 1. 1. llkes shade; 4. 1. 2. grows tall in shade but has inferior timber; 4. 1. 3. grows high on mountains, but not tall; 4. 4. 1. (?) distribution in Asia; 4. 5. 1. in list of Northern trees; 4. 5. 3. does not grow in Pontus; 4. 15. 3. effects of stripping bark at various seasons; 4. 16. 1. topping fatal; 4. 16. 1-2. not injured by cut-

ting for resin; 4, 16, 4, said to perish if entirely deprived of its heart-wood; 5. 1. 1. time when timber is of best colour; 5. 1. 2. time of cutting timber; 5. 1. 4. do.; 5. 1. 5-6. timber comp. with wirvs; 5. 1. 7. uses of timber; 5. 1. 8. growth and character; 5. 1. 9-10. methods of cleaving; 5. 3. 3. character of wood; 5. 3. 5. used for doors; 5. 4. 4. less eaten by teredon than πεύκη; 5. 4. 6. wood, if barked just before time of budding, does not decay in water: story in proof of this; 5. 5. 1. knotty parts of wood hard to work; 5. 5. 2. core most obvious in e.; 5. 6. 1. wood good for struts: behaviour under pressure; 5, 6, 2, strongest of all woods; 5, 7, 1-2, use wood in ship-building: 5.7.4-5. uses of wood in housebuilding and crafts: the most generally useful of woods: more so than $\pi \epsilon \hat{\nu} \kappa \eta$; 5. 9. 8. wood has a peculiar exudation; 9.1.2. sap gummy; 9. 2. 1. production of resin (pntivn); 9. 2. 2. quality of resin.

λάτη (2), silver-fir, Abies pectinata
5. 8. 1. grows to great size in
Latium, but finer still in
Corsica; 5. 8. 3. grows in hill-

country of Latium.
'ἐλάτη' (3), 'sea-fir,'
Abies-marina Custoseira

4. 6. 2. peculiar to certain waters; 4. 6. 7-8. described.

έλατήριον, see σίκυος ὁ ἄγριος 4. 5. 1. in list of northern plants. έλειοσέλινον (= σέλινον τὸ έλειον),

marsh celery, Apium graveolens 7. 6. 3. comp. with σέλινον: medicinal use.

έλειόχρυσος, gold-flower, Helichry-sum siculum

6. 8. 1. flowering time; 9. 19. 3. alleged magic properties: described: medicinal use.

έλελίσφακος, salvia, Salvia triloba 4. a spineless wild under-shrub;
 5. 2. 5. like wild σφάκος: leaf described.

έλένιον, calamint, Calamintha incana

2. 1. 3. propagation; 6. 1. 1. in list of under-shrubs; 6, 6, 2, a cultivated under-shrub: a coronary plant: the whole plant scented; 6. 6. 3. woody: only one form; 6. 7. 2. said by some to have no fruit: 6, 7, 4, roots described.

ekschrod.

ελίκη, crack willow, Salix fragilis
3.13.7. Arcadian name for willow.
ελιέ, ivy, Hedera Helix
3. 18. 7–8. described: does not develop into κιττός; 3. 18. 8. kinds; 7. 8. 1. stem 'clasping.' η λευκή, white-berried ivy, Hedera Helix

3. 18. 8. several kinds. ελιξ ή ποικίλη $(=\dot{\epsilon}.\dot{\eta})$ Θρακία, 3.18.8.), ivy, Hedera Helix 3.18.8. several kinds.

ελιξ η χλοερά, ivy, Hedera Helix 3. 18. 8. described.

έλλεβορίνη, rupture-wort, Herniaria glabra

9. 10. 2. seed mixed with ἐλλέβορος ο λευκός to make an emetic.

έλλέβορος, hellebore, Helleborus cyclophyllus and Veratrum album

belongs to 'ferula-like' plants: has a hollow stem; 9.8.4. what part of root cut for medicinal use: 'bulbous' part a purge for dogs; 9. 8. 6. poisonous effect on those who dig it; precautions; 9. 9. 2. medicinal use; 9. 10. 1-4. kinds distinguished (see below); 9. 14. 1. how long drug will keep; 9. 17. 1-3, the drug can be made ineffectual by use: instances.

έλλέβορος ὁ λευκός, white hellebore,

Veratrum album

9. 10. 1. has nothing in common with è. ὁ μέλας except the name: divergent accounts given of the resemblances between the two plants: described; 9. 10. 2. not poisonous to sheep; when in season: distribution; 9.10.3-4. very local: local varieties, Οιταίος, Ποντικός, Έλεάτης, Μαλιώτης, Παρνάσιος, Αίτωλικός; Oiraios the best: properties of Έλεάτης; 9. 15. 5. grows in Arcadia; 9. 18. 2. restores scorpion to life when it has been killed with σκορπίος (3).

ἐλλέβορος ὁ μέλας (drug σησαμοειδές 9. 14. 4.), hellebore, Helleborus cyclophyllus

- 9. 8. 8. superstition as to gathering; 9. 10. 1. (see under è. ὁ λευκός): described; 9. 10. 2. poisonous to animals: 9, 10, 3, grows everywhere: some localities specified; 9. 10. 4. called by some έκτομον τὸ μελαμπόδιον: uses for purification and as charm; 9.14. 4. use of fruit in medicine; 9. 15. 5. grows in Arcadia; 9.16.6. leaf of ἐφήμερον comp.
- ěλυμος, Italian millet, Setaria italica 4. 4. 10. inflorescence of δρυζον comp.; 8. 1. 1. in list of 'summer crops' distinct from cereals and pulses; 8. 11. 1. seed keeps well.

ένθρυσκον, chervil, Scandix australis

ένθρυσκον, οποιτις, 7.7.1. a λάχανου. έπετίνη (?) (? πιτυίνη), Ajuga Ira 7.8.1. stem 'clasping,' but, for of support, 'on the ground.

Επιμενίδειος, εεε σκίλλα ή Έ.

ἐπίπετρον, stone-crop, Sedum anopetalum

7. 7. 4. flowerless.

ἐρέβινθος, chick-pea, Cicer arietinum 2. 4. 2. seed soaked before sowing; 2. 6. 6. size of some dates comp. 4.4.4. size of fruit of συκή ή Ἰνδική comp.; 4. 4. 9. not found in India; 6. 5. 3. leaf of a kind of τρίβολος comp.; 8. 1. 1. in list of pulses; 8. 1. 4. sown both early and late; 8. 2. 1. germination described; 8. 2. 3. comes up with several leaves: deep-rooting; 8. 2. 5. flowering time; 8. 2. 6. time of maturing seed; 8. 3. 2. stem; 8. 5. 1. several kinds: three mentioned, κριοί, ὑροβιαΐοι, οἱ ἀνὰ μέσον: White forms sweetest; 8. 5. 2. pod round: seeds comparatively few; 8.5.4. attachment of seed;

8. 6. 5. rain hurtful when ¿, is in flower: three kinds mentioned, μέλας, πυρρός, λευκός; 8. 7. 2. comp. with other pulses: destroys weeds: suitable soil: grows well after κύαμος; 8.9.1. exhausts the soil most of pulses; 8. 10. 1. diseases and pests; 8. 10. 5. infested by caterpillars; 8. 11. 2. only seed which does not engender 'worms' etc.: seed keeps well; 8, 11, 6, do. especially in hill country.

έρείκη, heath, Erica arborea

1. 14. 2. bears fruit on the top; 9. 11. 11. λιβανωτὶς ἡ ἄκαρπος grows where ε, is abundant. έρευθεδανόν, madder, Rubia tinc-

torum

6. 1. 4. a spineless wild undershrub; 7.9.3. roots red; 9.13.4. do.; 9. 13. 6. described: habit: habitat: medicinal use.

έρινεός, wild fig, Ficus Carica
1. 8. 2. has more knots than συκή;

1. 14. 4. wild form of συκή; 2. 2. 12. cannot be made into συκή by cultivation; 2. 3. 1. sometimes changes to συκή For the contraction of the cont lived than $\sigma v \kappa \hat{\eta}$; 4. 14. 4. not liable to diseases of $\sigma v \kappa \hat{\eta}$; 5. 6. 2. wood tough and easy to bend: uses; 5. 9. 5. wood makes pungent smoke.

(δένδρον τὸ) έριόφορον, cotton-plant,

Gossypium arboreum 4. 4. 8. (not named) clothes made

from it; 4.7.7-8. described. ερπυλλος (1) (ε. ο ημερος), tufted thyme, Thymus Sibthorpii

1. 9. 4. evergreen; 2. 1. 3. propagation; 6. 1. 1. in list of undershrubs; 6. 6. 2. a cultivated under-shrub: a coronary plant: whole plant scented; 6. 6. 3. woody: only one form; 6. 7. 2. said by some to have no fruit: 6.7.4. roots described; 6.7.5. growth peculiar: wild forms (see ερπυλλος (2)); 6. 7. 6. cultivation.

ερπυλλος (2) ὁ ἄγριος, Attic thyme, Thumus atticus

 7. 2. produces seeds, unlike
 (1); brought from Hymettus; sometimes quite like θύμος; 6. 7. 5. has various forms.

έρύσιμον, Sisymbrium polyceratium 8. 1. 4. sown later than cereals and pulses, a 'summer crop'; 8. 3. 1. leaf; 8. 3. 3. flower; 8. 6. 1. rain not beneficial after sowing; 8. 7. 3. doubtful if eaten green by animals: described.

έρυσίβη (cf. έρυσιβάν, έρυσιβώδης 8. 3. 2.), wheat-rust, Puccinia

graminis

8. 10. 1. a pest common to all crops (cereals, pulses etc.). έτυμόδρυς (= ήμερίς (2) 3. 8. 2. = δρῦς ή ήμερος), true oak, Quercus

Robur 3. 8. 2. one of the five kinds of oak (Mt. Ida) : = ημερίς : fruit ;

3. 8. 7. one of the four Macedonian kinds: has sweet acorns. Εὐβοϊκόν, see καρύα ή Εὐβοϊκή.

εύζωμον, rocket, Eruca sativa 1. 6. 6. root woody; 7. 1. 2-3.

time of sowing and germina-tion; 7. 2. 8. root described; 7. 4. 1. only one kind; 7. 4. 2. leaf of a kind of paparis comp.; 7. 5. 5. seed keeps well; 9. 11. 6.

leaf of στρυχνὸς ὁ μανικός comp. εὐθύφλοιος (δρῦς), see δρῦς (4). εὐώνυμος, spindle-tree, Ευουγπιιε europaeus

[3. 18. 13. described].

saffron, Colchicum parnassicum 9. 16. 6. a poison which has an antidote: described: effects.

ζειά, rice-wheat, Triticum dicoccum 2. 4. 1. seed, unless bruised, produces πυρός; 4. 4. 10. ἄρυζον comp.; 8. 1. 1. in list of cereals; 8. 1. 2. sown early; 8. 8. 3. ζ. and τίφη only plants which can change into something quite different (cf. 2. 4. 1.); 8. 9. 2. exhausts the soil: reason: likes rich soil: ζ . and $\tau i\phi \eta$ the cereals most like πυρός.

ζυγία, maple, Acer campestre 3. 3. 1. a mountain tree; name for mountain form of σφένδαμvos; 3. 4. 2. time of budding; 3. 6. 1. slow growing (?); 3. 11. 1-2. described; 5. 1. 2. time of cutting timber; 5. 1. 4. do.; 5. 3. 3. character of wood; 5. 7. 6. uses of wood.

ζωστήρ, see φῦκος (1) τὸ πλατύφυλλον.

ήδύοσμον, (= μίνθη), green mint, Mentha viridis

Menta viruis
7.7.1. a λάχανον.
ηλιοτρόπον, Heliotropium villosum
7.3.1. length of flowering season
of ώκιμον comp.; 7.8.1. stem
on the ground; 7.9.2. long
in flower; 7.10.5. evergreen;
7.15.1. flowering depends on
the heavenly hodies. the heavenly bodies.

ημερίς (1), gall-oak, Quercus infectoria

3.8.2. one of the five 'Idaean' kinds of oak: fruit; 3.8.4. habit and timber; 3.8.6. galls, ημερίς (2), (so-called by some) (= δρῦς η ημερος = ἐννμόδρος 3.8.2.), true oak, Quercus δοῦς του δρῦς του δρῦς στο κατά του δρῦς του δρῦς στο δροῦς στο δρο

3. 8. 2. bears sweet fruit.

ημεροκαλλές, Martagon lily, Lilium Martagon

6. 1. 1. in list of under-shrubs (see note); 6. 6. 11. grown from seed: a coronary plant.

ημιόνιον, milt-waste, Asplenium Ceterach

18. 7. properties of leaf: described: habitat: mules fond

ηρακλεία (= μήκων ή 'Ηρακλεία), Silene venosa

9. 15. 5. an Arcadian drug.

ηρακλεωτική (καρύα), see καρύα 'Ηρακλεωτική.

ήριγέρων, groundsel, Senecio vulgaris 7. 7. 1. a λάχανον: classed as 'chicory-like' from its leaves: 7. 7. 4. prolonged floweringtime; 7.10.2. flowers in winter. ηρύγγιον, eryngo, Eryngium campestre

6. 1. 3. has spines on the leaves:

a wild under-shrub.

θαψία, Thapsia garganica

9. 8. 3. most powerful juice from root: 9.8.5. superstition as to method of cutting; 9. 9. 1. root and juice used; 9. 9. 5. medi-cinal use; 9. 9. 6. described; 9. 11. 2. leaf of πάνακες το 'Ασκληπίειον comp.; 9. 20. 3. medicinal use: grows specially in Attica: properties; effect on foreign and native cattle.

θέρμος, lupin, Lupinus alba (puos, lūpin, Lupinus alba

1.3. 6. refuses cultivation; 1.7.3. seed roots through undergrowth; 3.2. 1. fruits better in wild state; 4.7.5. fruit of an unnamed Arabian tree (see App. (13)) comp.; 4.7.6. fruit of an unnamed Persian tree (see App. (13)) comp.; 4.7.7. fruit of a tree of the island of Tylos (see App. (13)) comp.; 8.1.3. sown early; 8.2.1. germination described; 8.5.2. seeds in compartments; 8.5.4. seeds in compartments; 8.5.4. attachment of seed; 8.7.3. not eaten green by any animal; 8. 11. 2. seed keeps well: like a wild plant; 8. 11. 6. seed keeps specially well in hill country; 8. 11. 8. peculiarities about sowing seed.

θηλύγονον (= αρρενόγονον), dog-mer-cury, Mercurialis perennis

9. 18. 5. properties: described. θηλυκράνεια, cornel, Cornus sanguinea

1.8.2. has less knots than κράνεια ; 3. 3. 1. tree of mountain and plain; 3. 4. 2. time of budding; 3. 4. 3. time of fruiting; 3. 4. 6. fruit inedible; 3. 12. 1-2. described; 5. 4. 1. less fruitful than κράνεια.

θηλύπτερις, bracken, Pteris aquilina 9. 18. 8. properties: distinguished

from πτερίς.

θηλύφονον (= ἀκόνιτον = μυόφονον = σκορπίος (3) 9. 18. 2.), wolf's bane, Acontium Anthora
 9. 18. 2. properties: habit: fatal

to the scorpion.

θήσειον. Corydalis densiflora 7. 12. 3. root bitter: medicinal use.

θραύπαλος, joint-fir, Ephedra campulopoda

3. 6. 4. very shallow-rooting: many roots; 4.1.3. likes shade θριδακίνη (properly, but not always

distinguished from θρίδαξ), wild

lettuce, Lactuca scariola
1. 10. 7. time of leaf-growth: stem presently spinous; 1.12.2. taste of sap; 7. 1. 2-3. time of sowing and of germination; 7. 3. 2. seeds; 7. 4. 1. several kinds; 7. 4. 5. do. viz. λευκή, πλατύκαυλος. στρογγυλόκαυλος, Λακωνική: differences; 7. 5. 4. pests; 7. 6. 2. wild form distinguished: medicinal 9. 8. 2. juice of stalk collected, with a piece of wool: 9.11.10.

to θ. ή πικρά. θρίδαξ, lettuce, Lactuca sativa

7. 2. 4. grows again when stem is cut: effect on flavour; 7. 2. 9. root described; 7. 5. 3. bears transplanting.

θρυαλλίς, Plantago crassifolia

7. 11. 12. flowers more or less in a

leaf of λιβανωτίς ή ἄκαρπος comp.

spike. θρύον, (a grass), Imperata arundi-

nacea 11. 12. foliage of some κάλαμοι

comp. θρύορον (?) (= στρυχνὸς ὁ μανικός 9. 11. 6.), thorn-apple, Datura Stramonium.

θυία (θύεια), odorous cedar, Juni-

perus foetidissima 1. 9. 3. evergreen; 3. 4. 2. time of budding; 3. 4. 6. time of fruiting; 4. 1. 3. grows on hill-tops. (60µa,? a madrepore 4. 7. 1. grows in Atlantic: turns

to stone).

θύμβρα (θύμβρον), savory, Satureia

Thymbra

1. 3. 1. (?) a typical under-shrub; 1. 12. 1. taste of fruit; 1. 12. 2. taste of sap; 6. 1. 4. a spineless wild under-shrub; 6. 2. 3. seed conspicuous: not, like $\theta \dot{\nu} \mu o s$, particular as to situation; 6.7.5. a wild form of ερπυλλος comp.; 7. 1. 2-3. time of sowing and of germination; 7. 1. 6. germina-

tion; 7. 5. 5. seed keeps well; 7. 6. 1. wild form distinguished. θύμον (1) (θύμος), Cretan thyme, Thirmbra capitata

1. 12. 2. taste of sap; 3. 1. 3. reproduces itself without seed; 6. 2. 3. two forms, black and white: seed inconspicuous:

6. 2. 4. requires sea-breezes.
 ('θύμον' (2), ? a madrepore

4.7.2. a marine plant which turns to stone: described).

θύον (θύα), thyine-wood, Callitria quadrivalvis

5. 3. 7. described: character and use of wood; 5.4.2. wood proof against decay.

ιασιώνη, bindweed. Convolvulus sepium

1. 13. 2. flower consists of one leaf.

ϊκμη, ? duckweed, Lemna minor 4. 10. 1-2. in list of plants of Lake Copais; 4.10.4. requires further investigation

ifia (1), oak-mistletoe, Loranthus europaeus

3.7.6. grows on oak and other trees; 3. 16. 1. grows on πρίνος. iξία (2) (= ἄκανθα (9) = ἄκανος == ιξίνη = χαμαιλέων ο λευκός), pine-

thistle, Atractylis gummifera
9. 1. 3. Cretan: produces a gum.
iξίνη(gum (ἀκανθική) μαστίχη 6. 4. 9.,

9. 1. 2.) (= ἄκανθα (9) = ἄκανος = ἰξία (2) = χαμαιλέων ὁ λευκός), pine-thistle, Atractylis gummifera

6. 4. 3. a 'thistle-like' plant; 6. 4. 4. time of growing; 6. 4. 9. described; 9. 1. 2. produces a gum called μαστίχη.

ιον (= ιωνία = ιον το λευκόν), gilliflower, Matthiola incana

1. 9. 4. evergreen; 2. 1. 3. propagation; 4. 7. 4. colour and scent of unnamed Arabian tree (see App. (12a)) comp.; 6. 1. 1. in list of under-shrubs; 6. 6. 1. a cultivated undershrub: a coronary plant; sweetscented; 6.6.5. sweetest-scented at Cyrene; 6. 6. 11. grows from

seed; woody; 6.8.5. position and climate important for fragrance: flowers very early in Egypt; 6. 8. 6. on mountains blooms well, but has inferior scent; 7. 6. 4. wild form quite distinct, alike only in leaf.

ίου το λευκόν (= λευκόϊον (1) = ίωνία ή λευκή), gilliflower, Matthiola in-

 18. 13. flower of εὐώνυμος comp.; 4. 7. 8. flower of an Arabian tree (see App. (15)) comp; 6. 6. 3. several colour forms; 6. 6. 7. distinguished from i. το μέλαν; 6. 8. 1-2. flowering-tine; 6. 8. 5. plant lives three years at most: degracers with age; 7. 8. generates with age: 7, 8, 3, leaves 'on the stem.

τον τὸ μέλαν (= ἰωνία ἡ μέλαινα),

violet, Viola odorata
1. 13. 2. has a 'twofold' flower;
6. 6. 3. only one form; 6. 6. 7. distinguished from i. τὸ λευκόν: 6. 8. 1-2. a coronary plant: flowering time.

ιπνον, ? marestail, Hippuris vulgaris 4. 10. 1-2. in list of plants of Lake Copais; 4.10.4. requires further

investigation.

ἰππομάραθον (≔μαγύδαρις), Prangos

ferulacea

6. 1. 4. a spineless wild undershrub: belongs to 'ferula-like plants.

iπποσέλινον, Alexanders, Smyrnium Olusatrum

1. 9. 4. evergreen; 2. 2. 1. propagation; 7. 2. 6. root of τεύτλιον comp.; 7. 2. 8. root; 7. 6. 3. comp. with έλειοσέλινον: medicinal use; 9. 1. 3. root produces a gum: which is like σμύρνα; 9. 1. 4. propagated from a δάκρυον: a popular error about è. and σμύρνα; 9.15.1. grows in Arcadia.

iπποφαές, see τιθύμαλλος.

ιππόφεως, spurge, Euphorbia acan-

thathamnos

6. 5. 1. in list of spinous plants which have leaves as well as spines; 6. 5. 2. has no spines on the leaves.

lpis, iris, Iris pallida, etc.

1. 7. 2. root fragrant; 4. 5. 2. grows best in Illyria on shores of Adriatic: 6. 8. 3. a coronary plant: flowering time; 7. 13. 1. 7. 13. 2. leaves described; flower-stem not the only stem : stem comp. with ἀσφόδελος; 9.7.3. in list of apmuara; 9.7.4. only European αρωμα; best in Illyria: preparation; 9. 9. 2. perfume.

ίσχαιμος, Andropogon Ischaemum 9. 15. 3. Thracian: properties.

ισχάς (= ἄπιος (2) 9. 9. 5. = ράφανος opeia), spurge, Euphorbia Apios.

ίτέα, willow, Salix spp.

1. 4. 2. lives near water; 1. 4. 3. 'amphibious'; 1. 5. 1. crooked and low; 1. 5. 4. wood light; 3. 1. 1. propagation; 3. 1. 2. seems to have no fruit, yet reproduces itself: instance; 3.1.3. sheds its fruit unripened; 3.3.1. tree of mountain and plain; 3.3.4. a question if it bears fruit; 3. 4. 2. time of budding; 3. 6. 1. quick or slow grower?; 3. 13. 7. described: kinds (see below): called in Arcadia έλίκη; 3. 14. 4. leaf of κολυτέα comp.; 4. 1. 1. likes wet ground; 4. 5. 7. common in some Mediterranean regions; 4.8.1. grows partially in water; 4. 10. 1. in list of plants of Lake Copais; 4. 10. 6. grows both on land and in water; 4. 13. 2. shorter-lived by water; 4. 16. 2. grows again after being cut or blown down; 4. 16. 3. instance of a tree which survived the lopping of its branches; 5. 3. 4. character of wood; 5. 7. 7. uses of wood; 5. 9. 4. wood makes an evil smoke when burnt for charcoal.

ίτέα ἡ λευκή, white willow, Salix alba

3. 13. 7. described.

ίτέα ἡ μέλαινα, Salix amplexicaulis
3. 13. 7. described.

ίφυον, spike-lavender, Larandula Spica

6. 6. 11. a coronary plant: grown

from seed; 6. 8. 3. flowering time.

τψος, ? cork-oak, Quercus Suber (G. from Plin. 16. 98. Hesych. has ϊψος = κισσός)

3. 4. 2. time of budding.

ὶωνία (=ἴον q.v.=ὶωνία ἡ λευκή = ἰόν τὸ λευκόν = λευκόῖον (1)), gilliflower, Matthiola incana.

ίωνία ή λευκή (= ίωνία = ἴον q.v.), gilliflower, Matthiola incana. ιωνία η μέλαινα (= ιον τὸ μέλαν φ.υ.), violet, Viola odorata.

κάκτος (1), cardoon, Cynara Cardunculus

10-11. a 'thistle-like' 6. 4. plant: described: peculiar to Sicily.

κάκτος (2), artichoke, Cynara Scoly-

 4. 11. has erect 'stalk' called πτέρνιξ: described; edible; base of receptacle called σκαλίας.

κάλαμος, reed 1. 5. 2. bark fibrous: bark in one layer; 1. 5. 3. stem jointed; 1.6.2. core membranous; 1, 6, 7, 1.0.2. Core immorations; 1.6. 7. root jointed; 1.6. 10. roots large and numerous; 1.8. 3. joints analogous to 'knota': 1.9. 4. evergreen; 1.10. 5. leaves end in a point; further described; 1.10. 9. leaves made of fibre: leaf-stalk made of fibre; 2. 2. 1. (a kind of) propagation; 4. 8. 1. in list of τὰ λοχμώδη; 4. 8. 7. κύαμος ο Aiγύπτιος comp.; 4. 8. 8. thickness of root of κύαμος ο Αίγύπ-Tios comp.; 4. 9. 1. class of rivers in which κ . grows; 4.9.3. has 'side-growths'; 4.10.1. in list of plants of Lake Copais; 4.10.6. grows both on land and in water; 4.10.7. effect of drought: 4. 11. 1. distinguished from k. ὁ αὐλητικός (see below): a stout and a slender form (ὁ χαρακίας and ὁ πλόκιμος) (see below); 4.11.10-13. other forms; 6.2.8. setting of leaves of νάρθηξ and ναρθηκία comp.; 9.16.1. δίκταμνον kept έν καλάμφ.

κάλαμος ὁ αὐλητικός (= κ, ὁ Λακωνικός

 $= \kappa$. δ συριγγίας $= \kappa$. δ τοξικός = κ. ο χαρακίας = δόναξ, polereed, Arundo Donax

4. 10. 1. in list of plants of Lake Copais; 4. 10. 6. grows only in water; 4. 11. 1. distinguished from the ordinary form of κ .; 4. 11. 2. not true that it takes nine years to grow; 4. 11. 3. conditions of growth; 4. 11. 4. described by contrast with other κάλαμοι; 4. 11. 4-7. manufacture of the mouthpleces of pipes; 4. 11. 8-9. distribution in region of Lake Copais.

κάλαμος ὁ είλετίας, Ammophila arundinacea

4. 11. 13. the 'male kind' of k. eπίγειος, so called by some.

κάλαμος (ἐπίγειος), bush-grass, Calamogrostis Enigeios

4.11.13. described : growth comp. to ἄγρωστις.

κάλαμος ο εὐώδης, sweet flag, Acorus Calamus

4. 8. 3. grows in a Syrian lake: 9.7.1. habitat (east of Lebanon): described: fragrance; 9.7.3. in list of ἀρώματα.

κάλαμος ὁ Ἰνδικός, bamboo, Bambusa arundinacea

4. 11. 13. described. κάλαμος ὁ Ἰνδικός ('male'), Male bamboo, Dendrocalamus strictus 4. 11. 13. distinguished as solid.

κάλαμος ὁ Λακωνικός (= κ, ὁ αὐλητικός κ. ὁ συριγγίας = κ. ὁ τοξικός = κ. ὁ χαρακίας = δόναξ), polereed, Arundo Donax

11. 12. colour.

κάλαμος ὁ πλόκιμος, spear-grass, Phragmites communis

4. 11. 1. pliant reed; compared with κ. ὁ χαρακίας: grows on floating islands of Lake Copais.

κάλαμος ὁ συριγγίας (= κ. ὁ αὐλητικός = κ. ὁ Λακωνικός = κ. ὁ τοξικός = κ. ὁ χαρακίας = δόναξ), polered, Arundo Donax
4. 11. 10 described.

κάλαμος ὁ τοξικός (Κρητικός) (= κ. ὁ αὐλητικός = κ. ὁ Λακωνικός = κ. ὸ συριγγίας = κ. ὁ χαρακίας = δό-ναξ), pole-reed, Arundo Donax 4. 11. 11. described.

κάλαμος ὁ χαρακίας (= κ. ὁ Λακωνικός etc.), pole-reed, Arundo Donax

4. 11. 1. stout form: described: grows in reed-beds of Lake Conais.

κάλαμος (other kinds)
4. 11. 10. briefly described.

κάππαρις, caper, Capparis spinosa 1. 3. 6. refuses cultivation; 3. 2. 1. fruits better in wild state: 4. 2. 6. fruit of βάλανος comp.; 6. 1. 3, has spines on the shoots; 6. 4. 1. has spines on leaves as well as on stem; 6.5.2. described; 7.8.1. stem 'on the ground'; 7.10.1. grows and flowers entirely in summer.

κάρδαμον, cress, Lepidium sativum
1.12.1. taste of fruit; 7.1.2-3.
time of source and germination; 7.1.6. germination; 7.4.1. only one kind; 7.5.5. seed keeps well.

καρδάμωμον, cardamom, Elettaria Cardamomum

9. 7. 2. an ἄρωμα, Median or Indian; 9 7. 3. in list of ἀρώματα. καρύα (fruit κάρυον), hazel, Corylus avellana

1. 12. 1. taste of fruit; 3. 2. 3. evidence that it is really wild; 3.3.1.a mountain tree; 3.4.2. time of budding; 3.4.4. time of fruiting; 4.5.4. abundant on Tmolus and Mysian Olympus; 8.2.2. germination described.

καρύα ή Εὐβοϊκή, sweet chestnut, Castanea vesca var. (improved form) 1. 11. 3. seed in a leathery shell: 1. 5. 4. common in Euboea and Magnesia; 5. 4. 2. wood proof against decay; 5. 4. 4. wood does not decay in water (?) 5. 6. 1. wood makes a noise when about to split: instance: 5. 7. 7. uses of wood: does not rapidly decay; 5. 9. 2. charcoal of this wood used in iron-mines.

καρύα ή 'Ηρακλεωτική ('Ηρακλεώτις) (fruit κάρυον), filbert, Corylus avellana vars.

1. 3. 3. effect of not pruning; 1. 10. 6. leaves notched; 1.11.1. seed immediately within enve-

lope; 1.11.3. seed in a woody shell; 3. 3. 8. doubt whether it has a flower (ioulos); 3.5.5-6. catkins; 3. 6. 2. formation of buds; 3. 6. 5. deep-rooting according to Arcadians: etc.; 3. 7. 3. catkins; 3. 15. 1-2. described: kinds.

καρύα ή Περσίκη, walnut, Juglans

regia
3.6.2. formation of buds; 3.14.4. leaf of σημύδα comp.

κασία, cassia, Cinnamomum iners 4. 4. 14. in list of oriental aroma-

tic plants; 9. 4. 2. Arabian; 9. 5. 1. and 3. described; method of collection; 9. 7. 2. Arabian: 9. 7. 3. in list of ἀρώματα,

καυκαλίς, Tordylium apulum

 7. 7. 1. a λάχανον.
 κέγχρος, millet, Panicum miliaceum
 1. 11. 2. seeds in a husk; 4. 4. 10. inflorescence of ὅρυζον comp.; 4. 8. 10. fruit of λωτός (2) comp.; 10. 3. size of seeds of σίδη comp.;
 11. 1. in list of 'summer crops' distinct from cereals and pulses; 8.1. 4. sown later than cereals and pulses; 8.2.6. time of maturing seed; 8.3.2. stem; 8.3.3 flower; 8.3.4. seed abundant; 8.7.3. needs little water: comp. with $\mu \epsilon \lambda \iota \nu o s$; 8. 9. 3. reasons why it might have been expected to exhaust the soil: contrasted with pulses as to 'lightness'; 8.11.1. seed keeps well; 8.11.6. do. specially in hill-country; 9. 18. 6. fruit of κραταιγών comp.

κεδρίς, juniper, Juniperus communis 1. 9. 4. evergreen: a dwarf kind (cf. κέδρος 3. 13. 7.); 1. 10. 6. leaf spinous at tip; 1. 12. 1. taste of fruit.

κέδρος (1) (= ὀξύκεδρος 3. 12. 3.), prickly cedar, Juniperus Oxycedrus

 5. 3. wood not fleshy;
 1. 10. 6.
 leaf spinous at tip;
 6. 5. shallow-rooting according to Arcadians; 3.10.2. μίλος comp.; 3. 12. 3-4. described: two kinds, ή Λυκίη and ή Φοινική (? Φοινι-

κική) (see κέδρος (3)); distinguished from άρκευθος; 3. 13. 7. has a dwarf form (? κεδρίς, cf. 1.9. 4.); 4.3.3. size of fruit of παλίουρος ο 'Αιγώπτος comp.; 4.5.2. grows on Thracian and Phygian mountains; 4.16. 1. some think topping fatal; 5.3.7. images made from the wood; 5. 4. 2. wood proof against decay; 5.9.8. wood exudes moisture: hence 'sweating' statues; 9. 1. 2. sap gummy.
κέδρος (2), Syrian cedar, Juniperus

excelsa

3.2.6. characteristic of mountains of Cilicia and Syria; 4. 5. 5. grows in Syria and is used for ships; 5. 7. 1-2. use of wood in ship-building; 5. 7. 4. use of wood in house-building; 5.8.1. remarkably fine in some regions e.g. Syria.

κέδρος (3), ή Φοινικική (= ἄρκευθος), Phoenician cedar, Juniperus phoenicea

3, 12. 3. εεε κέδρος (1); 9, 2. 3. said to be burnt for pitch in Syria.

κέδρος (4) ή Δυκίη

3, 12, 3, a kind so distinguished by some from κέδρος (3). κενταύριον, centaury, Centaurea sa-

lonitana

1. 12. 1. taste of fruit; 3. 3. 6. only bears fruit in hill country; 4.5.1. seeks cold regions; 7.9.5. leaves and stems bitter, yet wholesome; 9. 1. 1. juice blood-red; 9. 11. 6. juice mixed with στρύχνος ὁ μανικός to make a potion.

κενταυρίς, feverwort, Erythraea Centaurium

9. 8. 7. superstition as to gathering; 9. 14. 1. how long drug will keep.

κεντρομυρρίνη (= μυάκανθος), butcher's broom, Ruscus aculeatus 3. 17. 4. bears fruit on its leaves.

κεράϊς (= ράφανος ή άγρία 9. 15. 5.), charlock, Raphanus Raphanistrum.

κέρασος (= λακάρη), bird-cherry, Prunus avium

3, 13, 1-3, described; 4, 15, 1, bark

can be stripped: 9. 1. 2. sap gummy.
κεραύνιον, 'thunder-truffle,' Tuber

aestivum

6. 5. has no roots.

κερκίς (1), Judas-tree, Cercis Siliquastrum

1. 11. 2. seeds in a pod.

κερκίς (2), aspen, Populus tremula 3. 14. 3. described.

κερωνία (= συκή ή Αίγυπτία 1. 11. 2.), carob, Ceratonia Siliqua

1. 11. 2. seeds in a pod; 1. 13. 2. bears on stem and branches; 4. 2. 4. described.

κήλαστρος (κήλαστρου), holly, Ilex Aquifolium

- 1. 3. 6. refuses cultivation; 1. 9. 3. evergreen; 3.3.1. tree of mountain and plain; 3. 3. 3. ever-green; 3. 4. 5-6. time of fruiting; said to lose fruit in winter;
 4. 1. 3. grows in very cold positions: 5, 6, 2, colour of wood of φιλύκη comp.; 5. 7. 7. wood used for walking-sticks.
- κινάμωμον, cinnamon, Cinnamomum Cassia
 - 4. 4. 14. in list of oriental aromatic plants; 9.4.2. Arabian; 9.5.1-2. two kinds, white and black, described: habitat: method of collection: a story; 9.7.2. in list of αρώματα.

κίσθος, rock-rose, Cistus app.

6. 1. 4. a spineless wild under-shrub; 6. 2. 2. described: two forms (see below).

κίσθος ὁ ἄρρην, Cistus villosus 6. 2. 1. described. κίσθος ὁ θηλυς, Cistus salvifolius

6. 2. 1. described.
 κιττός, ivy, Hedera Helix

1. 3. 2. a shrub which becomes 3. 2. a snrub which becomes tree-like; 1. 9. 4. evergreen; 1. 10. 1. leaves change shape with age of plant; 1. 10. 7. long leaf-stalk; 1. 13. 1. flower downy'; 1. 13. 4. attachment of flower; 3. 4. 6. time of fruiting; 3. 10. 5. fruit and leaf of φίλυρα comp.; 3. 14. 2. leaf of κερκίς (2) comp.; 3. 18. 6. kinds distinguished (see below); 3. 18. 7. distinguished from

έλιξ; 3. 18. 9-10. described: habit etc.; 3. 18. 11. cluster of berries of σμίλαξ (2) comp.: described; 4. 4. 1. distribution in Asia; 4. 16. 5. overgrowth of k. dangerous to trees; 5. 3. 4. character of wood; 5. 9. 6. wood said to make best fire-sticks: 5. 9. 7. the stationary fire-stick should be made of κ. or άθραγένη; 9. 13. 6. leaf of έρευθεδανόν comp.

KITTOS O EXIE, BEE EXIE

κιττὸς ὁ λευκός, white-berried ivy. Hedera Helix

3. 18. 6. described : several kinds : one = κορυμβίας, one = κ. ο 'Αχαρ-νικός; 3. 18. 9. roots; 3. 18. 10. fruit; 9.18.5. properties of fruit.
κιττὸς ὁ μέλας, black-berried ivy, Hedera Helix

3. 18. 6. several kinds; 3. 18. 9. roots: 3, 18, 10, fruit.

κιχόριον (κιχόρη), chicory, Cichorium

Intybus 1 1.00.7. attachment of leaves; 7.7.1. a λάχανον; a class of plants called 'chicory-like' from their leaves; 7.7.3. season of growing; 7.8.3. leaves 'en the ground' and 'on the stem'; etc.; 7.9.2. long in flower; 7.10.3. flowers borne in succession; 7.11.3. root, inflorescence and 7. 11. 3. root, inflorescence and seed-vessel described; 9. 12. 4. μήκων ἡ ροιάς comp. to κ. τὸ ἄγριον; 9. 16. 4. leaf of ἀκόνιτον comp

κλήθρα, alder, Alnus glutinosa

νήθρα, alder, Ainus giutinosa
1.4.3. 'amphibious'; 3.3.1. tree
of mountain and plain; 3.3.6.
does not always fruit; 3.4.2.
time of budding; 3.4.4. time
of fruiting; 3.6.1. slow growing (?); 3.6.5. roots slender
and 'plain,' according to Arcadians; 3.14.3. described; 3. 15. 1. leaf of καρύα η Ήρακλεωτική comp.; 4. 8. 1. grows partially in water.

κλινότροχος (?), Acer Pseudo-platanus 3. 11. 1. a form of σφένδαμνος. κλύμενον, honeysuckle, Lonicera

etrusca 9. 8. 5. superstition as to time of cutting; 9. 18. 6-7. properties of fruit.

κνέωρον (berry Κνίδιος κόκκος). Daphne Gnidium

6. 1. 4. a spineless wild under-shrub; 9. 20. 2. berry described; medicinal use and properties.

κνέωρος ο λευκός, Daphne oleoides
6. 2. 2. distinguished from κ. ο

μέλας; use of root. κνέωρος ὁ μέλας, Thymelaea hirsuta

1. 10. 4. leaves fleshy: 6, 2, 2, see

κ. ὁ λευκός.

 $\kappa \nu \hat{n} \kappa o s (= \kappa, \delta, \tilde{n} \mu \epsilon \rho o s = \kappa \rho \delta \kappa o s, \delta, \delta, \delta \alpha \nu$ θώδης), safflower, Carthamus tinctorius etc. (see below)

1. 13. 3. flowers attached above each seed; 6.1.3. a wild undershrub: has spines on the leaves; 6. 4. 3. a 'thistle-like' plant; 6. 4. 4. no side-growths; 6. 4. 5. three forms distinguished and described, one cultivated (see below); 6. 6. 6. seed of ρόδον comp.

κνήκος ή αγρία, Carthamus leucocaulos 6. 4. 5. distinguished from κ. ή ημερος.

κνήκος η άγρία (ἐτέρα), Cnicus benedictus

6. 4. 5. described.

κνήκος ή ήμερος, Carthamus tinctorius 6. 4 5. distinguished from wild kinds.

Κνίδιος κόκκος, 866 κνέωρον

(= κουκιόφορου), doum-palm, Hyphaene thebaica

1. 10. 5. reedy leaves; 2. 6. 10. a shrubby palm: Ethiopiau. κοκκυγέα, wig-tree, Rhus Cotinus 3. 16. 6. described.

κοκκυμηλέα (fruit κοκκύμηλον), plum-

tree, Prunus domestica
1. 10. 10. fruit made of flesh and fibre; 1. 11. 1. seed enveloped in flesh and stone; 1.12.1. taste of fruit; 1. 13. 1. flower 'leafy' 1. 13. 3. flower above fruit-case: 3. 6. 4. very shallow-rooting: few roots; 3. 6. 5. deep-rooting according to Idaeans; etc.; 4. 2. 3. size of fruit of συκή ή Κυπρία comp.; 4.2.5. fruit-stone of περσέα comp.

κοκκυμηλέα (ἡ Λίγυπτία) (sebesten), Cordia Myza

4. 2. 10. described.

κολοιτία (1) (κολουτέα 3. 17. 2.: cf. 3. 17. 3. n.), Cytisus acolicus 1. 11. 2. tree of Lipari islands:

seeds in a pod; 3. 17. 2. described.

acribed.

αλοιτία (2), Salix cinerea
3. 17. 3. Idaean: described.
κολοκύντη, gourd, Cucurbita maxima
1. 11. 4. seeds in a row: 1. 12. 2.
taste of sap; 1. 13. 3. flower
attached above fruit: 2. 7. 5.
use of dust; 7. 1. 2–3. time of
sowing and of germination;
7. 1.6 argmination: 7. 2.0 seed 7. 1. 6. germination; 7. 2. 9. rcot described; 7.4.1. several kinds; 7. 5. 5. seed does not keep well.

κολυτέα, bladder-senna, Colutea arborescens

[3. 14. 4. described.]

κόμαρος (fruit μεμαίκυλον 3. 16. 4.), arbutus, Arbutus Unedo

1. 5. 2. bark readily drops off; 1. 9. 3. evergreen; 3. 16. 4. de-scribed; 3. 16. 6. leaf of κοκκυγέα comp.; 5.9.1. wood makes good charcoal.

κόμη = τραγοπώγων 7. 7. 1. q.v. κόνυζα, Inula spp.

6. 1. 4. a spineless wild undershrub; 6. 2. 5. two kinds described and compared ('male' and 'female') (see below); 7. 10. 1. grows and flowers entirely in summer.

κόνυζα ή ἄρρην, Inula viscosa 6. 2. 5.

κόνυζα ή θήλεια, Inula graveolens 6. 2. 5.

κορίαννον, coriander, Coriandrum sativum

1. 11. 2. seeds naked; 7. 1. 2-3. time of sowing and germination: 7. 1. 6. germination; 7. 2. 8. root described; 7. 3. 2. seeds described; 7. 4. 1. only one kind; 7. 5. 4. effect of hot weather; 7. 5. 5. seed keeps well.

κορυμβίας, see κιττός ὁ λευκός. κόρχορος, blue pimpernel, Anagallis

caerulea

7. 7. 2. a Aáxavov: proverbial for bitterness.

κορωνόπους, hartshorn. Plantago Coronopus

7. 8. 3. leaves 'on the ground.'

κόστος, Saussurea Lappa 9. 7. 3. in list of ἀρώματα

κότινος (? = ἀγριέλαιος), wild olive, Olea Oleaster

1. 4. 1. more fruitful than ἐλάα; 1. 8. 1. many knots; 1. 8. 2. more knots than ἐλάα; 1. 8. 3. knots regular: knots opposite; 1. 8. 6. liable to excrescences; 1.14.4. wild form of ἐλάα; 2.2.11. cannot be made into ἐλάα by cultivation: effect of transplanting and removing top-growth; 2. 3. 1. occasionally changes to έλάα spontaneously; 3. 2. 1. produces more fruit than έλάα but ripens less; 3. 6. 2. knots opposite; 3. 15. 6. size of fruit of κράταιγος comp.; 4. 4. 11. Indian olive between κ. and λάα; 4. 13. 1. longer-lived than λάα; 4. 13. 2. story of a very old κ. at Olympia; 4. 14. 12. suffers less than λάα from special winds; 5. 2. 4. story of a tree at Megara; 5. 3. 3. character of wood; 5. 4. 2. wood proof against decay; 5. 4. 4. wood not eaten by teredon; 5. 7. 8. uses of wood for carpenter's tools.

κουκιόφορον (= κόιξ), doum-palm, Hyphaene thebaica

2. 6. 9. (not named) described; 4. 2. 7. described. κράνεια (fruit κράνεον 4. 4. 5.),

cornelian cherry, Cornus

1.6.1. core hard and close; 1.8.2. has more knots than θηλυ-κρόνεια; 3.2.1. fruit sweeter and better ripened in wild than in cultivated form; 3.8.1. tree of mountain and plain; 3.4.2. time of budding; 3. 4. 3. time of fruiting; 3. 6. 1. slow-growing (?); 3. 12. 1-2. described; 4. 4. 5. fruit of an unnamed Indian tree (see App. (6)) comp.; 5. 4. 1. more fruitful than θηλυκράνεια; 5. 6. 4. wood very strong.

κραταίγονος, willow-weed, Polygonum Persicaria

num Persicaria
9. 18. 6. properties: described.
κράταιγος (= κραταιγών 3. 15. 6.),
Crataegus Heldreichii

3. 15. 6. described: perhaps a

wild form of μεσπίλη.
κρηπίε, ox-tongue, Helminthia echi-

oeides
7. 8. 3. leaves on the stalk.

κριθή, barley, Hordeum sativum

1. 6 5. roots numerous; 1. 6. 6. do.; 1. 11. 5. each seed separately attached; 2. 2. 9. said to turn sometimes into wheat; wheat; 2. 4. 1. wild « turns into cultivated with cultivation; 4. 4. 9. India has a corresponding cereal and a wild form of κ ; 8. 1. 1. in list of cereals; 8. 1. 3. sown time of germination in Hellas (and in Egypt?); 8. 2. 1. germination described; 8. 2. 3. single leaf first appears: roots described; 8. 2. 6. time of maturing seed; 8. 2. 7. time of harvest in Hellas and in Egypt; 8. 2. 9. crop very early in island of Chalkia; 8.3.2. stem; 8.4.1–2. comp. in detail with other cereals: kinds distinguished distinguished (see below); 8.6.1. conditions for sowing; 8.6.4. suitable soil; 8.6.5–6. rain hurtful when κ . is in flower: and when it is ripe; 8. 7. 1. said to change into aipa under certain conditions; 8.7.5. in many places comes up again next year; etc.; 8. 8. 2. favour-able localities; 8.8.3. αἰγίλωψ (2) grows specially among k.; 8.9.1. grows specially among κ_i ; 8.9.1. exhausts the soil, but less than $\pi\nu\rho\delta$: reason; 8. 10. 2. wheat-rust; 8. 10. 3. effects of weather; 8. 11. 1. seed keeps less well than $\pi\nu\rho\delta$; 8. 11. 3. grain stored without drying; 8. 11. 7. at Babylon grain jumps on the threshing-floor: reason: 9. 11. 9. $\pi\thetai$. floor: reason; 9. 11. 9. πιθύ-μαλλος ὁ μυρτίτης gathered at time of barley-harvest; 9. 12. 4. μήκων ή ροιάς grows in fields

of K.

κριθαὶ ai ἀγρίαι (Indian), Sorghum halevense

4. 4. 9. can be used for bread. κριθαὶ αἰ 'Αχιλλεῖαι, barley, Hordeum

sativum var. 8. 4. 2. ear close to leaf; 8. 10. 2.

s. 4. 2. ear close to leat; 8. 10. 2. specially liable to wheat-rust. κριθαί αἰ Ἰνδικαί, barley, Hordeum sativum var.

8 4. 2. branching.

κριθών γένυς τρίμηνον, barley, Hordeum sativum var.

8. 1. 4. sown late.

κρίνον (= κρινωνία, cf. λείριον (1)), lily, Lilium candidum etc. 1.13.2. has a 'twofold' flower;

1. 13. 2. has a 'twofold' flower; 2.2.1. propagation from exudation; etc.; 4. 8. 6. an unnamed Egyptian plant (see App. (18)) comp.; 4.8.9. petals of flower of λωτός (2) comp.; 6. 6. 3. several colour forms; 6. 6. 8. do.; a coronary plant: described: propagation; 6. 6. 9. leaves of νάρκισσος (1) comp.; 6. 8. 3. flowering time; 9. 1. 4. cf. 2. 2. 1.

κρίνον τὸ πορφυροῦν, Turk's cap lily, Lilium chalcedonicum

6. 6. 3. (see κρίνον). κριοί, see ἐρέβινθος.

κρόκος, crocus, Crocus spp. etc. (see below)

6. 6. root fleshy; 1. 6. 7. do.
 6. 11. large fleshy root;
 7.7.1. leaf of τραγοπάγων comp.;
 7. 4. flowering time short: three kinds mentioned, εὐουμος, λευκός, ἀκαυθώδης (see below).
 7.9.4. root acorn-shaped; 7.10 2. flowers in winter;
 7.13.1. leaves described;
 7. 13.2. no stem except flower-stem.

κρόκος ὁ ἀκανθώδης (= κνῆκος = κ. ἡ ῆμερος), safflower, Carthamus tinctorius

7. 7. 4. (see κρόκος).

κρόκος ὁ ενοσμος, saffron crocus, Crocus sativus

4. 3. 1. abundant in Cyrenaica; 6. 6. 5. sweetest-scented at Cyrene; 6. 6. 10. a coronary plant: described: propagation; 6. 8. 3. flowering time: a wild (scentiess) and a cultivated

kind; 7. 7. 4. see κρόκος; 9. 7. 3. in list of ἀρώματα,

κρόκος ο λευκός, crocus, crocus cancellatus

7. 7. 4: 7. 10. 2. (see κρόκος). κρομυογήτειον, onion, Allium Cepa

4. 6. 2. root of φῦκος τὸ πλατύ comp.

κρόμυον, onion, Allium Cepa 1. 5. 2. 'bark' in layers; 1. 6. 7. roots; 1. 10. 7. attachment of leaves; 1. 10. 8. leaves hollow; 7. 1. 7. stem single; 7. 2. 1. propagation; 7.2.3. growth of γήθυον and πράσον comp.: off-sets specially numerous; 7. 3. 4. seed borne at top; 7. 4. 7-10. kinds distinguished, Σάρδιον, Κνίδιον, Σαμοθράκιον, σητάνιον, σχιστόν, 'Ασκαλώνιον: cultivaσχιστόν, 'Ασκαλώνιον: cultiva-tion and special points of σχιστόν (see below), 'Ασκαλώνιον: further local varieties; 7. 4. 12 formation of roots of σκόροδον contrasted; 7.5.1. likes water; 7.5.2. said to dislike rainwater; 7.8.2. stem smooth, not branched; 7.9.4.cf. 1.6.7; 7. 13. 4. grows in colonies because of offsets; 9. 15. 7. root of μῶλυ comp.

κρόμυον τὸ σχιστόν, shallot, Allium Cepa var.

4. 7-10. distinguished from other varieties of κρόμυον: cultivation.

κρότων, castor-oil plant, Ricinus communis

1. 10. 1. leaves change shape with age of plant; 3. 18. 7. do.

κύαμος, bean, Vicia Faba 3. 13. 3. Size of fruit of κέρασος comp.; 3. 15. 3. fruit of $\tau \epsilon \rho \mu \iota \nu \theta$ os comp.; 3. 17. 6 size of berry of αμπελός (2) comp.; 4. 3. 1. size of fruit of λωτός (4) comp.; 7. 3. 1. length of flowering of ωκιμον comp.; 8. 1. 1. in list of pulses; 8. 1. 3-4, sown early, but can be sown late; 8. 1. 5. time of germination: very slow; 8. 2. 1. germination described; 8. 2. 3. comes up

with several leaves: roots and side-growths contrasted with other pulses; 8.2.5. flowering time; 8.2.6. time of maturing seed; 8.3.1. leaf; 8.3.2. stem; 8.5.1. more than one kind: white form sweetest; 8. 5. 4. attachment of seed; 8. 6. 1. rain attachment of seed; 8.6.1. rain not beneficial after sowing; 8.6.5. likes water when in flower, but not later; 8.7.2. makes ground fertile for έρέβιθος; 8.8.6 causes etc. of & becoming 'cookable' or 'uncookable'; 8.9.1 improves the soil (cf. 8.7.2.); 8.10.5. Infested by lues; 8.11.1. seed does not keep; 8.11. 3. seed keeps well in some localities. keeps well in some localities.

κύαμος (ὁ Λίγύπτιος), Nelumbium spe-

4. 8. 7-8 described; 4. 8. 9. stalk. leaves and growth of fruit of λωτός (2) comp.

κυδώνιος (fruit μήλον κυδώνιον) (= στρουθίον (1)), quince, Cydonia vulgaris

2. 2. 5. produced from seed of στρούθιον; 4. 8. 11. size of root of λωτός (2) comp. to μήλον κυδώνιον. κυϊέ, ?

7.13. 9. (in defective sentence): belongs to τὰ βολβώδη.

κυκλάμινος, cyclamen, Cyclamen graecum

7. 9. 4. root has 'bark'; 9. 9. 1. root and juice used; 9. 9. 3. use in medicine and as charm; 9. 18. 2. leaf of σκορπίος (3) comp.

κύμινον, cummin, Cuminum Cyminum

1. 11. 2. seeds naked; 7. 3. 2-3. seeds described: popular belief about sowing; 7. 4. 1. several kinds; 8. 3. 5. seed very abundant and small; 8. 6. 1 rain not beneficial after sowing; 8. 8. 5. a plant parasitic on root (see App. (25)); 8. 10. 1. diseases; 9. 8. 8. cf. 7. 3. 2-3. κυνόρροδον, dog-rose, Rosa canina

4. 4. 8. an unnamed Indian tree (cotton plant) comp.

κυνόσβατος, wild rose, Rosa sempervirens

3. 18. 4. described: 9. 8. 5. superstition as to method of cutting. κύνωψ, rib-grass, Plantago lanceolata

7.7.3. time of growing: 7.11.2. (?) flowers in a spike.

κυπάριττος, cypress, Cupressus

sempervirens

1. 5. 1. erect and tall: 1. 5. 3. wood not fleshy; 1.6.4. shallow rooting; 1.6.5. roots not branching; 1.8.2. 'male' has more knots than 'female' 1. 9. 1. growth chiefly upwards; 1. 9. 3. evergreen; 1. 10. 4. leaves fleshy; 2. 2. 2. propagation; 2. 2. 6. seed of 'female' produces 'male' trees; 2.7.1. dislikes manure and water; 3. 1. 6. comes up spontaneously in Crete; 3. 2. 3. evidence that it is really wild (at least 'male' form); 3. 2. 6. characteristic of the Cretan Ida; 3. 12 4. bark of κέδρος (1) comp.; 4.1.3. grows very high on Cretan mountains; 3. 1. grows in Cyrenaica;
 4. 5. 2. abounds in Crete, Lycia, Rhodes;
 4. 18. 12. beer (βρυτός) made from k. in Egypt; 4,16.1. some think topping fatal; 5.3.7. θύον comp.: images made from the wood; 5. 4. 1. the 'male' form the more fruitful; 5. 4. 2. wood proof against decay: an instance: takes a fine polish; 5, 7, 4, use of wood in housebuilding. *

κύπειρον, Cyperus rotundus 9. 7. 3. in list of ἀρώματα.

- κύπειρος, galingale, Cyperus longus
 - 1.5.3. stem very smooth; 1.6.8. a stout root and also fibrous roots; 1.8.1. no knots; 1.10.5. leaves end in a point; further described; 4.8.1. in list of $\tau \hat{a}$ λοχμώδη; 4. 8. 12. leaves of μαλιναθάλλη comp.; 4. 10. 1. in list of plants of Lake Copals; 4. 10. 5. described; 4. 10. 6. grows both on land and in water: grows on the floating islands of Lake Copais:

11. 12. foliage of some κάλαμοι

κύτινος, δεε ρόα.

κύτισος (1), laburnum, Laburnum

vulgare

1. 6. 1. core hard and close; 4. 4. 6. habit of ἐβένη comp.; 5. 3. 1. wood of the core very close and heavy.

κύτισος (2), tree-medick, Medicago arborea

4. 16. 5. dangerous to trees.

κώμακον, Ailanthus malabarica 9. 7. 2. an Arabian ἄρωμα (i.e. imported through Arabia . mixed with other ἀρώματα:

[name also given to fruit of a different plantl.

κώνειον, hemlock, Conium maculatum

1.5.3. stem fleshy; 6.2.9. belongs to 'ferula-like' plants: has a to 'ferula-like' plants: has a hollow stem; 7. 6. 4. leaf of ορειοσέλινον comp.; 9.8.8. most powerful juice from root; 9. 15. 8. localities; 9. 16. 8. medical experience; 9. 16. 9. treatment in Keos; 9. 20. 1. πέπερι an antidote to κ .

κωνόφορος, εεε [πεύκη ή] κωνόφορος

λάθυρος, Lathyrus sativus
8. 3. 1. leaf; 8. 3. 2. stem;
8. 10. 5. infested by 'worms.'

λακάρη (λάκαρα) (= κέρασος ? Macedonian name), bird-cherry, Prunus **a**vium

3. 3. 1. a tree of mountain and plain; 3. 6. 1. slow-growing (?). λάπαθος (λάπαθον), monk's rhubarb,

Rumex Patientia

1. 6. 6. root single; 7. 1. 2. time of sowing; 7. 2. 7-8. root described; 7. 4. 1. only one kind; 7. 6. 1. wild form distinguished (see below); 9. 11. 1. leaf of πάνακες το Χειρώνειον comp.

λάπαθον τὸ ἄγριον, dock, Rumex conglomeratus

7. 6. 1. distinguished from λ. τὸ ημερον; 7.7.2. a λάχανον; needs cooking.

λειμωνία, (? = σκόλυμος 6. 4. 3.). golden thistle, Scolymus hispanicus

thistle-like ' plant: 4. 3. a leaves spinous.

λείριον (1) (= κρίνον q.v.) Madonna lily, Lülium candidum 3. 13. 6. flower of ἀκτῆ has the

heavy scent of λ ; 3. 18. 11. scent of flower of $\sigma \mu i \lambda \alpha \xi$ (2) comp.; 9.16.6.(?) leaf of ἐφήμερον comp.

(2), narcissus, Narcissus λείριον

spp. (see below)
13. 2. flower consists of one 13. 2. flower consists of one 'leaf' only partly divided.
 λείριον, polyanthus narcissus, Nar-

cissus Tazetta 6. 8. 1. flowering time; 7. 13. 4.

grown from seed.

λείριον (τὸ ἔτερον 6.8.3.) (=νάρκισσος (1) 6.6.9.), narcissus. Narcissus serotinus

6. 6. 9. a coronary plant: described: flowering time.

λέμνα, water chickweed, Callitriche verna

4, 10, 1, in list of plants of Lake Copais. λευκάκανθα, milk-thistle, Silybum

marianum

6. 4. 3. a 'thistle-like' plant.

λεύκη, abele, Populus alba 1. 10. 1. leaves change shape with age of tree: leaves inverted in summer; 3. 1. 1. propagation; 3. 3. 1. tree of mountain and plain; 3. 4. 2. time of budding; 3. 6. 1. quickgrowing; 3. 14. 2. described; 3. 18. 7. cf. 1. 10. 1; 4. 1. 1. likes wet ground; 4. 2. 3 stem of συκή ή Κυπρία comp.; 4.8.1. grows partially in water; 4.8.2. scarce on Nile; 4.10.2 flower of έλαίαγνος comp.; 4.13.2. shorter-lived by water; 4. 16. 3. instance of a tree which grew again after falling down; 5. 9. 4. wood makes an evil smoke when burnt for char-

 $\lambda \epsilon \nu \kappa \delta i o \nu (1) (= i o \nu \tau \delta \lambda \epsilon \nu \kappa \delta \nu = i \omega \nu i \alpha$ ή λευκή), gilliflower, Matthiola incana

λευκόϊον (2), snowdrop, Galanthus nivalis

7. 13. 9. (in defective sentence) belongs to τὰ βολβώδη.

λιβανωτίς 9.9 5. medicinal use; 9.11.10-11.

two kinds (see below).

λιβανωτὶς ἡ ἄκαρπος, Lactuca graeca 9. 11. 10-11. described: medicinal use: habitat.

λιβανωτίς ή κάρπιμος, (fruit κάχρυ 9. 11. 10.), Lecokia cretica

9. 11. 10. described: medicinal use: habitat: prevents moth in clothes.

λιβανωτός, (gum λίβανος, frankin-cense: also λιβανωτός 9. 4. 4-9. etc.), frankincense-tree, Bos-

wellia Carteri

4. 4. 14. in list of oriental ἀρώματα; 9. 1. 6. time of tapping; 9. 4. 1. collection of gum; 9. 4. 2. Arabian: described: habitat; 9.4.3. another account; 9.4.4-10. accounts of travellers; 9. 11. 3 scent of πάνακες το Ηράκλειον comp.; 9. 11. 10. scent of root of λιβανωτίς ή κάρπιμος comp.; 9. 20. 1. an antidote to κώνειον.

λίνον, flax, Linum usitatissimum 3. 18. 3. seeds oily; 8. 7. 1. said

to change into aipa. λίνον πύρινον

9. 18. 6. growth of κραταίγονος comp. λινόσπαρτον, Spanish broom, Spar-

tium junceum

1. 5. 2. bark in layers. λυχνίς, rose-campion, Lychnis coron-

6, 8, 3, a coronary plant: flowering time.

(λωτός) 7. 15. 3. many plants called by this name which have nothing in common but the name.

λωτός (1), nettle-tree, Celtis australis

1. 5. 3. wood not fleshy; 1. 6. 1. core hard and close; 1. 8. 2. few knots; 4. 2. 5. colour of wood of περσέα comp.; 4. 2. 9. wood of olive of Thebaid comp.; 4. 2. 12. wood of an unnamed tree (? κοκκυμηλέα ή Αίγυπτία)

comp.; 5. 3. 1. wood very close and heavy; 5. 3. 7. images made from the wood; 5. 4. 2. wood proof against decay: turns black when old; 5. 5. 4. core not obvious but exists; 5. 5. 6. treatment of core for making door-hinges; 5. 8. 1. grows in some places exceptionally fine.

λωτός (2) (aquatic) (root κόρσιον), Nile water-lily, Nymphaea stel-

4. 8. 9-11. described.

λωτός (3) (herb), trefoil, Trifolium

fragiferum
7. 8. 3. leaves 'on the stem';
7. 13. 5. seed sometimes takes two years to germinate.

λωτός (4) (Libyan tree), Zizyphus Lotus

4 3.1. common in Libya; 4.3.1-2. described; 4. 3. 4. further described.

λωτός (5) (aromatic) (= μελίλωτος), Trigonella graeca

9. 7. 3. in list of ἀρώματα.

μαγύδαρις (= ἱππομάραθον), Prangos ferulacea

 6. 12. root most characteristic part; 6. 3. 7. distinct from σίλφιον: described: distribu-

[6. 3. 4. name also given to seed of σίλφιον].

μαδωνάϊς (= νυμφαία 9.13.1.), yellow water-lily. Nuphar luteum μαλάχη (1), mallow, Lavatera ar-

borea

1. 3. 2. a herb which becomes tree-like under cultivation; 1. 9. 2. do.; 4. 15. 1. outer bark can be stripped; 9. 18. 1. leaf fruit and taste of stem of αλθαία comp.

μαλάχη (2), cheese-flower, Malva

silvestris

 7. 7. 2. a λάχανον; needs cooking: 7. 8. 1. stem on the ground μαλάχη (3) ἡ ἀγρία (= ἀλθαία 9.15.5.), marsh-mallow, Althaea officinalis

μαλιναθάλλη (= μνάσιον), Cyperus esculentus 8. 12. described.

μανδραγόρας (1), mandrake, Mandra-

gora officinarum 9. 8. 8. superstition as to gathering; 9. 9. 1. root and juice used: medicinal use of leaf and root.

μανδραγόρας (2), ? deadly night-shade, Atropa Belladonna 6. 2. 9. belongs to 'ferula-like'

plants: has hollow stem: fruit

described.

μάραθον, fennel, Foeniculum vulgare 1. 11. 2. seeds naked; 1. 12. 2. taste of sap; 4. 6. 3. φῦκος τὸ τριχόφυλλον comp.; 6.1.4. a spineless wild under-shrub: belongs to 'ferula-like' plants; 6. 2. 9. do.: has a fibrous stem; 7. 3. 2. seeds described; 9. 9. 6. leaf of θαψία comp.

μάσπετον, εεε σίλφιον

μελαγκρανίς (= σχοίνος ο κάρπιμος 4. 12. 1.), bog-rush, Schoenus nigricans

μελάμπυρον (μελάμπυρος), Neslia

paniculata

8. 4. 6. infests πυρὸς ὁ Σικελός: contrasted with αίρα; 8. 8. 3. (μελάμπυρος ὁ Ποντικός), speci-

3. 6. 1. slow growing (?); 3. 6. 5 roots numerous matted and run deep, according to Arcadians; 3. 11. 3-4. described; two kinds, see βουμέλιος; 3.17.1. leaf of φελλός comp.; 4. 5. 3. grows in Pontus; 4. 8. 2. grows in Pontus; 4. 8. 2. common on Nile; 5. 1. 2. time of cutting timber; 5. 6. 4. wood moist': used for elastic bedsteads; 5. 7. 3. wood used for bent-wood work: use in shipbuilding; 5. 7. 8. uses of wood for carpenter's tools.

μελίλωτος (= λωτός (5)), Trigonella graeca

7. 15. 3. one of the many diverse plants called λωτός. μέλινος, (in other authors μελίνη:

see 8. 1. 1. n.), Italian millet, Setaria italica

8. 1. 4. sown later than cereals and pulses; 8. 2. 6. time of maturing seed; 8. 3. 2. stem; 8. 3. 3. flower; 8. 7. 3. needs little water: comp with κέγχρος.
μελισσόφυλλον, balm, Melissa offici-

nalis 6. 1. 4. a spineless wild undershrub.

μεμαίκυλον, 800 κόμαρος

μεσπίλη (fruit μέσπιλον) (= μ. σατάνειος), medlar, Mespilus germanica

3. 12. 5-6. described: three kinds (Idaean account, see below); 3. 13. 1. leaf of képagos comp. 3. 15. 6. leaf, bark and taste of fruit of κράταιγος comp.; 3.17.5. flower of συκή ή Ίδαία comp. also taste of fruit; 4.2.10, fruit of κοκκυμηλέα comp.; 4.8.12. μαλιναθάλλη comp.; 4. 14. 10. fruit gets worm-eaten.

μεσπίλη ή ανθηδονοείδης, hawthorn,

Crataegus Oxyacantha

3. 12. 5. described.

μεσπίλη ή ἀνθήδων, oriental thorn, Crataegus orientalis

3. 12. 5. described.

μεσπίλη ή σατάνειος, medlar, Mespilus germanica 3. 12. 5. described.

Μηδική (πόα), lucerne, Medicago sativa

8, 7, 7, destroyed by sheep sleeping on it.

μήκων, poppy etc., Papaver spp. etc. (see below)

1. 9. 4. evergreen; 1. 11. 2. seeds in a vessel; 4. 8. 7. size of flower of κύαμος ο Αλγύπτιος comp.; 4. 8. 10. size of 'head' of λωτός (2) comp.; 4.10.3. σίδη comp. (?); 9.8.2. juice of 'head' collected; 9. 12. 3-5. kinds (see below) having nothing in common but the name; 9. 16. 9. medical experience; 9. 20. 1. seeds of one kind of mémepi comp.

μήκων ή 'Ηρακλεία (= 'Ηρακλεία), Silene venosa

9. 12. 5. described: medicinal 1180.

μήκων ή κερατίτις, horned poppy, Glaucium flavum var. Serpierii 9. 12. 3. described: medicinal use:

habitat.

μήκων ή μέλαινα, Papaver Rhoeas 9. 11. 9. mixed with τιθύμαλλος ο μυρτίτης to make a medicine. μήκων (ή οπώδης), opium poppy,
Papaver somniferum

1. 12. 2. juice.

μήκων ή ροιάς, Papaver hybridum 9. 12. 4. described: edible: habitat: medicinal use.

μηκώνιον (= τιθύμαλλος 9. 8. 2.). spurge, Euphorbia Peplus 9. 8. 2. collection of juice.

μηλέα (fruit μήλον), apple, Pyrus

Malus

1. 3. 3. a tree whose stem is not single; 1. 5. 2. bark smooth: bark readily drops off; 1. 6. 1. core fleshy; 1. 6. 3. few roots; 1. 6. 4. shallow rooting; 1. 8. 4. knots peculiar; 1. 9. 1. trunk divides low down; 1. 10. 4. (?) leaves fleshy; 1. 10. 5. leaves oblong; 1. 11. 4. seeds all together in a single case; 1. 11. 5. seeds in a membrane: 1. 12. 1. taste of fruit; 1. 12. 2. taste of sap; 1. 13. 1. flower 'leafy'; 1. 18. 3. flower above fruit-case; 1. 14. 1. bears on last year's wood: some kinds hast year s, wood: some kinds bear also on new wood; 1.14.4, many cultivated forms; 2.1.2. propagation; 2.2.4, degenerates from seed; 2.2.5. seed produces wild form; 2.5. 8. grafting; 2.5.6. trees should be planted fairly close together; 2.6.6. some dates round like $\mu \hat{\eta} \lambda a$; 2. 8. 1. apt to shed immature fruit; 3. 3. 1. tree of mountain and plain; 3. 3. 2. has better fruit and timber in lowlands; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 3. 11. 5. mountain and lowland forms compared; 4. 5. 3. abundant in Pontus; 4. 5. 4. grows on Mount Tmolus and Myslan Olympus; 4. 7. 7. size of fruit of δένδρον τὸ ἐριόφορον comp.; 4. 10. 2 leaf of έλαιαγνος comp. (?); 4. 10. 3.

size of flower of σίδη comp.; 4. 13. 2. short-lived, especially certain kinds; 4. 13. 3. after decaying shoots again from same stock; 4. 14. 2. apt to get worm-eaten; 4. 14. 10 fruit gets worm-eaten; 4. 14. 12. un-injured by special winds; 4.16.1. survives splitting of stem; 5.3.3. character of wood; 5, 4, 1, the less fruitful trees produce more solid wood; 6. 4. 9 'head' of iξίνη comp. to μήλον.

μηλέα ή γλυκεία, Pyrus Malus var.? 4. 13. 2. specially short-lived; 4. 14. 7. has specially weak constitution; a form of μ. ή ἐαρινή; 9. 11. 5. leaf of στρύχνος ὁ ὑπ-

νώδης comp.

μηλέα ή εαρινή, Pyrus Malus var.? 1. 3. propagation; 4. 7. 7. size of cotton-bearing vessel comp.; 4 13. 2. specially short-lived; 4. 14. 7. has weak constitution;

(cf. μ. ή γλυκεία). μηλέα ή όξεια, Pyrus Malus var.? 4. 13. 2. comparatively long-lived. μηλέα ή Περσική (Μηδική) citron, Cit-

rus Medica

1. 11. 4. seeds in a row; 1. 18. 4. only pistillate flower fruitful; 4. 4. 2. peculiar to Media and Persia: described.

μήλον το Κυδώνιον, see Κυδώνιος μήλωθρον (= ἄμπελος (4)), bryony, Bryonia cretica

3. 18. 11. fruit of σμίλαξ (2) comp.; 6. 1. 4. a spineless wild under-

shrub. μήνανθος, Limnanthemum nymphoide**s**

10. 1-2. in list of plants of Lake Copais; 4. 10. 4. requires further investigation.

 $\mu i \lambda a \xi (= \sigma \mu i \lambda a \xi (2)), smilax, Smilax$

aspera

1. 10. 5. leaf described; 1. 10. 6. leaf with spinous projections; 6. 8. 3. flower used in garlands.

μίλος, yew, Taxus baccata
1. 9. 3. evergreen; 3. 3. 1. a
mountain tree; 3. 3. 3. evergreen; 3. 4. 2. time of budding; 3. 4. 5. time of flowering and fruiting; 3. 4. 6. time of fruiting; 3. 6. 1. slow growing (?) 3. 10. 2. described; 4. 1. 3. likes shade; 5. 7. 6. uses of wood.

μίνθη (μίνθα) (= ἡδυόσμον), green mint, Mentha viridis

2. 4. 1. σισύμβριον turns into μ. unless often transplanted; 6.7.2. said by some to have no fruit.

μνάσιον (= μαλιναθάλλη), Cyperus

esculentus

4. 8. 2. used for food in Egypt; 4. 8. 6. described.

μυάκανθος (= κεντρομυρρίνη), but-cher's broom, Ruscus aculeatus 6. 5. 1. in list of spinous plants which have leaves as well as

μύκης, mushroom etc., Fungus
1. 1. 11. has not all the 'parts' of a plant; 1. 5. 3. stem very smooth; 1. 6. 5. no roots: 3. 7. 6. grows on roots of trees.

[4. 7. 2. marine growths which turn to stone];

[4.14.3. name given to a disease of exac.

μυόφονον (= ἀκόνιτον = θηλύφονον == σκορπίος (3)), wolf's bane, Aconitum Anthora

1. 4. a spineless wild under-shrub: belongs to 'ferula-like' plants; 6. 2. 9. do.; has a fibrous stem.

μυρίκη (1), tamarisk, Tamarix te-

trandra

1. 4. 3. 'amphibious'; 1. 9. 3. evergreen; 1. 10. 4. leaves fleshy; 3. 3. 1. tree of mountain and plain; 3. 3. 3. evergreen; 3. 16. 4. bark of κόμαρος comp.; 4. 2. 6. (?) leaf of βαλανος comp.; 4. 6. 7. leaf of δρῦς (7) comp; 6. 2. 1. leaf of kvéwpos à médas comp.; 6. 4. 8. flower of yanaiλέων comp.

μυρίκη (2), tamarisk, Tamarix articulata

5.4.8. Arabian : wood very strong. μυρρίνη (μύρρινος, μύρτος) (fruit μύρτον), myrtle, Myrtus communis

1. 3. 3. effect of not pruning; 1. 9. 3. evergreen; 1. 10. 2. leaves close-set and opposite; 1. 10. 4. leaves narrow; 1. 10. 8.

465

н н

THEOPH. VOL. II.

leaves regular; 1. 12. 1. taste of fruit; 1. 13. 3. flower above fruit-case; 1. 14. 1. bears on last year's wood: flowers borne on new wood not fertile; 1.14.4. many cultivated forms; 2. 1.4. propagation; 2.2.6. sometimes improves from seed; 2. 5. 6. propagation: trees should be planted close together; 2. 7 2. needs much pruning; 2. 7. 3. requires pungent manure and much water; 3.6 2. formation of buds; 3.12.4. fruit of κέδρος (1) comp.; 3. 15. 5. leaf of πύξος comp.; 3. 16. 4. flower of κόμαρος comp.; 4. 2. 6. (?) leaf of βάλανος comp.; 4. 3. 1. arrangement of fruit of λωτός (4) comp. 4. 5. 3. does not thrive in cold regions; 4. 5. 4. grows in Propontis; 5. 8. 3. grows in low-land parts of Latium: and on Circelan promontory (a dwarf kind); 6. 8. 5. very fragrant in Egypt; 9. 11. 9. leaf of τιθύ-

μαλλος δ μυρτίτης comp. μῶλυ, moly, Allium nigrum 9. 15. 7. localities in Arcadia:

said to be like the μ . of Homer: described: use as charm.

ναῖρον 9. 7. 3. in list of ἀρώματα. νᾶπν, white mustard, Brassica alba 1. 12. 1. taste of fruit; 7. 1. 2-3; time of sowing and of germina-tion; 7. 3. 2. seeds described; 7. 5. 5. seed keeps well. νάρδον, spikenard, Nardostachys Ja-

tamansi

9. 7. 2. an Indian ἄρωμα; 9. 7. 3. in list of ἀρώματα; 9. 7. 4. an unnamed Thracian plant (see App. (25)) comp.

ναρθηκία (= νάρθηξ see 6.2.7.), ferula. Ferula communis

6.1.4 spineless: belongs to 'ferulalike, plants; 6. 2. 7. perhaps. differs only in size from νάρθηξ; 6. 2. 8. described.

 $v\acute{a}\rho\theta\eta E$ (= $va\rho\theta\eta\kappa ia$ see 6, 2, 7.), ferula,

Ferula communis

 1. 2. 7. flesh turns to wood; 1.6.1. core fleshy; 1. 6. 2. core membranous; 6.2.7. perhaps differs only in size from $\nu a \rho \delta \eta_{\kappa}(\alpha)$ δ . 2. 8. described; 6. 3. 1. stall of $\sigma(\lambda \phi \omega \nu \text{ comp.}; 9.9.6. \text{ stem})$ of $\theta a \psi (\alpha \text{ comp.}; 9.10.1. leaf of both <math>\delta \lambda \lambda \delta \rho \omega \rho \omega$ comp. by some; 9. 16. 2. δίκταμνον kept έν νάρθηκι.

 ν άρκισσος (1) (= λ είριον (2) 6, 6, 9,), narcissus, Narcissus serotinus

6.6.9. a coronary plant: described; 7.13.1. leaves described; 7.13.2. no stem except the flower-stem; 7. 13. 5-7. stem appears before leaves, viz. flower-stem: sequence described and comp. with σκίλλα.

νάρκισσος (2), pheasant's eye nar-cissus, Narcissus poeticus

6. 8. 1. flowering time.

νάρτη 7. 8. in list of ἀρώματα. (νηπενθές) = μήκων ή οπώδης, opium poppy, Papaver somniferum.
9. 15. 1. mythical: supposed

effects.

νυμφαία (= μαδωνάϊς 9.13.1.), yellow water-lily, Nuphar luteum

13. 1. fragrant: habitat and localities: leaf described: medicinal use: called μαδωνάϊς in Boeotia.

ξίρις, gladwyn, Iris foetidissima 9.8.7. superstition as to gathering. ξίφιον $(=\xi i \phi \cos 7.13.1. = \phi a \sigma \gamma \alpha \nu \nu)$,

corn-flag, Gladiolus segetum 6. 8. 1. flowering time; 7. 13. 2. flower-stem not the only stem. $\xi i \phi o s (= \xi i \phi i o v 7, 13, 1. = \phi a \sigma y a v o v),$ corn-flag, Gladiolus segetum.

ογχνη, wild pear, Pyrus communis var. Pyraster

2. 5. 6. trees should be planted rather far apart.

ŏη (οἴη), sorb, Sorbus domestica

2. 2 10. becomes sterile in a warm place; 2. 7. 7. 'correcting' the tree; 3. 2. 1. fruit sweeter and better ripened in wild than in cultivated form: 3, 5, 5, winterbuds; 3. 6. 5. roots shallow but strong: thick according to Arcadians; 3 11. 3. leaf of μελία

comp.; 3. 12. 6-9. described; 3. 15. 4. leaf of τέρμινθος comp. οινάνθη (1), drop-wort, Spiraea Filipendula

6. 6. 11. a coronary plant: grown from seed;
 6. 8. 1-2. flowering

time: flower described. οινάνθη (2) ή άγρία, wild vine, Vitis

silvestris

9. 6. ἀθραγένη comp.

oloos (= ayvos), withy, Vitex Agnus-3. 18. 1-2. has two forms, 'white'

and 'black'; 6. 2 2. used for tying-up.

ολόσχοινος, εεε σχοίνος ο όλ. ολύρα, (cultural variety of ζειά), rice-wheat, Triticum dicoccum

8. 1. 3. sown early; 8. 4. 1. comp. in detail with other cereals; 8. 9 2. does not exhaust the

soil much: reason. ὀνοθήρας (= δάφνη ἡ ἀγρία), oleander, Nerium Oleander

9. 19. 1. effect on 'mind': described.

ονόπυξος, Onopordon illyricum 6. 4. 3. a 'thistle-like' plant.

ονοχειλές, bugloss, Echium diffusum 7. 10. 3. flowers borne in succes-

ονωνίς, rest-harrow, Ononis antiquorum

6. 1. 3. has leaves as well as spines: a wild under-shrub; 6. 5. 1. in list of spinous plants which have leaves as well as spines; 6. 5. 3-4. described: troublesome to farmers.

δξυάκανθος, cotoneaster, Cotoneaster Pyracantha

1. 9. 3. evergreen; 3. 3. 1. tree of mountain and plain; 3. 3. 3. evergreen; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 4.4.2. thorns of μηλέα ἡ Περσική comp.; 6. 8 3. fruit used in garlands.

δξύη (ὑξύα), beech, Fagus silvatica 3. 3. 8. doubt whether it has a

flower; 3. 6. 5. roots few slender and 'plain' according to Arcadians: shallow - rooting; 3. 10. 1. described; 3. 11. 5. mountain and lowland forms

compared; 5.1.2. time of cutting timber; 5.1.4. do.; 5.4.4. wood does not decay in water; 5. 6. 4. wood 'moist': used for elastic bedsteads; 5. 7. 2. wood used for keel etc.of small vessels; 5. 7. 6. other uses of wood; 5. 8. 8. grows very fine in low-land part of Latium.

οξύκεδρος (= κέδρος (1) 3, 12, 3.), prickly cedar, Juniperus Oxy-

cedrus

3. 12. 3. some, who call ἄρκευθος a κέδρος, distinguish κέδρος (1) as

οπιτίων, ? 7. 13. 9. (in defective sentence)

belongs to τὰ βολβώδη. ὑποβάλσαμον, εςε βάλσαμον.

ορειοσέλινον, parsley, Petroselinum sativum

7. 6. 3-4. distinguished from other forms of σέλινον: medicinal use. ορειπτελέα, wych-elm, Ulmus mon-

 3.14.1. distinguished from πτελέα. ορίγανον (ορίγανος) (= ο. ή μέλαινα), marjoram, Origanum viride etc.

1.9.4. evergreen (partly); 1.12.1. taste of fruit; 6.1.4. a spineless wild under-shrub; 6. 2. 3. two forms, 'black' and 'white' (see below): seed conspicuous: not, like $\theta \dot{\nu} \mu o s$, particular as to situation; 7. 1. 3. time of germination; 7. 1. 6. germination; 7. 2. 1. propagation; 7. 6 1. wild form distinguished.

υρίγανος ή λευκή, marjoram, Origanum heracleoticum

 2. 3. distinguished from δ. ή μέλαινα.

ορίγανος η μέλαινα (= ορίγανον), mar-

joram, Origanum viride 6. 2. 3. distinguished from δ. ή

λευκή. ὅρμινον, Salvia Horminum

 1. 4. sown later than cereals and pulses;
 2. 7. 3. doubtful if eaten green by animals: de-scribed: sown at same time as σησάμη.

δροβάγχη, dodder, Cuscuta europaea 8. 8. 4. grows specially among οροβοι: reason: απαρίνη comp.

οροβος, bitter vetch, Ervum Ervilia 2. 4. 2. more digestible if sown in spring; 7. 5. 4. used to prevent ψύλλαι in ραφανίς; 7.6.3. size of fruit of iπποσέλινον comp.: 8.1.4. sown both early and late; 8.2.5. flowering time; 8. 3. 2. stem; 8. 5. 1. more than one kind: white form sweetest; 8. 5. 2. seeds not in compartments; 8. 5. 3. shape of pod; 8. 8. 4. οροβάγχη grows specially among b.; 8. 10. 1. a pest; 8. 11. 2. seed keeps well; 8. 11. 6. do. specially in hill-country; 9.20.1. shape of one kind of πέπερι (fruit) comp.

ορτυξ (= στελέφουρος according to some, 7. 11. 2.), plantain, Plantago Lagopus.

ορυζον, rice, Oryza sativa

4. 4. 10. described.

ορχις (1) (μέγας), orchis, Orchis papilionacea

9.18.3. properties: leaf and stalk. ὄρχις (2) (μικρός), orchis, Orchis longicruris

9.18.3. properties: leaf and stalk. οστρύα (δστρυίς) (δστρυίς = δστρυς 3.10.3.), hop-hornbeam, Ostrya carpinifolia

than 'female' has more knots than 'female'; 3. 3. 1. tree of mountain and plain; 3. 6. 1. slow-growing(?); 3. 10. 3. described.

οστρυς (= ὀστρύα 3.10.3.), hop-hornbeam, Ostrya carpinifolia 3. 10 3. described.

ουίγγον, Colocasia antiquorum
1.1.7. 'fruit' underground; 1.6.9.
grows underground; 1.6.11. described.

πάδος (? = π ηδός (?)), Prunus Mahaleb 4. 1. 3. likes shade.

παιωνία (= γλυκυσίδη 9.8.6.), peony, Paeonia officinalis

9. 8. 6. superstition as to time of digging. παλίουρος (1), Christ's thorn, Pali-

urus australis

1. 3. 1. a typical 'shrub'; 1. 3. 2. becomes tree-like; 1.5.3. thorns on wood; 1. 10. 6. leaf with spinous projections; 1. 10. 7. stem presently spinous; 3. 3. 1. tree of mountain and plain; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 3. 11. 2. fruit of σφένδαμνος comp.; 3. 18. 3. kinds: described; 4. 8. 1. to some extent grows in marshes; 4.12.4. to some extent aquatic: 6.1.3. has spines on the shoots.

παλίουρος (2) (ὁ Αἰγύπτιος), Zizyphus Spina-Christi

3. 1-2, common in Libya; 4.3.3. described: distinguished from π , of Hellas.

πανάκεια (= πάνακες τὸ Ἡράκλειον). Opopanax hispidus

9. 15. 7. localities.

πάνακες (τὸ Σύριον ? 9.7.2: 9.10.1.), (juice χαλβάνη (?) 9.7.2: 9.9.2., see note), all-heal, Ferulago galbanifera

9. 1. 2. in list of plants whose juice is a gum; 9.7.2. Syrian: χαλβάνη made from π .; 9. 7. 3. in list of ἀρώματα; 9. 9. 1. root fruit and juice used; 9. 9. 2. uses for medicine and perfume; 9. 11. 1. kinds (see below); 9. 11. 4. two further kinds, one fine-leaved, the other not: medicinal use.

πάνακες τὸ 'Ασκληπίειον, Ferula nodos**a**

9. 8. 7. superstition as to gathering: 9. 11, 1. described: medicinal use.

πάνακες τὸ Ἡράκλειον (= πανάκεια),

Opopanax hispidus 11. 1. in list of kinds of w.: 9. 11. 3. described: medicinal

use. πάνακες τὸ Χειρώνειον, elecampane, Inula Helenium

9. 11. 1. described: habitat: medicinal use.

παντάδουσα, star-thistle, Centaurea Calcitrapa

6. 5. 1. in list of spinous plants which have leaves as well as spines.

πάπυρος (stalk πάπυρος), papyrus, Cyperus Papyrus

4. 8. 2. useful for food in Egypt; 8. 8-4. described: uses;

8. 5. stem of σάρι comp;
 3. 1. belongs to 'ferula-like'

plants. παρθένιον, bachelor's buttons, Pyre-thrum Parthenium

 7. 2. a λάχανον: needs cooking πέζις, bullfist, Lycoperdon Bovista 1. 6. 5. no roots.

πελεκίνος, axe-weed, Securigera Coro-

nilla 8. 8. 3. grows specially among

άφάκη: name explained. πενταπετές (= πεντάφυλλον 9.13.5.), cinquefoil, Potentilla reptans

13. 5. described.

πεντάφυλλον (= πενταπετές 9. 13. 5.), cinquefoil, Potentilla reptans.

(πέπερι), pepper, Piper nigrum 9. 20. 1. a fruit: two forms: de-

scribed: properties: antidote to κώνειον; 9. 20. 2. size of Κνίδιος κόκκος comp.
περδίκιον, 'partridge-plant,' Polygonum maritimum

1. 6. 11. large fleshy roots.
περιττός (? στρύχνος ὁ περιττός) (=
στρύχνος ὁ μανικός θ. 11. 6.),
thorn-apple, Datura Stramonium. περσέα (= πέρσιον), Mimusops Schim-

3. 3. 5. not fruitful everywhere; 4. 2. 1. peculiar to Egypt; 4. 2. 5. described; 4. 2. 8. com-

mon in Thebaid. πέρσιον (= περσέα), Mimusops Schim-

2. 2. 10. effects of climate.

πευκέδανον, sulphur-wort, Peuceda-num officinale

9. 14. 1. how long drug will keep;
9. 15. 1. grows in Arcadia;
9. 20. 2. properties of root: use in medicine: grows in Arcadia.

πεύκη, fir, Pinus spp.
1. 3. 6. refuses cultivation; 1.5. 1. 3. 6. refuses cultivation; 1. 5. 1. erect and tail; 1. 5. 4. wood has many knots; 1. 6.1. core woody; 1. 6. 3. root single; 1. 6. 5. roots not branching; 1. 8. 1. many knots; 1. 9. 3. evergreen (the wild and one cultivated kind); 1. 10. 4. leaves like teeth of comb (?); 1.10 6. leaf spinous at tip; 1. 12. 1. taste of fruit; 1. 12. 2. taste of sap; 2. 2. 2. propagated only by seed; 2. 5. 2. instance of very long roots; 3. 1. 2. grows only from seed: 3. 2. 3. evidence that it is really wild; 3. 3. 1. a mountain tree; 3. 3. 8. evergreen; 3. 3. 8. doubt whether it has a flower; 3. 4. 5. time of budding and fruiting; 3. 4. 6. time of fruiting; 3. 5. 1. do.; 3. 5. 5. winter-buds; 3. 5. do.; 3. 5. 5. winter-buds; 3. 5. do.; 3. 5. 6. 1. quick growing; even young tree fruits; 3. 6. 4. not deep-rooting; 3. 7. 1. dies if topped; 3. 7. 3. produces a 'tuft' (xúrrapes); 3. 9. 1-8. kinds according to various authorities (see below): distincauthorities (see below): distinction from $\pi(\tau \nu s)$; 3. 9. 4. timber, foliage; 3. 9. 5. further distinction from $\pi(\tau \nu s)$: the disease 'pitch-glut'; 3.9.7. comparison with λάτη; 3.9.8. do.: core and callus; 4.1.1 likes sun; 4.1.2. in shade has inferior timber; 4.5.1. in list of Northtimber; 4.5.1. in list of North-ern trees; 4.5.3. does not grow in Pontus; 4.15.3. effects of stripping bark at various sea-sons; 4.16.1. topping fatal; 4.16.1-2. not injured by cutting for tar; 4.16.4. said to perish if entirely deprived of its heart-wood; 5.1.2. time of cutting timber; 5.1.4.do.; 5.1.9-10. methods of cleaving; 5.4.2. wood (when resinous) proof wood (when resinous) proof against decay; 5. 4. 4. more eaten by teredon than ελάτη: 5. 4. 8. effect of salt water on different parts; 5. 5. 1. knotty parts of wood hard to work; 5. 6. 1. wood good for struts: 5. 6. 1. wood good for struts: behaviour under pressure; 5.6.2. takes glue best of all woods; 5. 7. 1-2. uses of wood in shipbullding; 5. 7. 4-5. uses in house-building and crafts; 5.8.1. grows to great size in Latium, but finer still in Corsica; 5.8.3. grows in hill-country of Latium; 5. 9. 3. charcoal of this wood preferred by smiths to that of δρύς; 9.1. 2. sap gummy; 9.1.6. time of tapping; 9. 2. 1. pro-

ductive of resin (λητίνη); 9.2.21 quality of resin; 9.2.3-4. Macedonians only burn the 'male' donians only burn the main for pitch (πίττα), and the roots of the 'female'; aspect etc. required for production of good pitch; 9. 2. 5. Idaean account different (see π. ή 'Ιδαία and π. ή περαλία); 9. 2. 6. filling up the holes; 9. 2. 7. do. Idaean account; 9. 2. 8. further rules for collecting pitch: age of tree; the 10. 3. 1.3 method of presents of the second s etc.; 9. 3. 1-3. method of preparing pitch; 9. 1. 4. does not grow in Syria.

πεύκη ή ακαρπος (= π. ή θήλεια = π. ή 'Idaía), Corsican pine, Pinus

Laricio

3. 9. 2. described; 3. 9. 4. one of three wild kinds (Idaean account).

πεύκη ή ἄκαρπος ('male'), Corsican pine, Pinus Laricio

3. 9. 2. comp. with 'female.' πεύκη η ακαρπος ('female'), Aleppo

pine, Pinus halepensis 3.9.2. comp. with 'male.'

πεύκη ἡ ἄρρην (= π . ἡ παραλία = π ίτυς in 3. 9. 5.), Aleppo pine, Pinus halepensis

3. 9. 3. timber: produces συκή (Mt. Ida); 3. 9. 4. one of three wild kinds (Idaean account).

πεύκη ἡ ἡμερος (= [π. ἡ] κωνόφορος), stone pine, Pinus Pinea 3. 9. 1. distinguished from other

kinds; 3. 9. 4. Arcadians say it πίτυς: timber, foliage, resin.

 $\pi \epsilon \acute{\nu} \kappa \eta \dot{\eta} \theta \acute{\eta} \lambda \epsilon \iota a (= \pi. \dot{\eta} \ddot{a} \kappa a \rho \pi o s = \pi. \dot{\eta}$ Idaia), Corsican pine, Pinus

Laricio

3. 9. 3. timber: contains aiyis;
3. 9. 4. one of three wild kinds (Idaean account). πεύκη ή 'Ιδαία (= π. ή ἄκαρπος = π. ή

θήλεια), Corsican pine, Pinus Larició

3. 9. 1-2, described; 9. 2. 5. Idaean account of pitch (πίττα).

[πεύκη ή] κωνόφορος (= π. ή ήμερος), stone pine, Pinus pinea 2.52.6. seeds true; 3.9.4. foliage: pitch (πίττα): Arcadians say it ία α πίτυς.

 $\pi \epsilon \nu \kappa n \dot{n} \pi \alpha \rho \alpha \lambda i \alpha (= \pi \cdot \dot{n} \dot{\alpha} \rho \rho \rho n \nu = \pi i \tau \nu c$ in 3. 9. 5), Aleppo pine, Pinus halepensis

3. 9. 1. described; 9. 2. 5. Idaean account of pitch (πίττα).

πήγανον (πηγάνιον), rue, Ruta graveo-

1. 3. 1. a typical under-shrub; 1.3.4. becomes tree-like; 1.9.4; evergreen; 1. 10. 4. leaves fleshly; 2. 1. 3. propagation; 6. 1. 1. may be classed as an under-shrub; 6. 5. 3. leaf of overis comp.; 6. 7. 3. strong plants of άβρότονον comp.; 7.2.1. propagation: seed slow to germinate; 7. 4. 1. only one kind; 7. 5. 1. dislikes manure; 7. 6. 1. wild form distinguished; 9. 4. 2. colour of leaf of λιβανωτός comp.; 9. 5. 1 leaf of βάλσαμον comp.; 9. 9. 6. leaf of ioxás (απιος (2)) comp.

πηδός (?) (? = πάδος 4. 1. 3.), Prunus

Mahaleb

7. 6. uses of wood. πικρίς, Urospermum picroeides

7. 11. 4. inedible: flowers in spring, but also throughout throughout winter and summer.

πίλος, Polyporus igniarius (?)
3. 7 4. produced by δρῦς; de-

scribed. πισός, pea, Pisum sativum

sos, pea, Pisum sattoum
8. 1. 1. in list of pulses; 8. 1. 4
sown late; 8. 2. 3. comes up
with several leaves; 8. 3. 1.
leaf; 8. 3. 2. stem; 8. 5. 2.
seeds not in compartments; seeds not in compartments; 8. 5. 3. shape of pod; 8. 10. 5. infested by worms.' πίτυς (= πίτυς ἡ ἀγρία = πεύκη ἡ ἀρρην in 3. 9. 5. = πεύκη ἡ παρα-λία in 3. 9. 5.), Aleppo pine,

Pinus halepensis

leaves like teeth of comb; 1. 10. 6. leaf spinous at tip; 1. 12. 1. taste of fruit; 2 2. 2 propagated only by seed; 3.1.2. do.; 3. 3. 3. evergreen; 3. 3. 8. doubt whether it has a flower (κύτταρος); 3. 4 5. time of budding and fruiting; 3.5.5. winterbuds; 3. 6. 1. slow growing (?):

even young tree fruits; 3.9 4-8. πεύκη and πίτυς; 3 9. 6. distinction from πεύκη; 3. 11. 1. bark of σφένδαμνος comp.; 3.17.1. bark of φελλός comp.; 4.5.3. does not grow in Pontus: 4.14.8. if topped, becomes barren, but is not destroyed; 4. 16 1. topping fatal; 5. 1. 2. time of cutting timber; 5. 1. 4. do.; 5. 1. 5-6. timber comp. with $\epsilon\lambda \dot{\alpha}\tau\eta$; 5 7. 1. used in Cyprus for ship building instead of πεύκη; 5. 7. 3. wood used for bent-wood work in triremes; 5. 7. 5. use of wood in shipbuilding and house-building: soon rots; 5. 7. 8. use of wood for carpenter's tools; 5. 9. 2. charcoal of this wood used in silver mines; 9.1.2. sap gummy; 9. 2. 1. production of resin $(\rho m(\nu \eta))$; 9. 2. 2. quality of resin; 9. 2. 3. said to be burnt for pitch (πίττα) in Syria.

πίτυς ή άγρία (= πίτυς = πεύκη ή άρρην = πεύκη ή παραλία in 3.9.5.) Pinus halevensis (mountain

form)

1. 9. 3 evergreen; 3. 3. 1. a mountain tree (Macedonian). πίτυς ἡ φθειροποιός, Pinus brutia

2. 2. 6. seeds come true.

πλάτανος, plane, Platanus orientalis
1. 4, 2. lives near water; 1. 6. 3.
roots many and long; 1 7. 1.
example of long roots; 1. 8. 5. diseased formation (κραδή); 9. 5. evergreen specimens;
 10. 4. leaves broad;
 1. 10. 7. attachment of leaf-stalk: 3.1.1. propagation; 3. 1 3. produces seed and seedlings; 3. 3. 3. everseed and seedings; 3.5. ever-green in some places; 3 4. 2. time of budding; 3. 6. 1. quick growing; 3. 11 1. leaf of σφέν-δαμνος comp.; 3. 11. 4. has a sort of winter-bud like that of µedia; 4. 5. 6. found at only one place on Adriatic coast: rare in Italy; 4. 5. 7. common in some Mediterranean regions; 4.7.4. size of unnamed Arabian tree (see App. 12a) comp.; 4. 8. 1. grows partially

in water: not common on Nile: 4. 13. 2, trees said to have been planted by Agamemnon; 4.15.2. bark cracks; 4. 16. 2. grows again after being cut or blown down: instances; 5. 8. 4. character of wood; 5. 7. 3. wood used for bent-wood work: soon decays; 5. 9. 4. wood makes an evil smoke when burnt for charcoal; 9, 11, 6, 'head' of στούχvos ò μανικός comp. to fruit of π.

πλατύφυλλος (δρύς) see δρύς (6).

πόα, grass 7. 8. 3. leaves 'on the ground.'

πόα ή Μηδική, see Μηδική. πόθος (1), larkspur, Delphinium orientale

6. 8. 3. a coronary plant: flowers in summer: flower like ὑάκινθος. πόθος (2) (= ἀσφόδελος), asphodel, Asphodelus ramosus

6. 8. 3. a coronary plant: flowers in summer: flower white: used in connexion with funerals.

πόλιον, hulwort, Teucrium Polium

 10. 4. leaves fieshy: prevents moth in clothes; 2. 8. 3. used for caprification; 7 10. 5. evergreen.

πολυάκαυθος, Carduus acanthoides 6. 4. 3. a 'thistle-like' plant. πολυπόδιον, polypody, Polypodium vulgare

9.13.6. peculiar shape: described: named from cuttle-fish (πολύmous), and used as charm to prevent polyp (πολύπους): other medicinal use: habitat; 9.20.4. comes up after rain: has no

πράσιον, Marrubium app.

6. 1. 4. a spineless wild under-shrub; 6. 2. 5. two kinds; see below.

πράσιον (1), horehound, Marrubium peregrinum

6. 2. 5. leaf described: used by druggists.

πράσιον (2), horehound, Marrubium vulgare

6. 2. 5. leaf described.

πράσον (1), leek, Allium Porrum 7. 1. 2-3. time of sowing and of germination; 7. 1. 6. germina-

tion; 7. 1. 7. bears fruit in second year: stem single; second year: sum amage, 7.2.2 S. root makes offsets; 7.3.4. seed borne at top: method of sowing; 7.4.10. coat often like yirttor; 7.4.11. size of 'head' of one year old oxoροδον comp.; 7. 5. 3. improved by transplanting; 7. 5. 4. pests; 7. 5. 5. seed keeps well; 7. 8. 2. stem smooth, not branched;
9. 10. 1. leaf of ἐλλέβορος ο λευκός comp. by some.

πράσον (2) (= ζώστηρ 4.6.2. = φῦκος(1)): grass-wrack. Posidonia

oceanica

4. 6. 2. = ζώστηρ, q.v. πράσου (3) (= φῦκος (2)), riband-weed,

Laminaria saccharina

4. 6. 4. carried by current from Atlantic into Mediterranean: described; 4.7.1. refers to 4 6.4. πρίνος, kermes-oak, Quercus coccifera

1. 6. 1. core hard and close; 1.6.2. core large and conspicuous; 1. 9. 3. evergreen; 1. 10. 6. leaves with spinous projections; 3. 3. 1. a mountain free; 3. 3. 3. 3. 3. 1. a mountain tree; 3. 3. 3. a evergreen; 3. 3. 6. does not al-ways fruit: 3. 4. 1. takes a year to ripen fruit; 3. 4. 4. 6; time of fruiting; 3. 6. 4. deep rooting; 3. 7. 3. produces a scarlet 'berry'; 3. 16. 1. described; 3. 16. 2. σμλαξ (1) comp.; 3. 16. 3. φελλόδρυς δρύς and π. comp.; 3. 16. 4. leaf of κόμαρος comp.; 4. 3. 1. leaf of λωτός (4) comp.; effect of stripping bark in winter; 5. 4. 8. wood of μυρέπ (2) comp. for strength; 5. 5. 4. core not obvious, but 5. 5. 4. core not obvious, but exists; 5. 7. 6. uses of wood: 5. 9. 7. wood used for fire-sticks: 9. 4. 3. leaf of σμύρνα comp. by some.

προύμνη (= σποδίας), bullace, Prunus insititia

9. 1. 2. sap gummy. πτελέα, elm, Ulmus glabra

1.8.5. diseased formation (κραδή): 1. 10. 1. leaves inverted in summer; 1. 10. 6. leaves notched; 3. 1. 1. propagation; 3. 1. 2. seems to have no fruit yet re-

produces itself: instance; 3.1.3. do.: proof; 3. 3. 1. tree of mountain and plain: 3.3.4. a question if it bears fruit; 3.4.2. time of budding; 3. 6. 1. quick growing; 3. 7. 3. produces a βότρυς and leaf-galls; 3. 11. 5. mountain and lowland forms comp.; 3. 14. 1. description: kinds; 3. 15. 4. leaf-galls of τέρμινθος comp.; 3. 17. 3. leaf of Idaean κολοιτία comp.; 3. 17. 5. leaf of συκή ή 'Ιδαία comp.; 3.18.5. leaf of ροῦς comp.; 4. 2. 3. 1eas of συκῆ ἡ Κυπρία comp.; 4. 5. 3. grows in Pontus; 4. 5. 7. common in some Mediterranean regions; 4. 9. 2. leaf of τρίβολος (3) comp.; 4. 16. 2. survives stripping of bark; 5. 1. 2. time of cutting timber; 5. 3. 4. character of wood; 5. 3. 5. haracter of wood; 5. 3. 5. leaf of pous comp.; 4. 2. 3. leaf method of making door-hinges of the wood; 5. 4. 3. wood does not decay if exposed to air; 5. 6. 4. wood strong: used for door-hinges; 5. 7. 3. wood used door-ninges; 5. 7. 3. wood use in shipbuilding; 5. 7. 6. other uses of wood; 5. 7. 8. uses of wood for carpenter's tools; 9. 1. 2. sap gummy; gum contained in the 'bag'; 9. 4. 3. leaf of σμύρνα comp. (by some).

**repic, fern, **Nephrodium Frika' mas*
1. 10. 5. frond described; 4. 2. 11. leaf of an upward Memphis.

leaf of an unnamed Memphian shrub (see App. (2)) comp.; 8.7.7. destroyed by manure or by sheep sleeping on it; 9. 13. 6. leaf of πολυπόδιον comp. to π. η μεγάλη; 9. 18. 8. distinguished from θηλύπτερις; 9. 20. 5. root only of use: medicinal use:

time of gathering. πύξος, box, Buxus sempervirens

1. 5. 4. wood heavy; 1. 5. 5. do. because of close grain; 1. 6. 2. core not conspicuous; 1.8.2. few knots; 1. 9. 3. evergreen; 3. 3. 1. a mountain tree; 3. 3. 3. evergreen; 3.4 6. time of fruiting: fruit inedible; 3.6.1. slow growing (?); 3.15.5. described; 4.4.1. hard to grow in Baby-

lon; 4.5.1. in list of Northern trees; 5. 3. 1. wood very close and heavy; 5. 3. 7. images made from the wood; 5. 4. 1. wood hard and heavy; 5. 4. 2. wood proof against decay; 5. 4. 5. wood not attacked by σκώληξ; 5. 5. 2. core not obvious: wherefore wood not apt to 'draw': 5. 5. 4. core not obvious but exists; 5. 7. 7. uses of wood: that grown on Mount Olympus useless; 5. 7. 8. uses of wood for carpenter's tools; 9. 20. 4. wood of esevos comp.

πυρός, wheat, Triticum vulgare
1.5.2. bark fibrous; 1.6.5. roots
numerous; 1.6.6. do.; 1.11.2.
seeds in a husk; 1 11.5. each seed separately attached; 2.2.9. said to turn sometimes into κριθή; 2.4.1. turns into aipa: wild π . turns into cultivated with cultivation; 4.4.9. πυροί grow in India; 4.10. 8. taste of seeds of India; 4. 10. 8. taste of seeds of σίδη comp.; 7.11.2. inflorescence and general appearance of στελέφουρος comp.; 8. 1. 1. in list of cereals; 8. 1. 8. sown early, but after κριθή; 8. 1. 4. one kind sown late; 8. 1. 5. time of germination; 8. 2. 1. germination described; 8. 2. 3. single leaf first appears: roots described; 8. 2. 6. time of maturing seed; 8. 2. 7. time of harvest in Hellas and in Egypt: 8. 3. 2. stem: and in Egypt; 8. 3. 2. stem; 8. 4. 1-2. compared in detail with other cereals; 8. 4. 3-5. kinds distinguished, local and other (see below); 8. 6. 1. conditions for sowing; 8. 6. 4. suitable soil; 8. 6. 5-6. rain hurtful when π . is in flower: and when it is ripe, but less so than to $\kappa \rho \iota \theta \dot{\eta}$; 8. 7. 1. said to change into $\alpha \iota \rho a$ under certain conditions; 8. 7. 4. effect of cutting down or grazing young crop in Thessaly and in Babylon; 8.7.5. in many places comes up again next year; etc.; 8.8.2. favourable localities; 8.8.3. degenerates into aipa,-or else aipa is specially apt to grow among π .;

8.9.1. exhausts the soil most of cereals; 8. 10. 1. a pest of π .: 8. 10. 2. wheat-rust: 8. 10. 3. effects of weather; 8.10.4. effects of 'worms' in various localities: 8. 11. 1. seed keeps better than κριθή; 8. 11. 3. grain stored without drying; 8. 11. 7. effect of mixing earth with the grain in some places: at Babylon grain jumps on the threshingfloor: reason.

πυρός ὁ Αἰγύπτιος, Triticum vulgare VAI.

 4. 3. in list of varieties of π.; 8. 4. 6. escapes αίρα. πυρὸς ὁ ᾿Αλεξάνδρειος, Triticum vul-

gare var. 8. 4. 3. in list of varieties. πυρὸς ὁ ᾿Ασσύριος, Triticum vulgare var.

8. 4. 3. in list of varieties. πυρός ο Βοιώτιος. Triticum vulgare

8. 4. 5. heaviest grain. πυρός ὁ Θράκιος, Triticum vulgare

8. 4. 3. grain has many coats.

πυρὸς ὁ καχρυδίας8. 4. 3. thick stem. πυρός ὁ κριθανίας 8. 2. 3. branching. πυρός ὁ Λακωνικός 8. 4. 5. grain light.

πυρός ὁ Διβυκός 8. 4. 3. grain not long in husk thick stem. πυρὸς ὁ Ποντικός

8. 4. 3.-4. lightest grain: 8. 4. 5. variation in grain; 8.4.6. escapes αĭρα. πυρός δ σιτανίας

8. 2. 3. branching. πυρός ὁ Σικελός

8. 4. 3., 8. 4. 5. heaviest grain of kinds imported to Hellas; 8.4.6. fairly free from aipa, especially that of Akragas; infested with μελάμπυρον.

πυρὸς ὁ στλεγγύς 8. 4. 3. in list of varieties, see note

ράμνος, buckthorn, Rhamnus spp. 1 5. 3. stem fleshy; 1. 9. 4. evergreen; 3. 18. 12. cluster of ber-

ries of σμίλαξ (2) comp.; 5. 9. 7. wood used for fire-sticks, especially for the stationary piece. ράμνος ή λευκή, buckthorn, Rhamnus graeca

3. 18. 2. distinguished from ρ. ή μέλαινα.

ράμνος ή μέλαινα, buckthorn, Rham-

nus oleoides 3. 18. 2. distinguished from p. n

papavis, radish, Raphanus sativus

1. 2. 7. flesh of root turns to wood; 1. 6. 6. root fleshy; 1. 6. 7. root of 'bark' and flesh; 7. 1. 2-3. time of sowing and of germination; 7.1.5. do.; 7.1.7. germination; 7.2.5. survives and increases in size under a heap of soil; 7. 2. 5–6. root described; 7. 2. 8. do.; 7. 3. 2. seeds described; 7. 3. 4. seed borne at side; 7. 4. 1–2. several kinds (described) Κορινθία, Κλεωναία, λειοθασία, (or Θρακία), άμωρέα (see below), Βοιωτία, and one with leaf like ευζωμον; 7.4.3. effects of weather; 7.5.3. im-proved by transplanting; 7.5.4. pests; 7.6.2. root of wild γογ-γυλίς comp.; 7.6.3. root of ίπποσέλινον comp.; 7.8.2. stem branched; 9.9.1. method of cutting root of μανδραγόρας (1) comp.; 9.12.1. method of cut-ting un γωμαλέρω ή λευνές for Κλεωναία, λειοθασία, (οτ Θρακία), ting up χαμαιλέων ὁ λευκός for medicinal use comp.

ραφανὶς ἡ ἀμωρέα, horse-radish (?) 7. 4. 2. in list of varieties of ρ.

pápavos, cabbage, Brassica cretica 1. 3. 4. becomes tree-like; 1. 6. 6. root single; 1. 9. 4. evergreen; 1. 10. 4. leaves fleshy; 1. 14. 2. bears fruit on top and at side; 4.4.12. size of an unnamed Asian shrub (see App. (10)) comp.; 4.16.6. spoils flavour of grape: vine-shoot turns away from ρ; whence use of \(\delta\). as cure for effects of wine; 6. 1. 2. may be as an under-shrub; 7. 1. 2-3. time of sowing and of germination; 7. 2. 1. propagation; 7. 2. 4. grows again when stem is cut: effect on flavour; 7. 4. 1. several kinds; 7. 4. 4. three kinds distinguished, οὐλόφυλλος, λειόφυλλος, ἀγρία (see below); 7. 5. 3. bears trans-planting; 7.5.4. pests; 7.6.1-2. wild form distinguished,

ραφανος ή άγρία (= κεράϊς 9. 15. 5.), charlock, Raphanus Raphanis-

 4. 4. see ράφανος;
 7. 6. 1-2. see ράφανος;
 9. 15. 5. Arcadian: drug: also called repais.

ράφανος η ορεία (= ἄπιος (2)= ἰσχάς). spurge, Euphorbia Apios

9. 12. 1. used to kill a pig, mixed with χαμαιλέων ο λευκός. ροά (ροιά), (flower κύτινος), pome-

granate, Punica Granatum 1. 3. 3. a tree which has not however a single stem; 1. 5. 1. crooked and low; 1. 6. 1. core fleshy; 1. 6. 3. few roots; 1. 6. 4. shallow rooting; 1. 6. 5. roots branching upwards; 1. 9. 1. much branched; 1. 10. 4. leaves narrow; 1. 10. 10. fruit made of fibre and skin; 1. 11. 4. seeds all together in a single case; 1. 11. 5. each fruit separately attached (?); 1.11.6. arrangement of stones: 1.12.1. taste of fruit; 1.13.1. flower red; 1.13.3. flower above fruit-case; 1.13.4. some kinds sterile; 1.13.5. flower described; 1.14.1. bears on last year's wood; 1.14.4. many cultivated forms; 2. 1. 2-3. propagation; 2. 2. 4 degenerates from seed; 2. 2. 5. degenerates from seed; z. z. 5.
do. details; z. z. 7. in some
places improves from seed;
z. 2. 9. effects of cultivation;
z. 2. 10. ref. to z. 2. 9; z. 2. 11
effect of good cultivation; z. 3.1. sometimes changes character; 2. 3. 2. ref. to 2. 2. 7.; 2. 3. 3. sometimes bears fruit on the stem; 2. 5. 5. propagation; 2. 5. 6. trees should be planted close together; 2. 6. 8. size of fruit of a kind of φοίνιξ (1) comp.; 2. 6. 12. cuttings set upside down; 2.7.1. water-loving; 2.7.3. requires pungent manure and much water; 2.8.1. apt to

shed immature fruit; 3. 5. 4. autumn budding; 3. 6. 2. formation of buds; 3. 18. 4. fruit and growth of κυνόσβατος comp.; 3.18.13. size and leaf of εὐώνυμος comp.; 4. 3. s. stones eaten with fruit; 4. 5. 3. stones eaten with fruit; 4. 5. 3. grows well in Pontus with shelter; 4. 5. 4. grows on Mt. Tmolus and Mysian Olympus; 4. 10. 3. flower of σίδη comp.: seeds of σίδη contrasted; 4. 13. 2. short-lived, especially the stoneless form (see below); 4.13.3. after decaying shoots again from same stock; 4.14.10. fruit gets worm-eaten; 4. 14. 12. unin-jured by special winds; 4. 16. 1. survives splitting of stem; 6.1.3. has spines on the shoots; 7.13.4. fruit kept by inserting stalk in bulb of σκίλλα; 9. 5. 2. size of βάλσαμον comp.

ροά ή απύρηνος, Punica Granatum

var.

4. 13. 2. specially short-lived. ροδωνία (flower ρόδον, fruit μήλον 6. 6. 6.), rose, Rosa centifolia

1. 9. 4. evergreen; 1. 13. 1. colour alluded to; 1.13.2. has a 'twofold 'flower; 1. 13. 3. flower above fruit-case; 1.13.5. flower of ρόα comp.; 2. 2. 1. propagation; 4. 8. 7. colour of flower of κύαμος ὁ Αἰγώπτιος comp.; 4.10.3. sepals of σίδη comp.; 6. 1. 1. in list of under-shrubs; 6. 1.3. has spines on the shoots; 6. 6. 4-6. cultivated under-shrub. a cultivated under-shrub: coronary plant: many kinds: localities: propagation and cultivation; 6.8.2 flowering time; 6.8.5. bush lives five years and then degenerates unless pruned: position and climate important for fragrance: flowers very early in Egypt; 6. 8. 6. blooms well on mountains. but has inferior scent; 9.19.1. colour of flower of δνοθήρας comp.

όδου τὸ ἄγριου, wild rose, Rosa dumetorum

6. 2. 1. flower of κίσθος comp.

poùs (drug poùs 3. 18. 5.), sumach Rhus Coriaria

3. 18. 1. has more than one form (see below); 3. 18. 5. 'male' and 'female' forms: described: used for dyeing: produces a drug called pous.

ρους ή λευκή 3. 18. 2. distinguished from $\dot{\rho}$. $\dot{\eta}$

μέλαινα.

ροῦς ἡ μέλαινα 3. 18. 2. distinguished from ρ. ἡ λευκή

ρύτρος, globe-thistle, Echinops spin-

6. 4. 4. a 'thistle-like' plant: branches from the top.

σάρι (stalk σάρι), Cyperus auricomus 4. 8. 2. useful for food in Egypt; 4. 8. 5. described.

σέλινον, celery, Apium graveolens 1.2.2. takes two years to mature; 1. 6. 6. root single, but with large side-growths; 1.9.4. evergreen (partly); 1. 10. 7. time of leaf-growth; 1. 12. 2. taste of sap; 2. 4. 3. effect of trampling and rolling in seed; 6. 3. 1. leaf of $\sigma(\lambda\phi)$ comp.; 7. 1. 2–3. time of sowing and germination; 7. 1. 6. germination; 7. 1. 7. bears fruit in second year; 7. 2. 2. root makes offsets; 7. 2. 5. root described; 7. 2. 8. do.; 7. 3. 4-5. methods of sowing and transplanting; 7.4.6. kinds distinguished; 7.5.3. kinds distinguished; 7. 5. 3. bears transplanting; 7.6.3. wild forms (ἰπποσέλινον, ἐλειοσέλινον, ορειοσέλινον) distinguished.

σέλινον το έλειον (= ελειοσέλινον), marsh celery, Apium graveo-

8. 1. in list of marsh plants;
 9. 11. 1. leaf of λιβανωτὶς ἡ κάρ-

πιμος comp. σέσελι, hartwort, Tordylium officinale

- 9. 15. 5. an Arcadian drug. σημύδα (?), Judas-tree, Cercis Siliquastrum
 - 3. 14. 4. described; 5. 7. 7. (?) wood used for walking-sticks.

σήσαμον (seed σήσαμη), sesame, Sesamum indicum

1.11.2. seed-vessel; 3.13.6. seeds of berry of ἀκτη comp.; 3.18.13. fruit of εὐώνυμος comp.; 4.8.14. size of fruit of an unnamed Egyptian plant (see App. (20)) comp.; 6. 5. 3. seed of a kind of τρίβολος comp.; 8.1.1. in list of summer crops' distinct from cereals and pulses; 8.1.4. sown later than cereals and pulses; 8 2. 6. time of maturing seed; 8. 3. 1. leaf; 8. 3. 2. stem; 8. 3. 3. flower; 8. 3. 4. seed abundant; 8. 5. 1. a white kind, which is the sweetest; 8. 5. 2. seeds in compartments; 8. 6. 1. rain not beneficial after sowing; 8. 7. 3. not eaten green by any animal: ἐρύσιμον comp.: sown at same time as ὅρμινον; 8.9.3. exhausts the soil; 9.9.2. fruit of ἐλλέβορος comp.; 9.14.4. do.

σίοη, waterilly, Nymphaea alba
4. 10. 1-2. in list of plants of
Lake Copais; 4. 10. 3-4. described; size of fruit of βούτομος comp. (to seed of σ .); 4. 10. 6. grows only in water; 4. 10. 7.

part used for food.

σικύα, bottle-gourd, Lagenaria vul-

garis

1. 11. 4. seeds in a row; 1. 13. 3. flower attached above fruit; 7. 2. 9. root described; 7. 3. 5. takes shape of vessel in which it is grown.

σίκυος (σίκυον), cucumber, Cucumis

sativus

1. 10. 10. fruit made of flesh and fibre; 1. 12. 2. taste of sap; 1. 13. 3. flower attached above 1. 13. 3. nower attached above fruit; 1. 13. 4. some flowers sterile; 2. 7. 5. use of dust; 7. 1. 2-3. time of sowing and germination; 7. 1. 6. germination; 7. 2. 9. root described; 7. 3. 1. long in flower; flower described; 7. 3. 5. effect of soaking seed in milk; 7. 4. 1. several kinds; 7. 4. 6. do. vis. Λακωνικός, σκυταλίας, Βοιώτιος; 7. 5. 2. said to dislike rainwater; 7. 5 8. bears transplanting; 7. 5. 5. seed does not keep well; 7. 5. 6. seed not liable to pests; 7. 13. 1. leaf of άρον comp.

σίκυος ὁ ἄγριος, (drug ἐλατήριον 9. 9. 4.), squirting cucumber, Ecballium Elaterium

4. 5. 1. in list of Northern plants: 7. 6. 4. quite distinct from cultivated σ .; 7. 8. 1. stem 'on the ground'; 9. 9. 4. medicinal use: έλατήριον made from seed 9. 4. 1-2. how long drug will keep: conditions; 9. 15. 6. grows in Arcadia.

σίλφιον (leaf μάσπετον 6. 3. 1), (seed φύλλον, μαγύδαρις 6. 3. 4), silphium, Ferula tingitana

1. 6. 12. root most characteristic o. 12. root most characteristic part; 3. 1. 6. comes up spontaneously; 3. 2. 1. fruits better in wild state; 4. 3. 1. grows in Cyrenaica; 4. 3. 7. consideration postponed; 6. 3. 1-2. described; belongs to 'ferulalike' plants: rules as to cutting and treatment; 6. 3. 3. distri-bution; 6. 3. 4-6. another ac-count, inconsistent in some points; 6.5.2. grows in mountain country; 7. 3. 2. seeds of ἀδράφαξυς comp.; 9. 1. 3. stem and root produce a gum; 9.1.4. gum pungent; 9.1.7. time of tapping: details about juices of stem and root.

σισύμβριον, bergamot-mint, Mentha

aquatica

1.3.1. (?) a typical 'under-shrub'; 2. 1. 3. propagation; 2. 4. 1. turns into μίνθη, unless often transplanted; 6.1.1. in list of under-shrubs; 6.6.2. a culti-vated under-shrub; a coronary plant: the whole plant scented; 6. 6. 3. woody: only one form; 6. 7. 2. said by some to have no fruit: but the wild form certainly seeds; 6. 7. 4. roots described; 6. 7. 6. cultivation; 9. 16. 3. leaf of δίκταμνον (ἔτερον) comp.

σισυρίγχιον, Bai Sisyrinchium Barbary nut, Iris

10. 7. attachment of leaves;

7. 13. 9 (in defective sentence) belongs to τὰ βολβώδη: peculiar growth of root: upper part edible.

σκαλίας, εθε κάκτος (2).

σκαμμωνία, scammony, Convolvulus Scammonia

4. 5. 1. seeks cold regions; 9. 1. 3. root produces a gum; 9. 1. 4. gum has medicinal properties; 9. 9. 1. root and juice used; 9. 20. 5. juice only used.

σκάνδιξ, wild chervil, Scandix Pec-ten-Veneris

7.7.1. a λάχανον; a class of plants called σκανδικώδη; 7.8. 1. stem on the ground.
σκίλλα, squill, *Urginea maritima*1. 6. 7. root in scales; 1. 6. 8.

root fleshy and bark-like: root not tapering; 1. 6. 9. no side roots; 1. 10. 7. no leaf-stalk; attachment of leaves; 2. 5. 5. attachment of leaves; 2. 5. 5. the bulb. of σ ; 7. 2. 2. root makes offsets; 7. 4. 12. formation of roots of $\kappa\rho\delta\mu\nu\nu\rho\nu$ comp.; 7. 9. 4., cf. 1.6.7; 7.12.1. root edible (of the kind called $\dot{\eta} \cdot E\pi\mu\nu\nu\nu\delta\epsilon\nu\rho$) (see below); 7. 13. 1. leaves described; 7. 13. 2. flower-stem pot the only stem; 7. 13. 3. not the only stem; 7. 13. 3. Successive flowering of ἀσφό-δελος comp.; 7. 13. 4. very tenacious of life: hence various uses: use as a charm; 7.13.5-7. stem appears before leaves: sequence described and comp.

with that of νάρκισσος (1); 9, 18, 3, leaf of ὅρχις comp. σκίλλα ἡ Ἐπιμενίδειος, French spar-row-grass, Ornithogalum pyren-

7. 12. 1. see above.

σκόλυμος (=? λειμωνία 6.4.3.), golden

λομος = Γλειμανία 0.4.3.), golden thistle, Scolymus hispanicus 4. 3. a 'thistle-like' plant: leaves spinous; 6. 4. 7. described; 7. 4. 5. leaf of θριδακίνη ή Λακωrun comp.; 7. 10. 1. grows and flowers entirely in summer; 7. 15. 1. flowering depends on the heavenly bodies; 9. 12. 1. leaf of χαμαιλέων ὁ λευκός comp.;

9. 13. 4. an unnamed plant of Tegea comp.

σκολόπενδρον, hart's tongue, Scolopendrium vulgare

9. 18. 7. leaf of ημιόνιον comp. σκόροδον (σκόρδον), garlic, Allium

sativum

 6. 9. no side-roots;
 1. 10. 7.
 attachment of leaves;
 7. 1. 7. stem single; 7. 2. 1. propagation; 7. 2. 3. offsets specially numerous; 7. 4. 1. several kinds; 7. 4. 7. do; 7. 4. 11. do; e.g. τὸ Κύπριον: cultivation etc.; 7. 4. 12. formation of roots of κρόμους contrasted; 7. 8. 2. stem smooth, not branched; 7. 18. 4. grows in colonies because of offsets; 9. 8. 6. eaten as precaution by diggers of έλλέβορος.

σκορπίος (1), Genista acanthoclada 6. 1. 3. has spines for leaves; 6. 4. 1. one of very few plants which are altogether spinous: 6. 4. 2. described.

σκορπίος (2), leopard's bane. Doro-

nicum cordatum

9. 13. 6. peculiar shape, resembles scorpion, and is useful against its sting. σκορπίος (3) (? σκορπίου) (= ἀκουίτου

= θηλύφονον 9.18.2. = μυόφονον),

wolf's bane, Aconitum Anthora
18. 2. properties: habitat:
fatal to scorpion.

σκυθική (= γλυκεία, sc. ρίζα 9.13.2.), liquorice, Glycyrrhiza ylabra 9. 13.2. fragrant: grows on Lake Maiotis: medicinal use: use

against thirst.

σμίλαξ (1), holm-oak, Quercus Ilex

typica
3. 16. 2. described.

σμίλα ξ (?) (2) (= μίλα ξ), smilax, Smilax aspera
3. 18. 11-12. described; 7. 8. 1.

stem clasping.

σμύρνα (gum σμύρνα 9.1.2.), myrrh, Balsamodendron Myrrha

4.4.12. gum of an Arian ἄκανθα (see App. (9)) comp.; 4. 4. 14. in list of Oriental ἀρώματα; 9.1.2. sap gummy (called σμύρνα); 9. 1. 6 time of tapping; 9. 4. 1. collec-

tion of gum; 9. 4. 2. Arabian: habitat; 9. 4. 3. described (two accounts); 9. 4. 4-10. accounts of travellers: 9, 7, 3, in list of

αρώματα.

σόγκος, sow-thistle, Sonchus Nymani 4.6.10. growth of φοῖνιξ (2) comp.; 6. 4. 3. a 'thistle-like' plant, but has not the characteristic 'head' of such plants; 6. 4. 5. stalk of a kind of akavos comp.; 6. 4. 8. root

σπάλαξ (?) (= ἐφήμερον), meadow saffron, Colchicum parnassicum

1. 6. 11. large fleshy roots. σπειραία, privet, Ligustrum vulgare
1.14. 2. bears fruit at top; 6.1.4.

a spineless wild under-shrub. (σπογγία, sponge 4. 6. 5. found on North Coast of

Crete; 4. 6. 10. distinguished from 'plants.')

σποδιάς (= προύμνη), bullace, Prunus insititia

3. 6. 4. very shallow rooting: few roots.

στελέφουρος (= ἀρνόγλωσσον 7.11.2. according to some) (= δρτυξ 7. 11. 2. according to some), plantain, Plantago Lagopus

στοιβή (= $\phi \epsilon \omega_S$ 6. 1. 3.), Poterium spinosum

1. 10. 4. leaves fleshy; 6. 1. 3. has leaves as well as spines: a wild under-shrub; 6. 5. 1. in list of such plants; 6. 5. 2. has

no spines on the leaves. στρουθίον (1) = (κυδώνιος), quince,

Cydonia vulgaris

2. 5. seed produces κυδώνιος. στρουθίον (2) (= στρούθος), 80ap-wort, Saponaria officinalis

6. 4. 3. a 'thistle-like' plant, but has not the characteristic 'head' of such plants; 6.8.3. a coronary plant; flowering time; scentless.

 $\sigma \tau \rho \circ \hat{\upsilon} \theta \circ s = \sigma \tau \rho \circ \upsilon \theta \circ \upsilon (2), so a p-wort,$

Saponaria officinalis

9. 12. 5. leaf of μήκων ή 'Ηρακλεία

comp.

στρύχνος 7. 15. 4. several plants called by this name, which have nothing in common but the name: three mentioned (see below); 9.11.5-6. kinds (see below); 9.15.5. two kinds grow in Arcadia.

στρύχνος ὁ ἐδώδιμος, garden night-

shade, Solanum nigrum 3. 18. 11. fruit of σμίλαξ (2) comp.:

7. 7. 2. a λάχανον: can be eaten raw; 7. 15. 4. more or less cultivated: has berries.

στρύχνος ὁ μανικός (= θρύορος 9.11 6. = περιττός 9.11.6) thorn-apple,

Datura Stramonium

7. 15. 4. causes madness, or, in a large dose, death; 9. 11. 6. described: medicinal use; 9. 19. 1. effect on 'mind.'

στρύχνος ὁ ὑπνώδης, Withania somnifera

7. 15. 4. induces sleep; 9. 11. 5. described: medicinal use: habi-

στύραξ, storax, Storax officinalis 9. 7. 3. in list of ἀρώματα.

συκάμινος, mulberry, Morus nigra 1. 6. 1. core hard and close: 1. 9. 7. time of leafing; 1. 10. 10. fruit made of fibre and skin;

1. 12. 1. taste of fruit; 1. 13. 1. flower 'downy'; 1. 13. 4. attachment of flower; 5. 3. 4. character of wood; 5. 4. 2. do.: wood little liable to decay: turns black when old; 5. 6. 2. wood towel and easy to hand: tough and easy to bend: uses; 5.7.3. wood used for bent-wood work: use in shipbuilding.

συκάμινος ή Αίγυπτία, sycamore

Ficus Sycamorus

1. 1. 7. position of fruit; 1. 13. 2. bears on stem; 4. 1. 5. barren in uncongenial climate; 4.2.1-2. peculiar to Egypt: described; 4. 2. 4. κερωνία distinguished.

συκή (1) (fruit σῦκον), fig, Ficus Carica

1. 3. 1. a typical 'tree'; 1. 3. 5. evergreen at Elephantine; 1.5.1. crooked and low; 1.5.2. bark smooth: bark in one layer; 1. 5. 3. wood fleshy; 1. 5. 3. wood not fibrous; 1. 6. 1. core fleshy; 1. 6. 3. roots many and long; 1. 6. 4. roots crooked; 1. 7 2. roots very long; 1. 8. 1.

no knots; 1.8.2. has less knots than ἐρυνεός; 1.8.5. young branches 'roughest'; 1.9.7. time of shedding leaves; 1.10.4. leaves broad; 1.10.5. leaf divided: etc.; 1.10.8. leaves made of 'bark' and flesh; 1.11. 4. seeds all together in a single seeds; 1. 11. 6. arrangement of seeds; 1. 12. 1. taste of fruit; 1. 12. 2. taste of sap; 1. 14. 1. bears on new shoots: sometimes also on old wood (?): 1. 14. 4. cultivated form of epiveos: many cultivated forms; 2. 1. 2. propagation; 2. 2. 4. degenerates from seed: etc.; 2. 2. 12. cannot be made out of ἐρινεός by cultivation; 2. 3. 1. sometimes changes to έρινεός spontaneously; 2. 3. 3. sometimes bears fruit from behind the leaves: other anomalies; 2. 5. 3. grafting; 2. 5. 4, propagation; 2. 5. 5. cuttings set in a bulb of σκίλλη; 2. 5. 6. easily propagated: trees should be planted far apart; 2. 5. 7. low ground suitable; 2. 6. 6. dates said to vary as much as figs in colour etc.; 2. 6. 12. cuttings set upside down: 2. 7. 1. effects of watering; 2. 7. 5. use of dust; 2. 7. 6. root-pruning etc.; 2. 8 1. apt to shed immature fruit: caprificasned immature runt: caprincation; 2.8.2-4. do. and pseudo-caprification; 3. 3. 8. sheds \$\dop\u00e4\u0 σ. ή Κυπρία comp.; 4. 4. 4. fruit of ή συκή Ἰνδική comp.; 4. 5. 3. grows well in Pontus with shelter; 4.7.7. size of marine trees of island of Tylos comp.; 4.13.1. shorter - lived than convers: shorter - lived than epiveos; 4.13.2. short-lived; 4.14.2. apt to get worm-eaten: young plants liable to 'sunscorch'; 4, 14, 3, destroyed by 'worms' which breed in it: gets scab in some regions; 4. 14. 4-5. other diseases; 4. 14. 8. effects on fruit of rain and drought; 4. 14. 10. infested by knips; 4. 14. 12. suffers most from special winds; 4. 15. 2. survives stripping of bark for some time: stripping of bark for some time; 4. 15. 2. instance of bark grow-4. 15. 2. instance of bark growing again; 4. 16. 1. survives splitting of stem; 5. 3. 3. character of wood; 5. 6. 1. wood strong only against a vertical strain; 5. 9. 5. wood makes pungent smoke; 5. 9. 6. wood good for kindling furnaces; 7. 13. 2. root of ἀσφόδελος eaten with fire. with figs.

συκή (2) ή Αίγυπτία (= κερωνία 1.11.2.), carob, Ceratonia Siliqua συκή (3) ή Ίδαία (fruit σῦκον), Amelanchier vulgaris

17. 4-5. described.
 συκή (4) ἡ Ἰνδική, banyan, Ficus bengalensis

1. 7. 3. roots again from branches: 4. 4. 4-5. described.

συκή (5) ή Κυπρία, sycamore, Ficus Sycamorus var.

4. 2. 3. described.

συκή (6) ή Λακωνική, Ficus Carica

var.
2. 7. 1. water-loving; 2. 8. 1. caprification not used. συκή (7) (ή 'Αραβική), Ficus Carica

var.? 4. 7. 8. an evergreen Arabian kind. (συκή (8) (? an alcyonidian polyp)

4. 6. 2. peculiar to certain waters; 4. 6. 9. described).

σφάκος, sage, Salvia calycina 6. 1. 4. a spineless wild undershrub; 6. 2. 5. like cultivated έλελίσφακος: leaf of one kind of πράσιον comp.

σφένδαμνος, maple, Acer monspessulanum

3. 3. 1. a tree of mountain and plain; 3. 8 doubt whether it has a flower; 3. 4. 4. time of fruiting; 3. 6. 1. slow-grow-ing (?); 3. 6. 5. roots shallow and few according to Arcadians; 3. 11. 1-2. described; 5. 1. 2. time of cutting timber; 5. 1. 4.

do.; 5. 3. 3. character of timber: 5, 7, 6, uses of wood.

σχίνος (fruit σχινίς 9.4.7.), mastich, Pistacia Lentiscus

9, 1, 2, produces a gum; 9, 4, 7. λιβανωτός comp. by some. σχοίνος (1), rush, Juncus spp. etc.

1. 5. 3. not jointed; 1. 8. 1. no knots; 4. 8. 1. in list of τὰ λοχμώδη; 4. 12. 1-3. kinds distinguished.

σχοίνος (2) (ὁ εὐώδης ?), ginger-grass, Cumbopogon Schoenanthus

9. 7. 1. habitat (E. of Lebanon): described: fragrance; 9. 7. 3. in list of ἀρώματα.

σχοίνος (3) ο κάρπιμος, (= μελαγκραvis, 4.12.1.), bog-rush, Schoenus

πίστίcans
4.12.1-3. described.
σχοῦνος (4) ὁ ὁλόσχοινος, Scirpus Holoschoenus

4. 12. 2. described: 9 12. 1. used for stringing pieces of χαμαιλέων ὁ λευκός.

σχοίνος (5) ὁ ὀξύς, Iuncus acutus 4, 12, 1-2, described.

σχοίνος (6) 7. 3. stone 'σχοινοι' in 'Red Sea.')

τέρμινθος (τερέβινθος), terebinth, Pis-

tacia Terebinthus 1. 9. 3. evergreen (wild form); 3, 2, 6, characteristic of Syria; 3. 3. 1. a mountain tree; 3. 3. 3. evergreen; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting: 3. 15. 3-4. described; 4. 4. 7. τ. ή Ἰνδική comp.; 4. 16. 1-2. not injured by cutting for resin; 5.3.2. character of wood: uses: 5. 7. 7. fruit and resin useful, wood not used in craits; 9.1.2 sap gummy; 9. 1. 6. time of tapping; 9. 2. 1. method of tapping; 9. 2. 2. produces best resin (ὀητίνη); 9.2.2. said to be burnt for pitch (πίττα) in Syria: cf. 3. 2. 6.; 9. 3. 4. method of preparing pitch in Syria; 9. 4. 7. σμύρνα comp. by some; 9. 4. 8. some say σμύρνα = τ.; 9. 5. 1. fruit of βάλσαμον comp.

τετραγωνία, Euonymus latifolius
3. 4. 2. time of budding; 3. 4. 6;

time of fruiting. τετράλιξ, yellow star-thistle, Cen-

taurea solstitialis
4. 4. a 'thistle-like' plant: time of growing.

τεῦτλις (= τεῦτλον)
7. 7. 2. a λάχανον; needs cooking. τεύτλον (τεύτλιον) = τεύτλις, beet, Beta maritima

1. 3. 2. becomes large in cultivation; 1.5.3. stem fleshy; 1.6.6. root single, but with large sidegrowths; 1. 6. 7. root fleshy; 1. 9. 2., cf. 1. 3. 2; 1. 10. 4. leaves fleshy; 7. 1. 2-3. time of seaves nesny; 7. 1. 2-3. time of sowing and germination; 7.1.5. do; 7. 1. 6. germination; 7.2.2 root makes offsets; 7. 2. 5-6. root described; 7. 2. 7. root of λάπαθον comp.; 7. 2. 8. root; 7. 3. 2. seeds; 7. 4. 1. several kinds; 7. 4. 4. two kinds, 70 λαμαρίς (Σταλικάν) and πλημά το λευκόν (Σικελικόν) and το μέλαν; 7. 5. 5. seed keeps well.

τήλις (= βουκέρας), fenugreek, Trigonella Foenum-graecum

3. 17. 2. leaf of κολουτέα (κολοιτία comp.

μαλλος (produces ίπποφαές? 9. 15. 6. see note) (= μηκώνιον τιθύμαλλος 9. 8. 2.), spurge, Euphorbia Peplus etc.

9. 8. 2. juice of stalk, how collected; 9. 11. 1. several kinds; 9. 11. 5. do.; leaf of στρύχνος δ υπνώδης comp.; 9. 11. 7-9. kinds (see below); 9. 15. 6. Arcadian: localities.

τιθύμαλλος ὁ ἄρρην, Euphorbia Sibthorpii

9. 11. 8. described: medicinal use.

τιθύμαλλος ο μυρτίτης (fruit κάρυον 9. 11 9.), Euphorbia Myrsinites 9. 11. 9. described: habitat: time of gathering: medicinal use.

τιθύμαλλος ὁ παράλιος, Bea-spurge,

Euphorbia Paralias 9, 11, 7, described: medicinal use.

τίφη, one-seeded wheat, Triticum monococcum

1. 6. 5. roots numerous; 2. 4. 1.

seed, unless bruised, produces $\pi\nu\rho\dot{\phi}_{5}$; 8. 1. 1. in list of cereals; 8. 1. 3. sown early; 8. 2. 1. germination described; 8. 2. 6. time of ripening seed; 8. 4. 1. comp. in detail with other cereals; 8. 8. 3. τ . and $\xi\epsilon\iota\dot{a}$ only plants which can change into something quite different (cf. 2. 4. 1.); 8. 9. 2. exhausts the soil less than any other cereal: reason: likes light soil: τ . and $\xi\epsilon\iota\dot{a}$ the cereals most like $\pi\nu\rho\dot{\phi}_{5}$.

τίφυον, autumn squill, Scilla autum-

nalis
7. 13. 7. flower appears before

leaves and stem. τραγάκανθα (1), tragacanth, Astra-

galus creticus

1. 3. produces a gum: now known to grow not only in Crete (see τ. (2));
 9. 8. 2. no cutting needed to collect gum.
 τραγάκανθα (2), tragacanth, Astra-

galus Parnassi

15. 8. abundant in Achaia and not inferior to the τ. of Crete. τραγοπώγων (= κόμη 7. 7. 1.) goat's beard, Tragopogon porrifolius 7. 7. 1. described; a λάγανον.

7. 7. 1. described: a λάχανον. τρίβολος (1), caltrop, Tribulus terrestris

restris
3. 1. 6 comes up spontaneously in damp places; 6. 1. 3. has leaves as well as spines; has spines on the fruit-vessel;

spines on the fruit-vessel; 6. 5. 3. distinguished from $\tau \rho \epsilon$ - $\beta o \lambda o s$ (2); 7. 8. 1. stem 'on the ground'; 8. 7. 2. (as a weed) destroyed by $\epsilon \rho \epsilon \beta \nu \rho o s$.

τρίβολος (2), caltrop, Fagonia cretica 6. 1. 3. has leaves as well as spines; 6. 4. 1. do.; 6. 5. 1. in list of such plants; 6. 5. 3. distinguished from τρίβολος (1);

grows near enclosures. τρίβολος (3), water chestnut, Trapa

natans

9. 1-3. described.
 τριπόλιον (?), Aster Tripolium
 9. 19. 2. use as charm.

9. 19. 2. use as charm. τριχομανές (? = ἀδίαντον τὸ λευκόν) 7. 14. 1., English maldenhair, Asplenium Trichomanes τύφη, bulrush, Typha angustata 1. 5. 3. not jointed; 1. 8. 1. no knots; 4. 10. 1. in list of plants of Lake Copais; 4. 10. 5. described; 4. 10. 6. grows both on land and in water: some doubt this; 4.10.7. part used for food.

υάκινθος ή ἀγρία, Scilla bifolia

6. 8. 1-2, flowering time. ὑάκινθος ἡ σπαρτή, larkspur, Delphinium Ajacis

6. 8. 2. flowering time: flower of

πόθος (1) comp. ΰδνον, truffle, Tuber cibarium

1. 1. 11. has not all the 'parts' of a plant; 1. 6. 5. no roots; 1. 6. 9. underground.

ύποχοιρίς, cat's ear, Hypochoeris radicala

7: 7. 1. a λάχανον; classed as chicory-like from its leaves; 7. 11. 4. growth contrasted with χόνδρυλλα.

υφεαρ, mistletoe, Viscum album 3. 16. 1. grows on πρίνος.

φακός, lentil, Ervum Lens
2. 4. 2. seed sown in dung; 3. 15. 3.
fruit of τέρμινθος comp.; 3. 17. 2.
slze of fruit of κολοιτία (1) comp.;

size of fruit of κορωνος comp.; 3. 17.2.

size of fruit of κολονία (1) comp.;
3. 18. 5. arrangement of fruit of ροῦς comp.; 4. 4. 9. not found in India; 4. 4. 10. a so-called φ. in India; 8. 1. 4. sown late; 8.3.2. stem; 8. 3. 4. seed; 8. 5. 1. several kinds; white form sweetest; 8. 5. 2. seeds comparatively few; 8. 5. 3. shape of pod; 8. 8. 3. ἄρακος grows specially among φ.; 8. 8. 4. so also ἀπαρίνη; 8. 8. 6. causes etc. of φ. becoming 'cookable' or 'uncookable.'

φάσγανον (= ξ ίφιον = ξ ίφος 7. 13. 1.), corn-flag, Gladiolus segetum

7. 12. 3. use of root in food: root described; 7. 13. 1. leaves described; 7. 13. 4. grown from seed.

φάσκος, tree-moss, Usnea barbata 3. 8. 6. borne only on αἰγίλωψ (1). φελλόδρυς (= ἀρία 3. 16. 3.), holmoak, Quercus Ilex var. agrifolia

1. 9. 3. evergreen; 3. 3. 3. do.;

ΙI

3. 16. 3. described: called apía by Dorians.

φελλός, (? = ίψος cf. Plin. 16. 98.). cork-oak, Quercus Suber

 2. 7. bark;
 5. 2. bark rough and fleshy;
 5. 4. wood light; 3. 17. 1. grows in Tyrrhenia; described; 4. 15. 1. is the stronger for having its bark stripped: 5. 3. 6. wood of point (1) comp.

φέως (= στοιβή 6. 1. 3.), Poterium spinosum.

φηγός (= δρῦς ἡ ἀγρία), Valonia oak,

Quercus Aegilops

3. 3. 1. a mountain tree; 3. 4. 2. time of budding; 3. 6. 1. slow-growing (?); 3. 8. 2. one of the five 'Idaean' kinds of oak: described: fruit; 3.8.3-4. acorns; 3. 8. 4. timber; 3. 8. 7. one of the four 'Macedonian' kinds of oak; 4. 13. 2. ancient trees at Ilium; 5. 1. 2. time of cutting timber.

φιλύκη, alaternus, Rhamnus Ala-

ternus

1. 9. 3. evergreen; 3. 3. 1. a mountain tree; 3. 3. 3. evergreen; 3. 4. 2. time of budding; 3. 4. 4. time of fruiting; 5. 6. 2. easiest wood for turning.

φίλυρα (= φίλυρα ἡ θήλεια), lime (or silver lime), Tilia platyphyllos,

(or tomentosa)

1. 5. 2. bark thin: bark in layers; 5. 2. bark thin: bark in layers; 1. 6. 5. wood pliable; 1. 10. 1. leaves inverted in summer; 1. 12. 4. leaves, but not fruit, eaten by animals; 3. 3. 1. a mountain tree; 3. 4. 2. time of budding; 3. 4. 6. time of fruiting: fruit inedible; 3. 5. 5-6. winter-buds; 3. 10. 4-5. described: 'male' and 'female' forms distinguished (see below). forms distinguished (see below); 3. 11. 1. bark of σφένδαμνος comp.; 3. 13. 1. bark of κέρασος comp.; 3. 13. 3. grows where κέρασος grows; 3. 17. 5. leaf of συκή ή Ίδαία comp.; 4.4.1. hard to grow in Babylon: 4. 5. 1. in list of Northern trees: 4. 8. 1. grows partially in water; 4.15.1. outer bark can be stripped;

4. 15. 2. survives stripping of bark for some time; 5. 1. 2. time of cutting timber: 5. 1. 4. do.; 5. 3. 3. character of wood; 5. 5. 1. wood easy to work; blunts tools; 5. 6. 2. wood soft and easy to work; 5.7.5. various uses of wood and bark; 5. 9. 7. wood used for fire-sticks.

φίλυρα ἡ ἄρρην (= φιλυρέα), mock-privet, Phillyrea media 3, 10, 4-5, distinguished from φ, ἡ

θήλεια.

φίλυρα ἡ θήλεια (= φίλυρα), silver-lime, Tilia tomentosa 3. 10. 4-5. distinguished from φ. ἡ

ἄρρην. φιλυρέα, mock-privet. Phillyrea

media 1. 9. 3. evergreen; 3. 4. 6. (?)

time of fruiting.

φλεώ (φλεώς), Erianthus Ravennae 4.8.1. in list of τὰ λοχμώδη; 4.10.1. in list of plants of Lake Copais; 4. 10. 4. described; 4. 10. 6. grows both on land and in water; grows on the floating islands of Lake Copais; 4. 10. 7. part used for food; 4. 11. 12. foliage of some κάλαμοι comp.

φλόγινον (= φλόξ), Wall-flower, Cheir-anthus Cheiri

8. 1-2, a coronary plant: flowering time.

φλόμος ή μέλαινα, mullein, Verbascum sinuatum

9. 12. 3. leaf of μήκων ή κερατίτις comp.

φλόξ (= φλόγινον), wall-flower.

Cheiranthus Cheiri 6. 6. 2. a cultivated under-shrub; a coronary plant: scentless; 6. 6. 11. grown from seed.

φοίνιξ (1), date-palm, Phoenix dacty-

lifera

1. 2. 7. 'flesh' turns to wood; 1. 4. 3. (?) tolerant of sea-water 1. 5. 1. few branches; 1. 5. 2. rough bark; 1. 5. 3 wood fibrous; 1. 6. 2. core not distinguishable; 1. 9. 1. growth chiefly upwards; 1. 9. 3. evergreen; 1. 10. 5. reedy leaves; 1.11.1. seed immediately within envelope: envelope not single;

1. 11. 3. seed solid and 'dry' 1. 11. 8. seed solid and 'dry' throughout; 1. 13. 5. 'male' only flowers; 1.14.2. bears fruit at top; 2. 2. 2. propagation; 2. 2. 6 seeds come true; 2. 2. 8. effect of locality; 2.2.10. effects of climate; 2. 6. 1. propagation from fruit; 2. 6. 2. propagation from 'head'; 2. 6. 3. cultivation; 2. 6. 4. do; 2. 6. 5. cultivation in Syria; 2. 6. 8. various kinds; (2. 6. 9. branching kind; see κουκόφορον]: [2. 6. 10. kinds; [2. 6. 9. branching kind: see κουκιόφορον]; [2. 6. 10. shrubby kind: see κοίξ]; 2.6.12. cuttings set upside down; 2. 8. 1. apt to shed immature fruit; 2. 8. 4. artificial fertilisation; 3. 3. 5. not fruitful wherever it grows; 3. 13. 7. dwarf form (? φ. ο χαμαιρριφής); 4.1.5. cf. 2. 2. 10; 4. 2. 7. κουκιόφορον comp.; 4. 3. 1. grows in parts of Libya; 4. 3. 5. grows well in waterless Libya: salt in soil, water supply; 4. 3. 7. kept alive by dew in dry regions; 4. 4. 3. sown in pots; 4.4.13. dangerous sown in pots; 4.4.13. dangerous sown in pots; 4.4.13. dangerous to eat unripe dates (in Gedrosia); 4. 7. 8. occurs on the island of Tylos; 4. 13. 2. story of the very old ϕ . on Delos; 4. 14. 8. if topped, becomes barren, but is not destroyed; 4. 15. 2. survives stripping of bark; 5. 3. 6. character of wood: used for images: 5. 6. 1. wood strong: behaviour under pressure: 5. 9. 4. wood makes a very evil smoke when burnt for charcoal; 6. 4. 11. seed-vessel of κάκτος (1), when stripped of seeds, comp. to 'brain' of φ.; 9. 4. 4. mats in Arabia made of leaves of ϕ .

φοινιξ (2), Nannorhops ritchiana

4. 4. 8. Bactrian.

φοινιξ (3), ο χαμαιρριφής, dwarf palm, Chamaerops humilis

2. 6. 11. described; 3. 13. 7. (?).

φοῖνιξ (4), Callophyllis laciniata
4. 6. 2. peculiar to certain waters;
4. 6. 10. described.

φόνος (= ατρακτυλίς 6. 4. 6.), distaffthistle, Carthamus lanatus

6. 4. 6. reason for name

φῦκος (1) τὸ πλατύφυλλον (= ζώστηρ 4. 6. 2. = πράσον (2)), grasswrack, Posidonia oceanica

4. 6. 2. occurs generally in Greek waters: root described.

φύκος (2) θαυμαστόν το μέγεθος, Γίband-weed, $(=\pi\rho\alpha\sigma\sigma\nu$ (3)), Laminaria saccharina

4. 6. 4. described: grows in Atlan-tic: washed into Mediterranean; 4. 7. 1. refers to 4. 6. 4.

φῦκος (3) τὸ πόντιον

4.6.4. collected by sponge-fishers. φῦκος (4) τὸ τριχόφυλλον, Cystoseira foeniculosa

4. 6. 3. described.

φῦκος (5), litmus, Roccella tinctoria 4, 6, 5, Cretan; dye described.

φῦκος (6), grass-wrack, Cymodocea nodosa (and Zostera marina) 4. 6. 6. described: comp. to

άνρωστις.

χαλβάνη, 808 πάνακες (τὸ Σύριον). χάλκειος, Carlina corymbosa 6. 4. 3. a 'thistle-like' plant.

χαμαίβατος (= βάτος), Rubus ulmifolius

3. 18. 4. described.

χαμαιδάφνη, periwinkle, Vinca her-

3. 18. 13. leaf of εὐώνυμος comp. χαμαίδρυς, germander, Teucrium Chamaedrys

9. 9. 5. medicinal use: described.

χαμαιλέων, chamaeleon 6. 4. 3. a 'thistle-like' plant, but leaves not spinous (see n. on 6. 4. 8.); 6.4.8. flower turns into 'down'; 9. 12. 1-2, kinds (see below).

χαμαιλέων ο λευκός (= ἄκανθα (9) 9. 12. 1. = ἄκανος = ἰξία (2) = ifin), pine-thistle, Atractylis

gummifera

9. 12. 1. described: medicinal use: fatal to dogs and pigs: how administered: grows every where,

χαμαιλέων ὁ μέλας, Cardopatium

corymbosum

9. 12. 2. described: medicina use: habitat: fatal to dogs medicinal 9. 14. 1. how long drug will

χελιδόνιον, greater celandine, Chelidonium maius

7. 15. 1. flowering depends on the heavenly bodies.

χόνδρυλλα, Chondrilla juncea

7. 7. 1. a λάχανον: classed as 'chicory-like' from its leaves;
 7. 11. 4. growth of ὑποχοιρίς contrasted.

ψευδοδίκταμνον, Ballota acetabulosa

 16.2. comp with δίκταμνον as to appearance and properties: said by some to be only a degenerate form of δίκταμνον: method of keeping.

ωκιμον, basil, Ocymum basilicum
1. 6. 6-7. root woody; 1. 10. 7.

time of leaf-growth; 7. 1. 2-3. time of sowing and germination; 7.2.1. propagation; 7.2.4. grows again when stem is cut; 7. 2. 7-8. root described; 7.3.1. long in flower; 7. 3. 2-3 seeds described; 7. 3. 4. seed very abundant: seed borne at top; 7. 4. 1. only one kind; 7. 5. 2. watered at mid-day; 7. 5. 4. effect of hot weather; 7. 5. 5. seed does not keep well; 7. 7. 2. leaf of κόρχορος comp.; 7. 9. 2. flowers borne in succession, 7. 7. 3. 1.; 9. 18. 5. leaf of ἀρρεο-σνον and of θηλύγονον comp. Σχος, Lathyrus Ochrus

8. 1. 3. sown early; 8. 3. 1. leaf; 8. 3. 2. stem; 8. 10. 5. infested by worms.

APPENDIX OF UNNAMED PLANTS

The following plants (arranged in the order of mention) which are described or indicated, but not named, in the Enquiry, seem possible to identify:—

- (ὅμοιον τῷ ἀράκῳ), tine-tare, Lathyrus tuberosus
 6. 12. root described.
- 2. (ὑλήμα ἴδιον τι περὶ Μέμφιν), Mimosa asperata
- 4. 2. 11. described: 'sensitive.'
 3. (δένδρον . . . μεγαλόκαρπον), Jackfruit, Artocarpus integrifolia

4. 4. 5. used for food by Indian sages.

- 4. (φύλλον . . . τοῖς τῶν στρουθῶν πτεροῖς ὅμοιον), banana, Musa sapientum 4. 4. 5. described.
- (καρπὸς σκολιὸς ἐσθιόμενος δὲ γλυκύς), mango, Mangifera indica
 - 4. 4. 5. fruit described: causes dysentery.
- 6, (καρπός όμοιος τοις κρανέοις), jujube, Zizyphus Jujuba 4, 4, 5.

- (ὅμοιον τερμίνθφ), pistachio-nut, Pistacia vera
 4. 4. 7. described.
- 8. (ὅμοιον τῆ ὅψει καὶ τὸ βούκερας),
 Phaseolus Mungo
 - 4. 9-10. called by Hellenes φακός, and similarly used.
- 9. (ἄκανθα ἐφ' ἦς γίνεται δάκρυον) (= ἄκανθα (4) ἡ Ἰνδική), Balsamodendron Mukul
 - 4. 4. 12. grows in Aria: de scribed.
- (ΰλημα ἥλικον ῥάφανος), Asafoetida, Scorodosma foetidum
 4. 12. described: fatal to
- horses. 11. (ὅμοιον τῆ δάφνη φύλλον ἔχον), Nerium odorum
- 4. 4. 13. effect on animals.
- 12a. (δένδρα μέγαλα), mangrove, Bruguiera gymnorhiza 4.7.4. described.

- 12b. (δένδρον φύλλον ἔχον ὅμοιον τῆ δάφνη) mangrove, Rhizophora mucronata
 - 4. 7. 4. described (as if identical with 12a.).
- 13. (καρπὸς ὅμοιος τοῖς θέρμοις), Aegiceras maius
- 4.7. 5, 6, and 7. described.
 14. (δένδρα ὅμοια τῆ ἀνδράχλη (= δάφνη (6) = ἐλάα (3)), white mangrove, Avicennia officinalis
 4.7. 5. described.
- (δένδρα τὸ ἄνθος ἔχοντα ὅμοιον τῷ λευκοίῳ) (= (16)), tamarind, Tamarindus indica
 - 4. 7. 8. grows in the island of Tylos.
- 16. (δένδρον πολύφνλλον) (= (15)), tamarind, Tamarindus indica
 - 4. 7. 8. grows in the island of Tylos: opening and closing of flower.
- 17. (συκή οὐ φυλλοροοῦσα), Ficus laccifera
- 4. 7. 8. grows in the island of Tylos.
- 18. (ὅμοιον τοῖς κρίνοις) Ottelia alismoides
 - 4. 8. 6. Egyptian marsh-plant: habitat and leaves: medicinal use.

- 19. (εν τι γένος εν ταις λίμναις), Saccharum biflorum
- 4. 8. 13. use for fodder. 20. (γένος παραφυόμενον εν τῷ σίτῳ),
- Corchorus trilocularis
 4. 8. 14. treatment as fodder:
- fruit described. 21. (δένδρον), Sissoo-wood, Dalbergia
- Sissoo 5. 3. 2. wood described: use for
- making furniture. 22. (ξύλον), teak, Tectona grandis
- 5. 4. 7. wood stands sea-water well.
- (δένδρον), calamander wood,
 Diospyros quaesita
 4. 7. wood described.
- (ὅμοιον ἀβροτόνψ), Artemisia camphorata
 6.3.6. properties: effect on sheep.
- τη νάρδφ προσεμφερή την δσμην έχον), Valeriana Dioscoridis
- 9. 7. 4. a Thracian ἄρωμα. 26. (ὑποφυόμενον εὐθύς ἐκ τῆς ῥίζης τῷ κυμένω), broom-rape, Orobanche versicolor
- 8. 8. 5. parasitic on κύμινον.
 (ρίζα θανατηφόρος), Somali arrowpoison, Acokanthera Schimperi
 - 9. 15. 2. Aethiopian: used for poisoning arrows.

KEY TO THE INDEX

I.—LIST OF PLANTS MENTIONED IN THE ENQUIRY UNDER BOTANICAL NAMES

Abies cephalonica
— pectinata
Acacia albida
— arabica
— tortilis
Acantha arabica
Acer campestre
— creticum

Aconitum Anthora

Acorus Calamus

Adiantum Capillus-Veneris Aegiceras majus Aegilops ovata Ailanthus malabarica Ajuga Ira Allium Cepa and vars.

— nigrum
— Porrum
— sativum
Alnus glutinosa
Althaea officinalis

Amaranthus Blitum Amelanchier vulgaris Amomum subulatum

έλάτη (1) **ἐλάτη (2)** ἄκανθα(1),etc. ἄκανθα(1),etc. ăκανθα (3) ἄκανθα (7) ζυγία γλείνος σφένδαμνος κλινότροχος App. (27) ἀκόνιτον, θηλύφονον, μυόφονον, σκορ-πίος (3) κάλαμος ὁ εὐώδης ἀδίαντον, ἀ. τὸ μέλαν App. (13) αἰγίλωψ (2) κώμακον ἐπετίνη γήθυου, γήτειον, κρόμυον. κ. τὸ σχιστόν, κρομυογήτειον μῶλυ πράσον (1) σκόροδον κλήθρα άλθαία, μαλάχη η άγρία

βλίτον

ἄμωμον

συκή ή Ίδαία

Anagallis caerulea Anchusa tinctoria Andropogon Ischaemum Anemone blanda

---- coronaria ---- pavonina

—— spp. Anethum graveolens

Anthemis chia

Antirrhinum Orontium Apium graveolens

Arbutus Andrachne
— hybrida
— Unedo
Aristolochia rotunda
Artemisia Absinthium
— arborescens
— camphorata
Artocarpus integrifolia
Arum italicum

Asparagus acutifolius Asphodelus ramosus

Arundo Donax

Asplenium Ceterach
— Trichomanes

κόρχορος ἄγχουσα ἴσχαιμος

ἀνεμώνη ἡ ὀρεία ἀ. ἡ λειμωνία

ά. ἡ λειμωνία ἀ. ἡ λειμωνία ἀνεμώνη ἄνηθον, ἄννη-

ἄνθεμον, ἀ. τὸ φυλλῶδες ἀντίρρινον

έλειοσέλινον, σέλινον, σ. τὸ ἔλειον ἀνδράχλη ἀφάρκη κόμαρος ἀριστολοχία ἀψίνθιον ἀβρότονον App. (24) App. (3)

άρον δόναξ, κάλαμος ὁ αὐλητικός etc. ἀσφάραγος ἀσφόδελος, πόθος (2) ἡμιόνιον ἀδίαντον τὸ

λευκόν, τρι-

χομανές 487

Aster Amellus

— Tripolium

Astragalus creticus

— Parnassi

Atractylis gummifera

Atriplex Halimus
— rosea
Atropa Belladonna

Avena sativa Avicennia officinalis

Balanites aegyptiaca Ballota acetabulosa

— pseudo-Dictamnus Balsamodendron Mu-

kul
— Myrrha
— Opobalsamum
Bambusa arundinacea

Beta maritima
Brassica alba
—— cretica
—— Rapa
Bruguiera gymnorhiza
Bryonia cretica

Buxus sempervirens

Calamintha incana
Calamogrostis Epigeios
Callitriche verna
Calitris quadrivalvis
Callophyllis laciniata
Calycotome villosa
Carpparis spinosa
Cardopatium corymbosum
Carduus arvensis
— acanthoides
Carex riparia
Carlina corymbosa
Carthamus lanatus

άστέρισκος τριπόλιον τραγάκανθα (1) τραγάκανθα (2) ά καν θα (8), άκανος, ίξία (2), ίξίνη, χαμαιλεών ὁ

(2), ίξίνη, χαμαιλεών ὁ λευκός
ἀλιμον ἀδράφαξυς
μανδραγόρας (2)
βρόμος
δ ά φνη (6), ἐλ ά α (3),
App. (14)

βάλανος ψευδοδίκταμνον δίκταμνον (έτερον)

ἄκανθα (4),
App. (9)
σμύρνα
βάλσαμον
κάλαμος ὁ Ἰνδικός
τεῦτλον
νάστι

ναπυ
ράφανος
γογγυλίς
Αpp. (12a)
αμπελος ἡ ἀγρία,μήλωθρον
πύξος

έλένιον κάλαμος (ἐπί-YELOS) λέμνα θύον φοίνιξ (4) άσπάλαθος κάππαρις χαμαιλεών μέλας άκανθα (2) πολυάκανθος Βούτομος χάλκειος άτρακτυλίς. φόνος

Carthamus leucocaulos κυῆκος ἡ ἀγρία

Castanea vesca

Celtis australis Centaurea calcitrapa — salonitana — solstitialis Ceratonia Siliqua Cerris Siliquastrum

VAT

Chamaerops humilis
Cheiranthus Cheiri
Chelidonium majus
Chondrilla juncea
Cicer arietinum
Cichorium Intybus
Cinnamomum Cassia
— iners
Cistus salvifolius
— villosus

Citrus medica

Clematis vitalba Cnicus Acarna — benedictus

Colchicum parnassicum Colocasia antiquorum Colutea arborescens Conium maculatum Convolvulus Scammonia

— sepium Corchorus trilocularis Cordia Myxa

Coriandrum sativum
Cornus Mas
— sanguinea
Corydalis densifiora
Cotoneaster Pyracantha
Corylus avellana
— var.

Crataegus Heldreichii —— orientalis κυῆκος ἡ ἀγρία κυῆκος, κ. ἡ ἡμερος, κρόκος ὁ ἀκανθώδης διοςβάλανος καρύα ἡ Εὐβοϊκή λωτός (1) παντάδουσα

κενταύριον τετράλιξ κερωνία κερκίς (1) σημύδα φοίνιξ (3) φλόγινον, φλόξ χελιδόνιον χόνδρυλλα ερέβινθος κιχόριον κινάμωμον κασία κίσθος ὁ θήλυς κίσθος, K. Ó aponv μηλέα ἡ Περσι-κή (Μηδική) άθραγένη **ἄκο**ρνα κνήκος ή άγρία (ἐτέρα) έφήμερον, σπάλαξ

ἰασιώνη
Αρρ. (20)
κοκκυμηλέα (ἡ
Αἰγυπτία)
κορίαννον
κράνεια
θηλυκράνεια
θήσειον
δένάκανθος

ούϊγγον

κολυτέα

κώνειον

σκαμμωνία

καρύα καρύα ή Ήρακλεωτική κράταιγος μεσπίλη ή ἀνθηδών

Crataegus oxyacantha	μεσπίλη ή ἀν- θηδονοείδης	Elettaria Cardamo-	καρδάμωμον
Crepis Columnae	ἀπαργία	Ephedra campylo-	θραύπαλος
Crocus cancellatus	κ, δ λευκός	poda	opaonato;
an dissura	κ. ὁ εὔοσμος	Erianthus Ravennae	φλεώ
spp.	κρόκος	Erica arborea	ἐρείκη
Cucumis sauvus	σίκυος	Eruca sativa	<i>ε</i> ύζωμον
Cucurbita maxima	κολοκύντη	Ervum Lens	φακός
Cuminum Cyminum	κύμινον	Eryngium campestre	ήρύγγιον
Cupressus semper- virens	κυπάριττος	Erythraea Centau- rium	κενταυρίς
Cuscuta europaea	ὀροβάγχη	Euonymus europaeus	εὐώνυμος
Cyclamen graecum	κυκλάμινος	latifolius	τετραγωνία
Cydonia vulgaris	κυδώνιον,	Euphorbia antiquo-	ἄκανθα (5), (6)
Cydolla vulgaris	στρουθίον(1)	rum	unusuu (0), (0)
Comphanagen Schoon			#= (9) !=
Cymbopogon Schoen-	σχοίνος (2)	Apios	απιος (2), iσ-
anthus			χάς, ράφανος
Cymodocea nodosa	φῦκος (6)	l ·	η όρεία
Cynara Cardunculus	κάκτος (1)	Myrsinites	τιθύμαλλος δ
Scolymus	κάκτος (2)	i	μυρτίτης
Cynodon Dactylon .	άγρωστις	paralias	τιθύμαλλος ο
Cyperus auricomus	σάρι	•	παράλιος
esculentus	μαλιναθάλλη,	Peplus	μηκώνιον, τιθύ-
Cooulonda	μνάσιον	Lopius	μαλλος
loneus		Sibthorpii	τιθύμαλλος δ
longus	κύπειρος	Sibiliorph	
Papyrus rotundus	πάπυρος		ãρρην ,
rotundus	κύπειρον	spp.	μηκώνιον, ὶπ-
Cystoseira Abies ma-	έλάτη (3)		ποφέως, τι-
rina			θύμαλλος
ericoides	δρῦς (7)		
foeniculosa	φῦκος (4)	Fagonia cretica	τρίβολος (2)
Cytisus aeolicus	κολοιτία (1)		δεύη
		Fagus silvatica	
Dalbergia Sissoo	App. (21)	Ferula communis	ναρθηκία, νάρ-
Daphne Gnidium	κνέωρον		$\theta\eta\xi$
- oleoides	κνέωρος ὁ λευ-	nodosa	πάνακες τὸ
Olcoldon	KÓS O KEU-		'Ασκληπίειον
Datura Stramonium		tingitana	σίλφιον
Davura Stramonium	θρύορον, περιτ-	Ferulago galbanifera	πάνακες
	τός, στρύχ-	Ficus bengalensis	συκή ή Ίνδική
D G 1	νος ὁ μανικός	Carica	συκή (1)
Daucus Carota	δαῦκον (1)	- var.	συκή ή Άρα-
Delphinium Ajacis	ὑάκινθος ἡ	- va.	βική η Αρα- βική
	σπαρτή	landfore	
orientale	πόθος (1)	laccifera	App. (17)
- Staphisagria	άσταφίς	Sycamorus	συκάμινος ή
Dendrocalamus stric-	κάλαμος ὁ Ίν-		Αίγυπτία
tus	δικός	var.	συκή ή Κυπρία
		Fraxinus excelsior	βουμέλιος
Diospyros Ebenum	έβένη (1)	Ornus	μελία
— Melanoxylon	ἐβένη (2)	Fucus spiralis	άμπελος (3)
Dracunculus vulgaris	δρακόντιον	Fungi	μύκης
Drypis spinosa	δρυπίς	- uBr	1-113
Ecballium Elaterium	σίκυος ὁ ἄγριος	Galanthus nivalis	λευκόϊον (2)
Echinops spinosus	ρύτρος	Galium Aparine	άπαρίνη
Echium diffusum	ὀνοχειλές	Genista acanthoclada	σκορπίος (1)
			489

Gladiolus segetum

Glaucium flavum var Serpierii Glycyrrhiza glabra

Gossypium arboreum

Hedera Helix Helichrysum siculum Heliotropium villosum Helleborus cvclophyllus Herniaria glabra Hippuris vulgaris Hordeum sativum and vars. Hyphaene thebaica

Ilex Aquifolium Imperata arundinacea Inula Helenium

graveolens

viscosa spp.

Iris foetidissima - pallida etc. - Sisyrinchium

Juglans regia

Juneus acutus - spp. Juniperus communis

excelsa - foetidissima - phoenicea

Oxycedrus

Laburnum vulgare Lactuca graeca sativa

- scariola 490

κύτισος (1) λιβανωτίς θρίδαξ θριδακίνη

ξίφιον, ξίφος, φάσγανον μήκων ἡ κερατίτις γλυκεῖα (ῥίζα). Σκυθική (δένδρον Tò) ἐριὀφορον

έλιξ, κιττός έλειόχρυσος ήλιοτρόπιον

έλλέβορος, έ. ὁ μέλας έλλεβορίνη ϊπνον κριθή

κόϊξ, κουκιόφοοον

κήλαστρος θρύον

πάνακες τò Χειρώνειον κόνυζα ή θήλεια κόνυζα ή ἄρρην κόνυζα

ξίρις lous σισυρίγχιον

καρύα ή Περσική σχοίνος ὁ ὀξύς σχοίνος κεδρίς κέδρος (2) θυία ἄρκευθος, ĸéδρος (3) κέδρος (1), ὀξύκεδρος

Lagenaria vulgaris Laminaria saccharina Lapidium sativum Lathyrus amphicar-

pus - Ochrus - sativus - tuberosus Laurus nobilis Lavandula spica Lavatera arborea Lecokia cretica

Lemna minor Ligustrum vulgare Lilium candidum etc.

— chalcedonicum

Martagon Limnanthemum nymphoides Linum usitatissimum Lolium temulentum Lonicera etrusca Loranthus europaeus Lupinus alba

Lychnis coronaria Lycoperdon Bovista giganteum

Malabaila aurea Malva silvestris Mandragora officinarum

Mangifera indica Marrubium peregrinum

Matthiola incana

Matricaria Chamomilla Medicago arborea - sativa

Melissa officinalis Mentha aquatica – Pulegium - viridis

Mercurialis perennis

σικύα φυκος (2) κάρδαμον ἀράχιδνα

ὧχρος λαθύρος App. (1) δάφνη (1) ζφυον μαλάχη (1) λιβανωτός 'n κάρπιμος ϊκμή σπειραία

κρίνον, κρινωνία, λείριον(1) κρίνον τὸ πορφυρούν ημεροκαλλές μήνανθος

λίνον αἴρα κλύμενον i£ía (1) θέρμος λυχνίς πέζις ἄσχιον

δαῦκον (2) μαλάχη (2) μανδραγόρας (1) App. (5) πράσιον

ίον τὸ λευκόν. ἰωνία (ἡ λευ÷ κή), λευκόϊον (1)

άνθεμον τò άφύλλανθες κύτισος (2) (πόα) ή Μηδική μελισσόφυλ-

λον σισύμβριον βληγώ ηδύοσμον, μίνθη

ἀρρενόγονον, θηλύγονον

Mespilus germanica μεσπίλη, μεσ-Orobanche cruenta ແມ່ນດ໌ດີພວດນ πίλη η σατά- versicolor App. (26) etc. DELOC. Orvza sativa ὄρυζον App. (2) Ostrya carpinifolia οστρύα, δστρυς Mimosa asperata Mimusops Schimperi περσέα. Ottelia alismoides App. (18) πέρσιον Paeonia officinalis Musa saplentum App. (4) γλυκυσίδη, Muscari comosum etc. παιωνία Βολβός Myrtus communis μυρρίνη Palinrus australis παλίουρος βολβὸς ὁ ἐριό-Pancratium maritimum φορος Nannorhops ritchiana Panicum miliaceum φοινιξ (2) κέγχρος Narcissus poeticus νάρκισσος (2) Papaver hybridum μήκων ή ροιάς – serotinus λείριον (2). - Rhoess μήκων ή μένάρκισσος(1) λαινα - somniferum --- Tazetta λείριον (2) μήκων (ἡ ὀπώλείριον (2) - spp. δης), νηπεν-Nardostachys Jataνάρδον θές μήκων ἀλσίνη mansi - spp. Parietaria cretica Nelumbium specioκύαμος ὁ Αίγύπτιος . Petroselinum sativum ορειοσέλινον gum Nephrodium Filix. TTEDÍS Peucedanum officiπευκέδανον nale Nerium Oleander Phillyrea media δάφνη ή άγρία, φιλυρέα δνοθήρας Phoenix dactylifera doivit (1) Phragmites commuκάλαμος ο χαodorum App. (11) Neslia paniculata μελάμπυρον nia ρακίας Pinus brutia Nuphar luteum μαδωνάϊς, νυμπίτυς ή φθειρόφαία ποιος Nymphaea alba σίδη --- halepensis πίτυς; see also - stellata λωτός (2) under πίτυς η άγρία --- Laricio πεύκη ή άκαρ-Ocymum basilicum ὥκιμον πος, Olea cuspidata έλάα (2) θήλεια, π. ή έλάα (1) Ίδαία - europaea --- Oleaster άγριέλαιος, κόπεύκη ἡ ἤμε-ρος, π. ἡ κω---- pinea τινος νόφορος Ononis antiquorum ονωνίς Onopordon illyricum δνόπυξος πεύκη Opoponax hispidus πανάκεια, πάν-Pimpinella Anisum άννησον akes to 'Ho-Piper nigrum πέπερι άκλειον Pistacia Lentiscus σχίνος Orchis longicruris τέρμινθος ὄρχις Terebinthus - papilionacea ὄρχις vera App. (7) Origanum Dictamnus Pisum sativum δίκταμνον πισός - heracleoticum δρίγανος Plantago Coronopus κορωνόπους λευκή - crassifolia θρυαλλίς - Lagopus — Majorana ἀμάρακον ορτυέ, στελέ-- viride etc. δρίγανος, δρίφουρος κύνωψ γανος ή μέ-λαινα ---- lanceolata - major ἀρνόγλωσσον

σκίλλα ή Έπι-

μενίδειος

βολβίνη

Ornithogalum pyre-

- umbellatum

naicum

πλάτανος

περδίκιον

Platanus orientalis

mariti-

Polygonum

Polygonum Persicaria		Ranunculus Ficaria	ἀφία
Polypodium vulgare	πολυπόδιον	Raphanus Raphani-	κεράϊς, ράφ
Polypogon mouspeli-	ἀλωπέκουρος	strum	νος ή άγρίο
ensis Polyporus igniarius	πίλος	sativus	ραφανίς
Populus nigra	αϊγειρος	Rhamnus alaternus —— graeca	φιλύκη
tremula	κερκις (2)	oleoides	ράμνος ή λευ
Portulaca oleracea	ἀνδράχνη	Oleoldes	ράμνος ἡ μ λαινα
Potentilla reptans	πενταπετές,	spp.	ράμνος
-	πεντάφυλλον	Rhizophora mucro-	App. (12b)
Poterium spinosum	στοιβή, φεώς	nata	rippi (rao)
Prangos ferulacea	ὶππομάραθον,	Rhus Coriaria	စ်ဝပိဋ
	μαγύδαρις	Cotinus	κοκκυγέα
Prunus Amygdalus	άμυγδαλή	Ricinus communis	κρότων
avium	κέρασος, λα-	Rosa canina	κυνόρροδον
domestica	κάρη	centifolia var.	ροδωνία
insititia	κοκκυμηλέα	dumetorum	ρόδον τὸ ἄγριο
msiviva	προύμνη, σπο- διάς	— sempervirens Rubus ulmifolius	κυνόσβατος
Mahaleb	πάδος (πηδός ?)	Trubus ulmitolius	βάτος, χαμαι βατος
Pteris aquilina	θηλύπτερις	Roccella tinctoria	φῦκος (5)
Puccinia graminis	έρυσίβη	Rumex conglomera-	λάπαθον τὸ ἄγ
Punica Granatum	ρόα	tus	ριον
var.	ρόα ή ἀπύρηνος	Patientia	λάπαθος
Pyrethrum Parthe- nium	παρθένιον	Ruscus aculeatus	κεντρομυρρίνη μυάκανθος
Pyrus amygdalifor- mis	ἀχράς	Hypophyllum	δάφνη ή 'Αλεξ ανδρεία
communis	ἄπιος (1)	Ruta graveolens	πήγανον
var. Pyraster	ὄγχ <i>ν</i> η		
Malus	μηλέα	Saccharum biflorum	App. (19)
vars.	μηλέα ή γλυ-	Salix alba	ίτεα ή λευκή
	κεία, μ. ή	amplexicaulis	ίτέα ἡ μέλαινο
	έαρινή, μ. ή	cinerea fragilis	κολοιτία (2)
	b feia	spp.	έλίκη ἰτέα
Quercus Aegilops	δρῦς ἡ ἀγρία,	Salvia calycina	σφάκος
•	φηγός	- Horminum	ὄρμινον
Cerris	aiγίλωψ (1),	triloba	έλελίσφακος
	ἄσπρίς `΄΄	Sambucus nigra	άκτέος, άκτη
coccifera	πρίνος	Saponaria officinalis	στρουθίον (2),
Ilex typica	σμίλαξ (1)		στροῦθος
var.	άρία, ϊψος,	Sargassum vulgare	δρῦς (8)
agrifolia —— infectoria	φελλόδρυς	Satureia Thymbra	θύμβρα
— lanuginosa	ήμερίς (1)	Saussurea Lappa	κόστος
lanuginosa	δρῦς ἡ πλατύ-	Scandix australis	ένθρυσκον
Pseudo-Robur	φυλλος δρῦς ἡ ἁλί-	— Pecten-Veneris Schoenus Holoschoe-	σκάνδιξ
230440 230241	φλοιος, δ. ή	nus	σχοίνος ὁ ὁλό-
	εὐθύφλοιος	nigricans	σχοινος
Robur	δρῦς, δ. ἡ ήμε-		μελαγκρανίς, σχοΐνος ὁ
	ρος, ἐτυμό-		κάρπιμος
	δρυς, ημερίς	Scilla autumnalis	τίφυον
g.,,,	(2)	bifolia	υάκινθος ή άγ-
Suber	φελλός, ἵψος (?)	l	ρία

Scolopendrium	vul-
gare	
Scolymus hispar	nicus

Scorodosma foetidum Securigera Coronilla Sedum anopetalum Sempervivum tectorum

Senecio vulgaris Sesamum indicum Setaria italica

Silene venosa

Silvbum marianum Smilax aspera Smyrnium Olusatrum Solanum nigrum

Sonchus Nymani Sorbus domestica Sorghum halepense

Spartium junceum Spiraea filipendula Storax officinalis

Tamarindus indica Tamarix articulata tetrandra Taraxacum officinale Taxus baccata Tectona grandis Teucrium Polium Thapsia garganica Thymelaea hirsuta

Thymbra capitata Thymus atticus - Sibthorpii Tilia platyphyllos tomentosa.

Tordylium apulum - officinale Tragopogon porrifolius τραγοπώγων

σκολοπένδριον λειμωνία σκό-

λυμος App. (10) πελεκίνος ἐπίπετοον ἀείζωον

> ηριγέρων σήσαμον έλυμος, μέλι-204

Ήρακλεία, μήκων η 'Ηρακλεία λευκάκανθα σμίλαξ (2) **ιπποσέλινον** στρύχνος ὁ ἐδώδιμος σόγκος

ŏ'n κριθαὶ αὶ ἀγρίαι (Indian) λινόσπαρτον οἰνάνθη (1) στύραξ

App. (15) (16) μυρίκη (2) μυρίκη (1) ἀπάπη μίλος

App. (22) πόλιον θαψία κνέωρος ὁ μέλας

θύμον (1) έρπυλλος (2) έρπυλλος (1) φίλυρα φίλυρα, φ. ή θή-

λεία

καυκαλίς σέσελι

Trapa natans Tribulus terrestris Trifolium fragiferum Trigonella Foenumgraecum - graeca

Triticum dicoccum - monococcum

vulgare VATS. Tuber aestivum cibarium Typha angustata

Ulmus glabra - montana Ulva Lactuca Urginea maritima Urtica urens Usnea barbata

Valeriana Dioscoridis Veratrum album

Verbascum sinuatum

Vicia angustifolia - Ervilia – Faba - Sibthorpii Vigna sinensis Vinca herbacea

Viola odorata

Viscum album Vitex Agnus-castus Vitis vinifera corin-

var. thiaca silvestris

Zizyphus Jujuba

Lotus - Spina-Christi τρίβολος (3) τρίβολος (1) λωτός (3) βουκέρας. λις μελίλωτος.

λωτός (5) ζειά, ὀλύρα τίφή πυρός πυρός κεραύνιον ΰδνον τύφη

πτελέα δρειπτελέα βρύον σκίλλα ἀκαλύφη φάσκος

App. (25) ἐλλέβορος λευκός φλόμος ή λαινα ἀφάκη οροβος κύαμος аракос δόλιχος χαμαιδάφνη ίον το μέλαν, iwvia n mé-

λαινα ΰφεαρ ayvos, oloos ἄμπελος (1) αμπελος (2)

οἰνάνθη ἡ ἀγρία

App. (6) λωτός (4) παλίουρος ο Αίγύπτιος

II LIST OF PLANTS MENTIONED IN THE ENQUIRY UNDER POPULAR NAMES

Bog-rush μελαγκρανίς, Abele ἄκανθα (1), (3) σχοίνος Acacias κάρπιμος Alaternus φιλύκη σικύα Bottle-gourd Alder κλήθρα **ιπποσέλινον** Box πύξος Alexanders Bracken θηλύπτερις Alkanet **ἄγχουσα** Brambles βάτος, χαμαίπάνακες Allheal άμυγδαλή βατος Almond Broadleaved oak δρύς ή πλατύ-Andrachne ανδράχλη άνεμώνη φυλλος Anemones μηλέα Broom-rapes αιμόδωρον Apples App. (26) κόμαρος Arbutus ἀφάρκη Brooms λινόσπαρτον, - hybrid Arrow-poison (So- App. (27) σκορπίος (1) αμπελος ή αγmali) Bryony ρία, μήλωθκάκτος (2) Artichoke ρον Asafoetida App. (10) ράμνος, φιλύκη Buckthorns Ashes βουμελιος, με-Bugloss ονοχειλές λία Bullfist πέζις ἀσφάραγος κερκίς (2) Asparagus Bulrush τύφη Aspen ἀσφόδελος, πό-Bush-grass κάλαμος (ἐπί-Asphodel yelos) θos (2) Butcher's broom κεντρομυρρίνη, πελεκίνος Axe-weed μυάκανθος

Bachelor's buttons παρθένιον μελισσόφυλ-Balm λον

Balsam of Mecca Bamboos

Banyan Barbary nut Barley Basil Bay (sweet) Bean Bedstraw Beet Bergamot-mint

Bird-cherry Birth-wort Bladder-senna

Bindweed

Blite

Βάλσαμον κάλαμος ὁ Ίνδικός συκη η Ινδική σισυρίγχιον κριθή ὥκιμον δάφνη (1) κύαμος ἀπαρίνη τεῦτλον, σισύμβριον ἰασιώνη λακέρασος, κάρη αριστολοχία κολυτέα Βλίτον

Cabbage Calamander-wood Calamint Calavance Caltrop Caper Cardamom - Nepaul Cardoon Carnation Carob

Cassia Castor-oil plant Cat's ear Cedar, odorous - prickly

--- Syrian

ράφανος App. (23) έλένιον δόλιχος τρίβολος(1),(2) κάππαρις καρδάμωμον αμωμον κάκτος (1) διόσανθος κερωνία, συκή η Αιγυπτία κασία κρότων ύποχοιρίς θυία κέδρος (1), ὀξύ-

κεδρος

κέδρος (2)

Celandine, greater	χελιδόνιον	Dittany	δίκταμνον
lesser	ἀφία	Dock	λάπαθον τὸ ἄγ-
Celery	σέλινον		ριον
Centaury	κενταύριον	Dodder	ὀροβάγχη
Chamaeleon	χαμαιλέων	Dog-mercury	αρρενόγονον,
Chamomile, wild	ανθεμον τὸ		θηλύγονον
•	ἀφύλλανθες	Dog-rose	κυνόρροδον
Charlock	κεράϊς, ραφανίς	Dog's tooth grass	ἄγρωστις
	η αγρία	Doum-palm	κόϊξ, κουκιό-
Chaste-tree	άγνος (οἶσος)	• • • • • • • • • • • • • • • • • • • •	φορον
Cheese-flower	μαλάχη (2)	Dropwort	οινάνθη (1)
Chervil	ἔ νθρυσκον	Duckweed	ϊκμη
— wild	σκάνδιξ	Dwarf palm	φοινιξ ο χα-
Chestnuts	διοσβάλανος,		μαιρριφής
CHODONAGO	καρύα ἡ Εὐ-		P PP - 7 - 1 -
	βοϊκή	Ebony	èβéνη
Chick-pea	ερέβινθος	Edder-wort	δρακόντιον
Chicory	κιχόριον	Elder	åktéos, åktñ
Christ's thorn	παλίουρος	Elecampane	πάνακες τὸ Χει-
Cinnamon	κινάμωμον	Dicominpuno	ρώνειον
Cinquefoil	πενταπετές,	Elms	πτελέα, ὀρει-
Cinqueton	πεντάφυλλον	min	πτελέα.
Citron		Erwago	
CALION	μηλέα ή Περ-	Eryngo	ήρύγγιον
Clamatia	σική	Fanuarook	0
Clematis	άθραγένη	Fenugreek	βουκέρας, τη-
Cork-oak	ίψος (?), φελ-	Ferns	, Aug
G1 1	λός	rerns	άδίαντον, ήμιό-
Coriander	κορίαννον		νιον, θηλυ-
Cornel	θηλυκράνεια		πτερίς, πολυ-
Cornelian cherry	κράνεια		πόδιον, πτ∈-
Corn-flag	ξίφιον, ξίφος,		ρίς, τριχομα-
	φάσγανον		νές, σκολο-
Corn-thistle	ἄκανθα (2)		πένδριον
Cotoneaster	οξυάκανθος	Ferula	ναρθηκία, νάρ-
Cotton-plant	(δένδρον τὸ)		ons
	έριόφορον	Feverwort	κενταυρίς
Crack willow	έλίκη	Fig, wild	έρινεός
Cress	κάρδαμον	Figs	συκή, σ. ή
Crocus	κρόκος		Αραβική, σ
Cuckoo-pint	ãρον		ή Ίνδική,
Cucumber	σίκυος		App. (17)
squirting	σίκυος ὁ ἄγριος	Filbert	καρύα η Ήρα-
Cummin	κύμινον		κλεωτική
Currant-vine	άμπελος (2)	Firs	πεύκη, ἐλάτη
Cyclamen	κυκλάμινος	Flax	λίνον
Cypress	κυπάριττος	Frankincense-tree	λιβανωτός
		French sparrow-	σκίλλα ή Έπι-
		grass	μενίδειος
Dandelion	ἀπάπη	Fungi	ἄσχιον, μύκης,
Darnel	αϊρα	_	πέζις
Date-palm	φοίνιξ (1)		•
Dill *	άνηθον, άννη-	Galingale	κύπειρος
	705	Gall-oak	ήμερίς (1)
Distaff-thistle	ἀτρακτυλίς,	Garden nightshade	
	φόνος	_	έδωδιμος
			•

Garlies	μῶλυ, σκόρο- δον	Junipers	ἄρκευθος, θυΐα, κεδρίς, κέ-
Germander	γαμαίδους		δρος, ὀξύκε-
Gilliflower	χαμαίδρυς ίον το λευκόν		δρος
Ginger-grass	σχοίνος ὁ εὐώ-		-Fo,
Carrier Branch	δης (?)		
Gladwyn	ξίρις	Kermes-oak	πρίνος
Globe-thistle	ρύτρος		
Goat's beard	τραγοπώγων	T - harman	/ /1>
Goat willow	έλαίαγνος	Laburnum	κύτισος (1)
Gold flower	έλειόχρυσος	Larkspurs	άσταφίς, πόθος
Golden thistle	λειμωνία (2),		(1), <i>ὐάκινθος</i>
	σκόλυμος	Tools	ή σπαρτή
Gourd	κολοκύντη	Leek	πράσον (1)
Grasses	αἰγίλωψ (2),	Lentil	φακός
	θρύον, πόα	Leopard's bane	σκορπίος (2)
Groundsel	ήριγέρων	Lettuce	θρίδαξ, θριδα-
Gum arabic	ἄκανθα (7)	T 414	κίνη
	unu, ou (1)	Lilies	ήμεροκαλλές,
	i		κρίνου, κ.
Hartshorn	κορωνόπους		τὸ πορφυ-
Hart's tongue	σκολοπένδριον		ροῦν, λεί-
Hartwort	σέσελι	* 1	ριον (1)
Hawk's beard	ἀπαργία	Liquorice	γλυκεία (ρίζα)
Hawthorn	μεσπίλη ή άν-		Σκύθίκή
	θηδονοείδης	Limes	φίλυρα, φ. ή
Hazel	καρύα	T14	θήλεια
Heath	έρείκη	Litmus	φύκος (5)
Hellebores	έλλέβορος	Lucerne	(πόα) ἡ Μηδική
Hemlock	κώνειον	Lupin	θέρμος
Holly	κήλαστρος		
Holm-oaks	ἀρία,ἵψος, σμῖ-	Madder	έρευθεδανόν
	λαξ (1), φελ-	Madonna lily	κρίνον, λείριον
	λόδρυς		(1)
Honeysuckle	κλύμενον	Maiden-hair	άδίαντον, ά. τὸ
(Greek)			μέλαν
Hop-hornbeam	δστρύα, δστρυς	English	ἀδίαντον τὸ
Horehounds	πράσιον		λευκόν
Horned poppy	μήκων ή κερα-	Mallow	μαλάχη (1)
	τίτις	Mandrake	μανδραγόρας
Horseradish	ραφανίς ή άμ-		(1)
	ωρέα (?)	Mango	App. (5)
House-leek	άείζωον	Mangroves	App. (12)
Hulwort	πόλιον	white	δάφνη (6), ἐλάα
			(3), App. (14)
Irises	2 Elma	Manna-ash	μελία
111909	ίρις, ξίρις, σι-	Maples	γλείνος, ζυγία,
Ivies	συρίγχιον	_	κλινότροχος,
T A 100	ëλιξ, κιττός		σφένδαμνος
		Marestail	ίπνον
Jack-fruit	App. (3)	Marjorams	ὀρίγανον
Joint-fir	θραύπαλος	sweet	άμάρακον
Jujube	App. (6)	Marsh celery	έλειοσέλινον
Judas-tree	κερκίς (1), ση-	Marsh mallow	ἀλθαία, μαλά-
	μύδα		χη η άγρία
_			

Wantagen lile	5	1 Onione	
Martagon lily Mastich	ήμεροκαλλές σχίνος	Onions	γήθυον, γή-
Meadow saffron	εφήμερον, σπά-		τειον, κρόμ- μυον, κρο-
Medde w banton	λαξ		μυογήτειον
Medlar	μεσπίλη, μ. ή	Opium poppy	μηκων (ἡ ὀπώ-
	σατάνειος		δης), (νηπεν-
Michaelmas daisy	ἀστέρισκος,		$\theta \hat{\epsilon}_{S})$
	(τριπόλιον)	Orach	ἀδράφαξυς
Milk-thistle	λευκάκανθα	Orchis	ὄρχις
Milt-waste Millet	ὴμιόνιον	Oriental thorn	μεσπίλη ή άν»
Italian	κέγχρος	Oyster-green	θηδών
Mints	έλυμος ἡδύοσμον, μίν-	Ox-tongue	βρύον κρηπίς
322205	θη, σισύμ-	Oil tonguo	xpijii is
	βριον		
Mistletoes	ίξία (1), ΰφεαρ	Palms	κόϊξ, κουκιό-
Mock-privet	φίλυρα η ἄρρην	_	φόρον, φοινιξ
Moly	μῶλυ	Papyrus	πάπυρος
Monk's rhubarb	λάπαθος	Parsley	ὀρειοσέλινον
Mulberry Mullein	σύκάμινος	Pea Pear	πισός
Munem	φλόμος η μέ-	wild	απιος (1)
Mushroom	λαινα μύκης	Pennyroyal	άχράς, δγχνη βληχώ
Mustard, white	μάπυ	Peony	γλυκυσίδη,
Myrrh	σμύρνα	1	παιωνία.
Myrtle	μυρρίνη	Pepper	πέπερι
		Periwinkle	χαμαιδάφνη
Manatana		Pimpernel, blue	κόρχορος
Narcissus	λείριον (2),	Pines Pine-thistle	πεύκη, πίτυς
Nepaul cardamom	νάρκισσος	rine-thistie	ἄκανθα (8) ,
Nettle	ακαλύφη		ἄκανος, ἰξία (2), ἰξίνη,
— tree	λωτός (1)	ĺ	χαμαιλέων
Nightshade, deadly	μανδρανόρας		δ λευκός
,	(2)?	Plane-tree	πλάτανος
garden	στρύχνος ὁ ἐδώ-	Plantains	άρνόγλωσσον,
	διμος		θρυαλλίς, κο-
	·		ρωνόπους, κύ-
Oaks	nl (1) (1)		νωψ, ὄρτυξ, στελέφουρος
Oaks	αί γίλωψ (1), ἀρία, ἄσπρις,	Plums	κοκκυμηλέα,
	δρῦς, ἐτυμό-		πάδος, πρού-
	δρυς, ήμερίς,		μνη, σποδίας
	ίψος, πρίνος,	Polypody	πολυπόδιον
	σμίλαξ (1),	Pole-reed	δόναξ, κάλαμος
	φηγός, φελ-		δ Λακωνικός
	λόδρυς, φελ-	Demonsta	etc.
Oak-mistletoe	λός 26/- (1)	Pomegranate Pomejog	ρόα
Oak-mistiewe	ίξία (1)	Poppies Poplar, black	μήκων αϊγειρος
Oleander	βρόμος δάφνη ἡ ἀγρία.	white (abele)	λεύκη
A. CAMPINGOT	ονοθήρας	Privet	σπείραια.
Olives	έλάα (1), (2)	Puff-ball	ἄσχιον
Olive, wild	άγριέλαιος, κό-	Purslane	άνδράχνη
,	TIVOS	Purse-tassels	βολβός

Soapwort

Spanish broom Spear-grass

Sow-thistle Spike-lavender

Spikenard

Spurges

Squills
Star-flower

Star-thistle

Stonecrop Storax Sulphur-wort Sumachs

Sweet bay

Sweet flag

Sycamore

Tamarind

Tamarisks

Tare

Teak Terebinth

Thistles

- --- yellow

Spindle-tree

Sorb Southernwood

Quince	κυδώνιον,
4	στρούθιον (2)
Radish	ραφανίς
Reeds	δόναξ, κάλα-
	μος, πάπυρος
Restharrow	ονωνίς
Rib-grass	κύνωψ
Rice	ὄρυζον
Rice-wheat	ζειά, όλυρα
Rocket	εύζωμον
Rock-roses	κίσθος
Rose-campion	λυχνίς
Roses	κυνόρροδον, κυ-
	νόσβατος, ρο-
	δωνία
Rue	πήγανον
Rupture-wort	έλλεβορίνη
Rushes	μελαγκρανίς,
	σχοίνος
Safflower	κνήκος, κρόκος
	ὸ ἀκανθώδης
Saffron crocus	κρόκος, κ. ο
	εὐώδης
Sage	σφάκος
Salvia	έλελίσφακος
Savory	θύμβρα
Scammon y	σκαμμωνία
Scrub oak	δρῦς ἡ πλατύ-
a	φυλλος
Sea-bark oak	δρῦς ἡ ἀλί-
	φλοιος, δ. ή
~	εὐθύφλοιος τιθύμαλλος δ
Sea spurge	
	παράλιος
Sea-weeds	άμπελος (3),
	βρύον, δρῦς (7), (8), ἐλά-
	(1), (8), EAA-
	τη (3), φοινιξ
0-1	(4), φῦκος κοκκυμηλέα ἡ
Sebesten	
Ca Ama	Αἰγυπτία
Sedge	βούτομος
Sesame	σήσαμον Κοόμμυον τὸ
Shallot	
Oll-blumi	σχιστόν σίλφιον
Silphium	σιλφιον ἐλάτη
Silver-fir	
Silver-lim '	φίλυρα, φ. ή θήλεια
Gianoo	
Sissoo	App. (21)

σμίλαξ (2)

ἀντίρρινον

λευκόϊον (2)

Smilax

498

Snowdrop

Snapdragon

στρούθιον (2), στρούθος άβρότονον λινόσπαρτον κάλαμος ὁ πλόκιμος, κ. ò χαρακίας σόγκος ιφύον νάρδον εὐώνυμος (τετραγωνία) ἄκανθα (5), (6), ἄπιος (2), ὶπποφέως, ίσχάς, μηκώ-νιον, ῥαφανὶς ἡ ὀρεία, τιθύ-μαλλος σκίλλα, τί-φυον, ὑάκιν-θος ἡ ἀγρία βολβίνη παντάδουσα τετράλιξ επίπετρον στύραξ πευκέδανον ρούς δάφνη (1) κάλαμος ὁ εὐώ• δης συκάμινος Αίγυπτία, σ. ή Κυπρία App. (15), (16) μυρίκη ἀφάκη App. (22) τέρμινθος ăκανθα (2), (8), äkavos, ἄκορνα, ἰξία (2), ifivn, κάκτος, λειμωνία (2). λευκάκανθα,

πολυάκανθος, ρύτρος, σκόλυμος, σόγ-

κος, χάλ-

keios, χαμαι-

λέων

Thorn, oriental	μεσπίλη ἡ ἀν- θηδών	Vine, wild Violet	οἰνάνθη ἡ ἀγρία ἴον τὸ μέλαν
Thorn-apple	θρύορον, περιτ- τός, στρυχ- νος ὁ μανικός	Wall-flower Walnut	φλόγινον, φλόξ καρύα ἡ Περ-
Thyine-wood	θύον		σική
Thyme, Attic	ἔρπυλλος ὁ ἄγ ∙	Water chestnut	τρίβολος (3)
	ριος	Water chickweed	λέμνα
Cretan	θύμον (1)	Water-lily, Nile	λωτός (2)
tufted	έρπυλλος (ό	white	σίδη
	ήμερος)	yellow	μαδωνάϊς, νυμ-
Tine-tare	App. (1)		φαία
Tragacanth	τραγάκανθα	Wheats	πυρός
Traveller's joy	άθραγένη	one-seeded	τίφη
Tree-medick	κύτισος (2)	Wheat-rust	ἐρυσίβη
Tree-moss	φάσκος	Wig-tree	κοκκυγέα
Trefoil	λωτός (3)	Willows	έλαίαγνος, έλί-
Truffles	κεραύνιον, ΰδ- νου		κη, ἰτέα, κο- λοιτία (2)
Turk's cap lily	κρίνον τὸ πορ-	Willow-weed	κραταίγονος
	φυροῦν	Withy	oloos
Turkey oak	αἰγίλωψ (1), ἄσπρις	Wolf's bane	ἀκόνιτον, θηλύ- φονον, μυό-
Turnip	γογγυλίς		φονον, σκορ- πίος (3)
Valonia oak	δρῦς ἡ ἀγρία,	Wormwood	ἀψίνθιον
	φηγός	Wych-elm	ορειπτελέα
Vetch, bitter	ὄροβος		
Vine	ἄμπελος (1)	Yew	μίλος

PRINTED IN GREAT BRITAIN BY
RICHARD CLAY AND SONS, LIMITED,
BRUNSWICK STREET, STAMFORD STREET, S.E.,
AND BUNGAY, SUFFOLK.

THE LOEB CLASSICAL LIBRARY.

VOLUMES ALREADY PUBLISHED.

Latin Authors.

APULEIUS. The Golden Ass. (Metamorphoses.) Trans. by W. Adlington (1566). Revised by S. Gaselee. I Vol.

CAESAR: CIVIL WARS. Trans. by A. G. Peskett, I Vol. CATULLUS. Trans. by F. W. Cornish; TIBULLUS. Trans. by J. P. Postgate; PERVIGILIUM VENERIS. Trans. by J. W. Mackail. 1 Vol.

CICERO: DE FINIBUS. Trans. by H. Rackham. I Vol. CICERO: DE OFFICIIS. Trans. by Walter Miller. 1 Vol.

CICERO: LETTERS TO ATTICUS. Trans. by E. O. Winstedt. Vols I and II.

CONFESSIONS OF ST. AUGUSTINE. Trans. by W. Watts (1631). 2 Vols.

HORACE: ODES AND EPODES. Trans. by C. E. Bennett. T Vol.

OVID: HEROIDES AND AMORES. Trans. by Grant Showerman. 1 Vol.

OVID: METAMORPHOSES. Trans. by F. J. Miller. 2 Vols.

PETRONIUS. Trans. by M. Heseltine; SENECA: APOCO-LOCYNTOSIS. Trans. by W. H. D. Rouse. I Vol.

Trans. by Paul Nixon. Vol. I. PLAUTUS.

PLINY: LETTERS. Melmoth's Translation revised by W. M. L. Hutchinson. 2 Vols.

PROPERTIUS. Trans. by H. E. Butler. 1 Vol.

SENECA: EPISTULAE MORALES. Trans. by R. M. Gummere, Vol. I.

SENECA: TRAGEDIES. Trans. by F. J. Miller. 2 Vols. SUETONIUS. Trans. by J. C. Rolfe. 2 Vols.

TACITUS: DIALOGUS. Trans. by Sir Wm. Peterson; AGRICOLA AND GERMANIA. Trans. by Maurice Hutton, I Vol.

TERENCE. Trans. by John Sargeaunt, 2 Vols.

Greek Authors.

ACHILLES TATIUS. Trans. by S. Gaselee. I Vol. APOLLONIUS RIIODIUS. Trans. by R. C. Seaton. I Vol. THE APOSTOLIC FATHERS. Trans. by Kirsopp Lake. 2 Vols.

APPIAN'S ROMAN HISTORY. Trans. by Horace White. 4 Vols.

DAPHNIS AND CHLOE. Thornley's Translation revised by J. M. Edmonds; PARTHENIUS. Trans. by S. Gaselee. I Vol.

DIO CASSIUS: ROMAN HISTORY. Trans. by E. Cary. Vols. I, II, III, IV, and V. EURIPIDES. Trans. by A. S. Way. 4 Vols.

GALEN: ON THE NATURAL FACULTIES. Trans. by A. J. Brock. I Vol.

THE GREEK ANTHOLOGY. Trans. by W. R. Paton. Vols. I, II, III, and IV.

THE GREEK BUCOLIC POETS (THEOCRITUS, BION, MOSCHUS). Trans. by J. M. Edmonds. I Vol.

HESIOD AND THE HOMERIC HYMNS. Trans. by H. G. Evelyn White. I Vol.

JULIAN. Trans. by Wilmer Cave Wright. Vols. I and II. LUCIAN. Trans. by A. M. Harmon. Vols. I and II. MARCUS AURELIUS. Trans. by C. R. Haines. I Vol. PHILOSTRATUS: THE LIFE OF APOLLONIUS OF TYANA. Trans. by F. C. Conybeare. 2 Vols.

PINDAR. Trans. by Sir J. E. Sandys. 1 Vol.

PLATO: EUTHYPHRO, APOLOGY, CRITO, PHAEDO, PHAEDRUS. Trans. by H. N. Fowler. I Vol. PLUTARCH: THE PARALLEL LIVES. Trans. by B.

PLUTARCH: THE PARALLEL LIVES. Trans. by B. Perrin. Vols. I, II, III, and IV.

PROCOPIUS. Trans. by H. B. Dewing. Vols. I and II. QUINTUS SMYRNAEUS. Trans. by A. S. Way. I Vol. SOPHOCLES. Trans. by F. Storr. 2 Vols.

ST. JOHN DAMASCENE: BARLAAM AND IOASAPH.
Trans. by the Rev. G. R. Woodward and Harold Mattingly.
1 Vol.

STRABO: GEOGRAPHY. Trans. by Horace L. Jones. Vol. I.

THEOPHRASTUS: ENQUIRY INTO PLANTS. Trans. by Sir Arthur Hort, Bart. 2 Vols.

XENOPHON: CYROPAEDIA. Trans. by Walter Miller.

2 Vols.

DESCRIPTIVE .PROSPECTUS ON APPLICATION.

London - - WILLIAM HEINEMANN. New York - - G. P. PUTNAM'S SONS.

on



FEB 9 785

DOC 5 1992.

SPRING 1979

- 13/9

SEP 1 0 1988 Stanford University Library Stanford, California

In order that others may use this book, please return it as soon as possible, but not later than the date due.



